

LIST OF FIGURES

- Figure 2.1.** Schematic of the two-stage model proposed for general folding of α -helical bundles (top) and the concerted model for the folding of β -barrels into the lipid bilayer. 11
- Figure 2.2.** Schematic illustrating the general cell envelope of *E.coli*. The inner membrane (IM) is separated from the outer membrane (OM) by the aqueous periplasm, which contains the peptidoglycan cell wall component. Inner membrane proteins contain α -helical domains while outer membrane proteins contain β -barrel transmembrane domains. Both the IM and OM contain lipoproteins that are attached to their periplasmic sides. Adapted from Ruiz et al., 2006. 14
- Figure 2.3.** The amino acid sequence of wt-OmpA is shown. The signal sequence prior to the OmpA sequence is shown in parentheses. Native Trp (W7, W15, W57, W102, W143) are highlighted in blue. The last residue in all truncated proteins is highlighted in red. A175 is shown in green and C290 and C302 that were mutated to serines are shown in orange. 20
- Figure 2.4.** Structure of the OmpA transmembrane domain (residues 1-171), solved by X-ray crystallography (Pautsch & Schulz, 1998). The 5 native Trp are shown along with the approximate dimensions and an illustration of the C-terminal tail in the periplasmic space. The approximate location of the transmembrane region is shown with the red, dashed lines. 21

Figure 2.5. Schematic of the oriented insertion of OmpA into the lipid vesicles in refolding <i>in vitro</i> experiments. The C-terminus tail of OmpA is located on the exterior of the vesicle while the extracellular loops are on the interior of the vesicle.	22
Figure 2.6. Cross section of the OmpA transmembrane structure displaying the Trp orientations. W102 is the only Trp that faces the OmpA pore. The other 4 Trp face the exterior of the bilayer and interact with the lipid bilayer.	23
Figure 2.7. Typical DLS scan showing the diameter of DMPC vesicles. Sizes range from 20-60 nm in diameter.	33
Figure 2.8. Scheme showing the major steps of OmpA extraction from outer membranes.	34
Figure 2.9. Representative FPLC trace for purification of OmpA mutants on a 35 HiTrap QFF column. A 0-200 mM NaCl gradient was used to elute unfolded protein, which comes out as a broad peak on the trace.	35
Figure 2.10. Typical SDS-PAGE of FPLC purified OmpA is shown on the left. The right figure shows unfolded (35 kDa) and folded (30 kDa) OmpA in vesicles.	36
Figure 3.1. Structures of OG detergent and DMPC phospholipid. Illustrations of a detergent micelle and unilamellar vesicles are shown, along with approximate sizes.	48
Figure 3.2. Schematic of the setup for time-resolved measurements using	52

the femtosecond Titanium:sapphire (Ti:Sap) laser and the picosecond streak camera. Not all optical parts are illustrated. Figure not drawn to scale. Abbreviation PD = photodiode.

Figure 3.3. CD spectra of wild-type, full-length, and truncated Trp mutants unfolded in urea (top) and buffer (bottom). 66

Figure 3.4. CD spectra of wild-type, full-length, and truncated Trp mutants folded in OG micelles (top), DMPC vesicles at 30 °C (middle), and DMPC vesicles at 15 °C (bottom). 67

Figure 3.5. Overlay of CD spectra for Trp mutants at 30 °C (solid) and 15°C (dotted). Truncated mutants display lower β -sheet signal than full-length. 15 °C data were averaged over the last 30-60 min of refolding. 30 °C data were averaged over 130-150 min into folding. 68

Figure 3.6. Steady-state fluorescence of NATA in urea, phosphate buffer, OG micelles, and DMPC vesicles at 30 °C. Spectra at 15 °C have the same emission maxima at 352 nm as the 30 °C spectra. 69

Figure 3.7. Steady-state fluorescence spectra of full-length (solid lines) and truncated (dotted lines) OmpA mutants unfolded in urea at 30 °C. Spectra were corrected for vesicle-only background and normalized to protein concentration. 70

Figure 3.8. Steady-state fluorescence spectra of full-length (solid lines) and truncated (dotted lines) OmpA mutants folded in OG micelles at 30 °C. Spectra were corrected for vesicle-only background and normalized to protein concentration. 71

Figure 3.9. Steady-state fluorescence spectra of full-length (solid lines) and truncated OmpA mutants folded in DMPC vesicles at 30 °C. Spectra were corrected for vesicle-only background and normalized to protein concentration.	72
Figure 3.10. Steady-state fluorescence spectra of full-length (solid lines) and truncated (dotted lines) OmpA mutants folded in DMPC vesicles at 15 °C. Spectra were corrected for vesicle-only background and normalized to protein concentration.	73
Figure 3.11. Comparison of steady-state fluorescence spectra of full-length and truncated OmpA mutants folded in DMPC vesicles at 30 °C (solid lines) and 15 °C (dotted lines). Top panel shows full-length mutants at both temperatures. Bottom panel shows truncated mutants at both temperatures. Spectra were corrected for vesicle-only background and normalized to protein concentration.	74
Figure 3.12. Steady-state fluorescence spectra of full-length (solid lines) and truncated (dotted lines) OmpA mutants unfolded in phosphate buffer at 30 °C. Spectra were corrected for vesicle-only background and normalized to protein concentration.	75
Figure 3.13 Time-resolved fluorescence of NATA in urea, phosphate buffer, micelles, and vesicles (30 °C and 15 °C).	77
Figure 3.14. Time-resolved fluorescence of Trp mutants in urea.	79
Figure 3.15. Time-resolved fluorescence of Trp mutants in OG micelles.	81
Figure 3.16. Time-resolved fluorescence of Trp mutants in DMPC	83

vesicles, 30 °C.

Figure 3.17. Fluorescence decays of full-length (left) and truncated 85

mutants (right), mutants unfolded in urea (black), folded in OG micelles

Figure 3.18. Time-resolved fluorescence of Trp mutants in DMPC 86

vesicles, 15 °C

Figure 3.19. Steady-state anisotropy of mutants in urea. 89

Figure 3.20. Steady-state anisotropy of mutants in micelles. 90

Figure 3.21. Steady-state anisotropy of mutants in DMPC. 91

Figure 3.22. Time-resolved anisotropy of full-length (top) and truncated 93

mutants (bottom) unfolded in urea. Measurements were taken in a 50 ns time window.

Figure 3.23. Time-resolved anisotropy of full-length (top) and truncated 94

mutants (bottom) folded in OG micelles. Measurements were taken in a 50 ns time window.

Figure 3.24. Time-resolved anisotropy of full-length (top) and truncated 95

mutants (bottom) folded in DMPC vesicles at 30 °C. Measurements were taken in a 50 ns time window.

Figure 3.25 Overlay of anisotropy decays for full-length and truncated 96

mutants for each Trp position.

Figure 3.26. Time-resolved anisotropy of full-length (top) and truncated 97

mutants (bottom) folded in DMPC vesicles at 30 °C. Measurements were taken in a 5 ns time window.

Figure 4.1. Structures of DMPC, 1-palmitoyl-2-stearoyl (6-7) dibromo- <i>sn</i> -glycero-3-phosphocholine (6,7-DiBr), and 1-palmitoyl-2-stearoyl (11-12) dibromo- <i>sn</i> -glycero-3-phosphocholine (11,12-DiBr). Mixed vesicles of DMPC/6,7-DiBr and DMPC/11,12-DiBr were prepared with a 0.25 molar fraction of DMPC:DiBr lipids.	108
Figure 4.2. Steady-state fluorescence of full-length (top) and truncated (bottom) mutants in DMPC (solid lines) and 6,7-DiBr (dashed lines) vesicles	113
Figure 4.3. Steady-state fluorescence of full-length (top) and truncated (bottom) mutants in DMPC (solid lines) and 11,12-DiBr (dashed lines) vesicles	114
Figure 4.4. Fluorescence decays of full-length (top) and truncated (bottom) Trp mutants folded in DMPC vesicles (solid lines) and 6,7-DiBr vesicles (dashed lines) at 30 °C.	115
Figure 4.5. Fluorescence decays of full-length (top) and truncated (bottom) Trp mutants folded in DMPC vesicles (solid lines) and 11,12-DiBr vesicles (dashed lines) at 30 °C.	116
Figure 4.6. Fluorescence quenching rates of full-length (top) and truncated (bottom) mutants in 6,7-DiBr vesicles at 30 °C.	117
Figure 4.7. Fluorescence quenching rates of full-length (top) and truncated (bottom) mutants in 11,12-DiBr vesicles at 30 °C.	118
Figure 4.8. Overlay of the quenching rate of full-length mutants (left) and truncated mutants (right) folded in the two different DiBr lipid positions.	119

- Figure 4.9.** Overlay of the quenching rates of full-length mutants (red) and truncated mutants (blue) folded in 6,7-DiBr (left) and 11,12-DiBr (right). 120
- Figure 5.1.** Schematic of the current model of OmpA folding into the lipid bilayer. The chart indicates the average distances of the Trp residues from the center of the bilayer for each intermediate and the native structure. Adapted from Kleinschmidt, 2003. 132
- Figure 5.2.** Fluorescence spectra of W143t immediately following greater than 120 fold dilution of 8 M urea into phosphate buffer (KPi). During the 120 minute reaction, the spectra showed no systematic shifts. 144
- Figure 5.3.** Fluorescence spectra of full-length W7 mutants immediately following protein injection to initiate folding into DMPC vesicles at 30 °C. During the 120 minute folding reaction, general changes in the spectra are observed in the form of a blue-shift in emission maximum and an increase in quantum yield. 145
- Figure 5.4.** Fluorescence spectra of W7t immediately following protein injection to initiate folding into DMPC vesicles at 30 °C. During the 120 minute folding reaction, general changes in the spectra are observed in the form of a blue-shift in emission maximum and an increase in quantum yield. 146
- Figure 5.5.** Relative changes in emission maxima (top) and emission intensity (bottom) as a function of folding time. A value of “1” indicates the final emission maximum or intensity at t = 120 minutes. 147

Figure 5.6. Comparison of the rate of blue-shifting and quantum yield increase for W7 and W7t. Dotted line indicates the approximate time when blue-shift is complete while quantum yield continues to increase.	148
Figure 5.7. Comparison of the rate of blue-shifting and quantum yield increase for W15 and W15t. Dotted line indicates the approximate time when blue-shift is complete while quantum yield continues to increase.	149
Figure 5.8. Comparison of the rate of blue-shifting and quantum yield increase for W57 and W57t. Dotted line indicates the approximate time when blue-shift is complete while quantum yield continues to increase.	150
Figure 5.9. Comparison of the rate of blue-shifting and quantum yield increase for W102 and W102t. Dotted line indicates the approximate time when blue-shift is complete while quantum yield continues to increase.	151
Figure 5.10. Comparison of the rate of blue-shifting and quantum yield increase for W143 and W143t. Dotted line indicates the approximate time when blue-shift is complete while quantum yield continues to increase.	152
Figure 5.11. Fluorescence intensity of Trp mutants monitored at 330 nm immediately following initiation of folding reaction in DMPC for full-length mutants. Kinetic traces were normalized to have equivalent maximum emission intensities at 120 minutes.	153
Figure 5.12. Fluorescence intensity of Trp mutants monitored at 330 nm	154

immediately following initiation of folding reaction in DMPC for truncated mutants. Kinetic traces were normalized to have equivalent maximum emission intensities at 120 minutes.

Figure 5.13. Overlay of fluorescence intensity at 330 nm for full-length and truncated mutants at each Trp position. 155

Figure 5.14. Fluorescence spectra (top) of W7t immediately following protein injection to initiate folding into DMPC vesicles at 15 °C. The intensity at 330 nm is shown on the bottom. Spectra are immediately blue-shifted upon addition of protein to the cold vesicles. During the 60 minutes of data collection, no changes in the spectra and 330 nm values are observed. 156

Figure 5.15. Fluorescence spectra (top) of W15t immediately following protein injection to initiate folding into DMPC vesicles at 15 °C. The intensity at 330 nm is shown on the bottom. Spectra are immediately blue-shifted upon addition of protein to the cold vesicles. During the 60 minutes of data collection, no changes in the spectra and 330 nm values are observed. 157

Figure 5.16. Fluorescence spectra (top) of W102t immediately following protein injection to initiate folding into DMPC vesicles at 15 °C. The intensity at 330 nm is shown on the bottom. Spectra are immediately blue-shifted upon addition of protein to the cold vesicles. During the 60 minutes of data collection, very small changes in the spectra and 330 nm values are observed. 158

- Figure 5.17.** Fluorescence intensity measured at 330 nm for all Trp mutants immediately following protein injection to initiate adsorption onto DMPC vesicles at 15 °C. During the 60 minutes of data collection, no changes in the 330 nm values are observed. 159
- Figure 5.18.** CD spectra of full-length Trp mutants immediately following initiation of a folding reaction in 30 °C DMPC (left) and 15 °C DMPC (right). The time scales for folding are different for 30 °C and 15 °C samples. As the protein refolds, the CD spectra evolved to form β -sheet signal for folding at 30 °C. In contrast, no changes in signal were observed for refolding at 15 °C. 160
- Figure 5.19.** CD spectra of truncated Trp mutants immediately following initiation of a folding reaction in 30 °C DMPC (left) and 15 °C DMPC (right). The time scales for folding are different for 30 °C and 15 °C samples. As the protein refolds, the CD spectra evolved to form β -sheet signal for folding at 30 °C. In contrast, no changes in signal were observed for refolding at 15 °C. 161
- Figure 5.20.** Top panel shows the molar ellipticity at 206 nm for full-length mutants as the proteins fold into DMPC vesicles at 30 °C. Bottom panel shows the molar ellipticity at 206 nm for truncated mutants as the proteins fold into DMPC vesicles at 30 °C. 162
- Figure 5.21.** Changes in molar ellipticity at 206 nm for full-length mutants as the proteins fold into DMPC vesicles at 30 °C. Blue traces are truncated mutants and red traces are full-length mutants. 163

Figure 6.1. Sample potential energy landscape for lysozyme folding. 168

Adapted from Dobson et al., 1998.

Figure 6.2. Schematic depicting the relationship between protein 173

conformations and fluorescence decay kinetics. The left side shows a simplified energy landscape funnel. At the top of the funnel, an ensemble of unfolded proteins will exhibit a broad distribution of distances ($P(r)$) between the FET donors and acceptors and slow excited-decay kinetics. At the bottom, an ensemble of folded proteins should exhibit a narrow distance distribution and faster excited-state decay kinetics. The distance distribution function can be transformed using Eq. 1 to a distribution of fluorescence decay rates (k). $P(k)$ can be transformed using Eq. 4 to a fluorescence decay intensity profile ($I(t)$). Figure obtained from Julia Lyubovitsky's dissertation (Lyubovitsky, 2003).

Figure 6.3. Overlap of NATA emission with normalized Dns absorption. 182

Figure 6.4. Reaction of a general cysteine with 1,5 IEADANS to produce 184

the dansylated cysteine on the protein. Note that cysteine was drawn as a free amino acid for simplicity. In reality, these cysteines are attached to the protein.

Figure 6.5. Structure of OmpA showing the positions and distance of W7 185

and A175C. The α -carbon position of W7 ~ 16 Å from to A175C, according to the NMR structure (PDB file IG90).

Figure 6.6. Absorption spectra of time points from a side-by-side labeling 186

reaction of W7/A175C and the control protein, W7/C290S/C302S (abbreviated as W7/0C), with IAEDANS. The ratio of Dns to protein indicates that 5 hr is sufficient for labeling.

Figure 6.7. Fluorescence spectra of full-length W7/A175C-Dns 187

immediately following protein injection to initiate folding into DMPC vesicles at 30°C. Two general processes are observed (top): blue-shift in emission maxima and quantum yield increase. Relative changes in emission maxima (bottom, blue trace) and emission intensity (bottom, green and red traces) are shown as a function of folding time. Traces were normalized so that a value of “1” corresponds to the emission maximum or intensity at $t = 2$ hr.

Figure 6.8. Fluorescence spectra of truncated W7t/A175C-Dns 188

immediately following protein injection to initiate folding into DMPC vesicles at 30 °C. Two general processes are observed (top): blue-shift in emission maxima and quantum yield increase. Relative changes in emission maxima (bottom, blue trace) and emission intensity (bottom, green and red traces) are shown as a function of folding time. Traces were normalized so that a value of “1” corresponds to the emission maximum or intensity at $t = 2$ hr.

Figure 6.9. Excitation of Dns in W7/A175C-Dns with 340 nm excitation 189

does not produce a continuous rise in emission.

Figure 6.10. Fluorescence spectra of W7/A175C (unlabeled) immediately 190

following protein injection to initiate folding into DMPC vesicles at 30°C.

Relative changes in emission maxima (bottom, blue trace) and emission intensity (bottom, red trace) are shown as a function of folding time.

Traces were normalized so that a value of “1” corresponds to the emission maximum or intensity at $t = 2$ hr.

Figure 6.11. Fluorescence spectra of W7t/A175C (unlabeled) 191

immediately following protein injection to initiate folding into DMPC vesicles at 30°C. Relative changes in emission maxima (bottom, blue trace) and emission intensity (bottom, red trace) are shown as a function of folding time. Traces were normalized so that a value of “1” corresponds to the emission maximum or intensity at $t = 2$ hr.

Figure 6. 12. Energy transfer rates are shown for the different folding 192

times for W7/A175C-Dns and W7t/A175C-Dns. The energy transfer rate has decayed to a constant by ~ 14 min for W7/A175C and between 10-60 min for W7t/A175C.

Figure 6. 13. Typical data sets collected for FET kinetics. Top, left graph 193

are measurements of Trp decays as the protein folds into DMPC vesicles;

Trp decays become only slightly faster as the protein folds. Top, right

graph are measurements of Dns emission using 290 nm excitation; energy

transfer is observed as a slow rise in intensity. Bottom graph is Dns

emission from W7/A175C-Dns in urea; no energy transfer to Dns is

observed.

Figure 6.14. CD spectra of W7/A175C, W7t/A175C, and wild-type 198

OmpA. Note that the ellipticities are lower than those from Chapter 3, most likely due to inaccurate protein concentrations used to determine molar ellipticities.