

Work Cited

- Aakalu, G., Smith, W. B., Nguyen, N., Jiang, C., and Schuman, E. M. (2001). Dynamic visualization of local protein synthesis in hippocampal neurons. *Neuron* 30, 489-502.
- Ainger, K., Avossa, D., Morgan, F., Hill, S. J., Barry, C., Barbarese, E., and Carson, J. H. (1993). Transport and localization of exogenous myelin basic protein mRNA microinjected into oligodendrocytes. *J. Cell. Biol.* 123, 431-441.
- Andersen, P., Sundberg, S. H., Sveen, O., and Wigstrom, H. (1977). Specific long-lasting potentiation of synaptic transmission in hippocampal slices. *Nature* 266, 736-737.
- Bach, M. E., Barad, M., Son, H., Zhuo, M., Lu, Y. F., Shih, R., Mansuy, I., Hawkins, R. D., and Kandel, E. R. (1999). Age-related defects in spatial memory are correlated with defects in the late phase of hippocampal long-term potentiation in vitro and are attenuated by drugs that enhance the cAMP signaling pathway. *Proc. Natl. Acad. Sci. USA* 96, 5280-5285.
- Banker, G., and Goslin, K. (1990). *Culturing nerve cells*. (Cambridge, MA, MIT Press).
- Bassell, G. J., Zhang, H., Byrd, A. L., Femino, A. M., Singer, R. H., Taneja, K. L., Lifshitz, L. M., Herman, I. M., and Kosik, K. S. (1998). Sorting of beta-actin mRNA and protein to neurites and growth cones in culture. *J. Neurosci.* 18, 251-265.
- Baudier, J., Deloulme, J. C., Van Dorsselaer, A., Black, D., and Matthes, H. W. (1991). Purification and characterization of a brain-specific protein kinase c substrate, neurogranin (p17). Identification of a consensus amino acid sequence between neurogranin and neuromodulin (*GAP43*) that corresponds to the protein kinase c phosphorylation site and the calmodulin-binding domain. *J. Biol. Chem.* 266, 229-237.
- Beach, D. L., and Bloom, K. (2001). Ash1 mRNA localization in three acts. *Mol. Biol. Cell* 12, 2567-2577.
- Berleth, T., Burri, M., Thoma, G., Bopp, D., Richstein, S., Frigerio, G., Noll, M., and Nusslein-Volhard, C. (1988). The role of localization of bicoid rna in organizing the anterior pattern of the *Drosophila* embryo. *EMBO J.* 7, 1749-1756.
- Bertrand, E., Chartrand, P., Schaefer, M., Shenoy, S. M., Singer, R. H., and Long, R. M. (1998). Localization of ash1 mRNA particles in living yeast. *Mol. Cell* 2, 437-445.
- Bito, H., Deisseroth, K., and Tsien, R. W. (1996). Creb phosphorylation and dephosphorylation: A Ca^{2+} - and stimulus duration-dependent switch for hippocampal gene expression. *Cell* 87, 1203-1214.

- Bliss, T. V. P., and Collingridge, G. L. (1993). A synaptic model of memory: Long-term potentiation in the hippocampus. *Nature* 361, 31-39.
- Bliss, T. V. P., and Gardner-Medwin, A. R. (1973). Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J. Physiol. (London)* 232, 357-374.
- Bonhoeffer, T., Staiger, V., and Aertsen, A. (1989). Synaptic plasticity in rat hippocampal slice cultures: Local "hebbian" conjunction of pre- and postsynaptic stimulation leads to distributed synaptic enhancement. *Proc. Natl. Acad. Sci. USA* 86, 8113-8117.
- Bouron, A., and Reuter, H. (1999). The D1 dopamine receptor agonist SKF-38393 stimulates the release of glutamate in the hippocampus. *Neuroscience* 94, 1063-1070.
- Bradshaw, K. D., Emptage, N. J., and Bliss, T. V. (2003). A role for dendritic protein synthesis in hippocampal late ltp. *Eur. J. Neurosci.* 18, 3150-3152.
- Brittis PA, L. Q., Flanagan JG (2002). Axonal protein synthesis provides a mechanism for localized regulation at an intermediate target. *Cell* 110, 223-235.
- Brown, A. (2003). Axonal transport of membranous and nonmembranous cargoes: A unified perspective. *J. Cell. Biol.* 160, 817-821.
- Brown, E. J., and Schreiber, S. L. (1996). A signaling pathway to translational control. *Cell* 86, 517-520.
- Burgin, K. E., Waxham, M. N., Rickling, S., Westgate, S. A., Mobley, W. C., and Kelly, P. T. (1990). *In situ* hybridization histochemistry of Ca/calmodulin dependent protein kinase in developing rat brain. *J. Neurosci.* 10, 1788-1789.
- Cao, Q., and Richter, J. D. (2002). Dissolution of the maskin-eif4e complex by cytoplasmic polyadenylation and poly(A)-binding protein controls Cyclin b1 mRNA translation and oocyte maturation. *EMBO J.* 21, 3852-3862.
- Casadio, A., Martin, K. C., Giustetto, M., Zhu, H., Chen, M., Bartsch, D., Bailey, C. H., and Kandel, E. R. (1999). A transient, neuron-wide form of CREB-mediated long-term facilitation can be stabilized at specific synapses by local protein synthesis. *Cell* 99, 221-237.
- Chao, S. Z., Ariano, M. A., Peterson, D. A., and Wolf, M. E. (2002). D1 dopamine receptor stimulation increases GluR1 surface expression in nucleus accumbens neurons. *J. Neurochem.* 83, 704-712.
- Chetkovich, D. M., and Sweatt, J. D. (1993). NMDA receptor activation increases cyclic AMP in area CA1 of the hippocampus via calcium/calmodulin stimulation of adenylyl cyclase. *J. Neurochem.* 61, 1933-1942.

- Collingridge, G. L., Kehl, S. J., and McLennan, H. (1983). Excitatory amino acids in synaptic transmission in the schaffer collateral-commissural pathway of the rat hippocampus. *J. Physiol. (London)* 334, 33-46.
- Crino, P. B., and Eberwine, J. (1996). Molecular characterization of the dendritic growth cone: Regulated mRNA transport and local protein synthesis. *Neuron* 17, 1173-1187.
- Cubitt, A. B., Heim, R., Adams, S. R., Boyd, A., and Gross, L. A. (1995). Understanding, improving and using green fluorescent proteins. *Trends Biochem. Sci.* 20, 448-455.
- Darnell, J. C., Jensen, K. B., Jin, P., Brown, V., Warren, S. T., and Darnell, R. B. (2001). Fragile X mental retardation protein targets G quartet mRNAs important for neuronal function. *Cell* 107, 489-499.
- Dash, P. K., Moore, A. N., and Dixon, C. E. (1995). Spatial memory deficits, increased phosphorylation of the transcription factor CREB, and induction of the AP-1 complex following experimental brain injury. *J. Neurosci.* 15, 2030-2039.
- Deisseroth, K., Bito, H., and Tsien, R. W. (1996). Signalling from synapse to nucleus: Postsynaptic CREB phosphorylation during multiple forms of hippocampal synaptic plasticity. *Neuron* 16, 89-101.
- DesGroseillers, L., Kuhl, D., Richter, D., and Kindler, S. (2001). Two rat brain Staufen isoforms differentially bind RNA. *J. Neurochem.* 76: 155-165.
- Ding, D., Parkhurst, S. M., and Lipshitz, H. D. (1993). Different genetic requirements for anterior RNA localization revealed by the distribution of adducin-like transcripts during *Drosophila* oogenesis. *Proc. Natl. Acad. Sci. USA* 90, 2512-2516.
- Engert, F., and Bonhoeffer, T. (1997). Synapse specificity of long-term potentiation breaks down at short distances. *Nature* 388, 279-284.
- Feig, S., and Lipton, P. (1993). Pairing the cholinergic agonist carbachol with patterned schaffer collateral stimulation initiates protein synthesis in hippocampal CA1 pyramidal cell dendrites via a muscarinic, NMDA-dependent mechanism. *J. Neurosci.* 13, 1010-1021.
- Ferrandon, D., Elphick, L., Nusslein-Volhard, C., and St Johnston, D. (1994). Staufen protein associates with the 3'UTR of bicoid mRNA to form particles that move in a microtubule-dependent manner. *Cell* 79, 1221-1232.
- Frey, D., Laux, T., Xu, L., Schneider, C., and Caroni, P. (2000). Shared and unique roles of cap23 and *GAP43* in actin regulation, neurite outgrowth, and anatomical plasticity. *J. Cell Biol.* 149, 1443-1454.

- Frey, U., Huang, Y. Y., and Kandel, E. R. (1993). Effects of cAMP simulate a late stage of LTP in hippocampal CA1 neurons. *Science* 260, 1661-1664.
- Frey, U., Krug, M., Reymann, K. G., and Matthies, H. (1988). Anisomycin, an inhibitor of protein synthesis, blocks late phases of LTP phenomena in the hippocampal CA region in vitro. *Brain Res.* 452, 57-65.
- Frey, U., Matthies, H., and Reymann, K. G. (1991). The effect of dopaminergic D1 receptor blockade during tetanization on the expression of long-term potentiation in the rat CA1 region in vitro. *Neurosci. Lett.* 129, 111-114.
- Frey, U., and Morris, R. G. (1997). Synaptic tagging and long-term potentiation. *Nature* 385, 533-536.
- Frey, U., Schroeder, H., and Matthies, H. (1990). Dopaminergic antagonists prevent long-term maintenance of posttetanic LTP in the CA1 region of rat hippocampal slices. *Brain Res.* 522, 69-75.
- Gardiol, A., Racca, C., and Triller, A. (1999). Dendritic and postsynaptic protein synthetic machinery. *J. Neurosci.* 19, 168-179.
- Gazzaley, A. H., Benson, D. L., Huntley, G. W., and Morrison, J. H. (1997). Differential subcellular regulation of NMDAR1 protein and mRNA in dendrites of dentate gyrus granule cells after perforant path transection. *J. Neurosci.* 17, 2006-2017.
- Greenough, W. T., Klintsova, A. Y., Irwin, S. A., Galvez, R., Bates, K. E., and Weiler, I. J. (2001). Synaptic regulation of protein synthesis and the Fragile X protein. *Proc. Natl. Acad. Sci. USA* 98, 7101-7106.
- Hebb, D. O. (1949). *The Organization of Behavior*, John Wiley & Sons.
- Huang, Y. S., Jung, M. Y., Sarkissian, M., and Richter, J. D. (2002). N-methyl-d-aspartate receptor signaling results in aurora kinase- catalyzed CPEB phosphorylation and alpha CamKII mRNA polyadenylation at synapses. *EMBO J.* 21, 2139-2148.
- Huang, Y. Y., and Kandel, E. R. (1995). D1/D5 receptor agonists induce a protein synthesis-dependent late potentiation in the CA1 region of the hippocampus. *Proc. Natl. Acad. Sci. USA* 92, 2446-2450.
- Huber, K. M., Kayser, M. S., and Bear, M. F. (2000). Role for rapid dendritic protein synthesis in hippocampal mGluR-dependent long-term depression. *Science* 288, 1254-1256.
- Isaac, J. T., Nicoll, R. A., and Malenka, R. C. (1995). Evidence for silent synapses: Implications for the expression of LTP. *Neuron* 15, 427-434.

- Jefferies, H. B., Reinhard, C., Kozma, S. C., and Thomas, G. (1994). Rapamycin selectively represses translation of the "polypyrimidine tract" mRNA family. *Proc. Natl. Acad. Sci. USA* 91, 4441-4445.
- Ju, W., Morishita, W., Tsui, J., Gaietta, G., Deerinck, T. J., Adams, S. R., Garner, C. C., Tsien, R. Y., Ellisman, M. H., and Malenka, R. C. (2004). Activity-dependent regulation of dendritic synthesis and trafficking of AMPA receptors. *Nat. Neurosci.* 7, 244-253.
- Kacharmina, J. E., Job, C., Crino, P., and Eberwine, J. (2000). Stimulation of glutamate receptor protein synthesis and membrane insertion within isolated neuronal dendrites. *Proc. Natl. Acad. Sci. USA* 97, 11545-11550.
- Kang, H., and Schuman, E. M. (1995a). Long-lasting neurotrophin-induced enhancement of synaptic transmission in the adult hippocampus. *Science* 267, 1658-1662.
- Kang, H., and Schuman, E. M. (1996). A requirement for local protein synthesis in neurotrophin-induced synaptic plasticity. *Science* 273, 1402-1406.
- Kang, H., Welcher, A. A., Shelton, D., and Schuman, E. M. (1997). Neurotrophins and time: Different roles for TRK-B signaling in hippocampal long-term potentiation. *Neuron* 19, 653-664.
- Kiebler, M. A., Hemraj, I., Verkade, P., Kohrmann, M., Fortes, P., Marion, R. M., Ortin, J., and Dotti, C. G. (1999). The mammalian Staufen protein localizes to the somatodendritic domain of cultured hippocampal neurons: Implications for its involvement in mRNA transport. *J. Neurosci.* 19, 288-297.
- Kiss, J. (1977). Synthesis and transport of newly formed proteins in dendrites of rat hippocampal pyramid cells. An electron microscope autoradiographic study. *Brain Res.* 124, 237-250.
- Knowles, R. B., Sabry, J. H., Martone, M. E., Deerinck, T. J., Ellisman, M. H., Bassell, G. J., and Kosik, K. S. (1996). Translocation of RNA granules in living neurons. *J. Neurosci.* 16, 7812-7820.
- Koenig, E., and Giuditta, A. (1999). Protein-synthesizing machinery in the axon compartment. *Neuroscience* 89, 5-15.
- Koenig, E., Martin, R., Titmus, M., and Sotelo-Silveira, J. R. (2000). Cryptic peripheral ribosomal domains distributed intermittently along mammalian myelinated axons. *J. Neurosci.* 20, 8390-8400.

- Kozubek, M., and Matula, P. (2000). An efficient algorithm for measurement and correction of chromatic aberrations in fluorescence microscopy. *J. Microsc.* 200 Pt 3, 206-217.
- Krichevsky, A. M., and Kosik, K. S. (2001). Neuronal RNA granules: A link between RNA localization and stimulation-dependent translation. *Neuron* 32, 683-696.
- Kullmann, D. M., and Siegelbaum, S. A. (1995). The site of expression of NMDA receptor-dependent LTP: New fuel for an old fire. *Neuron* 15, 997-1002.
- Lee, F. J., Xue, S., Pei, L., Vukusic, B., Chery, N., Wang, Y., Wang, Y. T., Niznik, H. B., Yu, X. M., and Liu, F. (2002). Dual regulation of NMDA receptor functions by direct protein-protein interactions with the dopamine D1 receptor. *Cell* 111, 219-230.
- Lerner, E. A., Lerner, M. R., Janeway, C. A., and Steitz, J. A. (1981). Monoclonal antibodies to nucleic acid-containing cellular constituents: Probes for molecular biology and autoimmune disease. *Proc. Natl. Acad. Sci. USA* 78, 2737-2741.
- Leung, L. S., and Shen, B. (1995). Long-term potentiation at the apical and basal dendritic synapses of CA1 after local stimulation in behaving rats. *J. Neurophysiol.* 73, 1938-1946.
- Levine, E. S., Dreyfus, C. F., Black, I. B., and Plummer, M. R. (1995a). Brain-derived neurotrophic factor rapidly enhances synaptic transmission in hippocampal neurons via postsynaptic tyrosine kinase receptors. *Proc. Natl. Acad. Sci. USA* 92, 8074-8078.
- Levine, E. S., Dreyfus, C. F., Black, I. B., and Plummer, M. R. (1995b). Differential effects of NGF and BDNF on voltage-gated calcium currents in embryonic basal forebrain neurons. *J. Neurosci.* 15(4), 3084-3091.
- Li, S., Cullen, W. K., Anwyl, R., and Rowan, M. J. (2003). Dopamine-dependent facilitation of LTP induction in hippocampal CA1 by exposure to spatial novelty. *Nat. Neurosci.* 6, 526-531.
- Li, Y. X., Xu, Y., Ju, D., Lester, H. A., Davidson, N., and Schuman, E. M. (1998). Expression of a dominant negative TRK B receptor, T1, reveals a requirement for presynaptic signaling in BDNF-induced synaptic potentiation in cultured hippocampal neurons. *Proc. Natl. Acad. Sci. USA* 95, 10884-10889.
- Li, Y. X., Zhang, Y., Lester, H.A., Schuman, E.M., Davidson, N. (1998). Enhancement of excitatory neurotransmitter release induced by BDNF in cultured hippocampal neurons. *J. Neurosci.* 18, 10231-10240.
- Liao, D., Zhang, X., O'Brien, R., Ehlers, M. D., and Huganir, R. L. (1999). Regulation of morphological postsynaptic silent synapses in developing hippocampal neurons. *Nat. Neurosci.* 2, 37-43.

- Liao, S., Hessler, N. A., and Malinow, R. (1995). Activation of postsynaptically silent synapses during pairing-induced ltp in cal region of hippocampal slice. *Nature* 375, 400-404.
- Lieberfarb, M. E., Chu, T., Wreden, C., Theurkauf, W., Gergen, J. P., and Strickland, S. (1996). Mutations that perturb poly(A)-dependent maternal mRNA activation block the initiation of development. *Development* 122, 579-588.
- Liu, F., Wan, Q., Pristupa, Z. B., Yu, X. M., Wang, Y. T., and Niznik, H. B. (2000). Direct protein-protein coupling enables cross-talk between dopamine D5 and gamma-aminobutyric acid A receptors. *Nature* 403, 274-280.
- Lyford, G. L., Yamagata, K., Kaufmann, W. E., Barnes, C. A., Sanders, L. K., Copeland, N. G., Gilbert, D. J., Jenkins, N. A., Lanahan, A. A., and Worley, P. F. (1995). Arc, a growth factor and activity-regulated gene, encodes a novel cytoskeleton-associated protein that is enriched in neuronal dendrites. *Neuron* 14, 433-445.
- Malenka, R. C. (2003). Synaptic plasticity and AMPA receptor trafficking. *Ann. N. Y. Acad. Sci.* 1003, 1-11.
- Malgaroli, A., and Tsien, R. W. (1992). Glutamate-induced long-term potentiation of the frequency of miniature synaptic currents in cultured hippocampal neurons. *Nature* 357, 134-139.
- Mangiavacchi S, W. M. (2004). D1 dopamine receptor stimulation increases the rate of AMPA receptor insertion onto the surface of cultured nucleus accumbens neurons through a pathway dependent on protein kinase A. *J. Neurochem.* 88, 1261-1271.
- Martin, K. C., Barad, M., and Kandel, E. R. (2000). Local protein synthesis and its role in synapse-specific plasticity. *Curr. Op. Neurobiol.* 10, 587-592.
- Martin, K. C., Casadio, A., Zhu, H., E, Y., Rose, J. C., Chen, M., Bailey, C. H., and Kandel, E. R. (1997). Synapse-specific, long-term facilitation of aplysia sensory to motor synapses: A function for local protein synthesis in memory storage. *Cell* 91, 927-938.
- Martin, K. C., and Kosik, K. S. (2002). Synaptic tagging - who's it? *Nat. Rev. Neurosci.* 3, 813-820.
- Matthies, H., Becker, A., Schroeder, H., Kraus, J., Hollt, V., and Krug, M. (1997). Dopamine D1-deficient mutant mice do not express the late phase of hippocampal long-term potentiation. *Neuroreport* 8, 3533-3535.
- Mayer, M. L., Westbrook, G. L., and Guthrie, P. B. (1984). Voltage-dependent block by Mg^{2+} of NMDA responses in spinal cord neurones. *Nature* 309, 261-263.

- Mayford, M., Baranes, D., Podsypanina, K., and Kandel, E. R. (1996). The 3'-untranslated region of CamKII-alpha is a cis-acting signal for the localization and translation of mRNA in dendrites. *Proc. Natl. Acad. Sci. USA* 93, 13250-13255.
- McAllister, A. K., Katz, L. C., and Lo, D. C. (1999). Neurotrophins and synaptic plasticity. *Ann. Rev. Neurosci.* 22, 295-318.
- Miesenbock, G., De Angelis, D. A., and Rothman, J. E. (1998). Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins. *Nature* 394, 192-195.
- Miller, S., Yasuda, M., Coats, J. M., Jones, Y., Martone, M. E., and Mayford, M. (2002). Disruption of dendritic translation of CamKII-alpha impairs stabilization of synaptic plasticity and memory consolidation. *Neuron* 36, 507-519.
- Miyashiro, K., Dichter, M., and Eberwine, J. (1994). On the nature and differential distribution of mRNAs in hippocampal neurites: Implications for neuronal functioning. *Proc. Natl. Acad. Sci. USA* 91, 10800-10804.
- Mori, Y., Imaizumi, K., Katayama, T., Yoneda, T., and Tohyama, M. (2000). Two cis-acting elements in the 3' untranslated region of the alpha-CamKII regulate its dendritic targeting. *Nat. Neurosci.* 3, 1079-1084.
- Nathans, D. (1964). Puromycin inhibition of protein synthesis: Incorporation of puromycin into peptide chains. *Proc. Natl. Acad. Sci. USA* 51, 585-592.
- Nayak, A., Zastrow, D. J., Lickteig, R., Zanister, N. R., and Browning, M. D. (1998). Maintenance of late-phase LTP is accompanied by PKA-dependent increase in AMPA receptor synthesis. *Nature* 394, 690-683.
- Nguyen, P. V., Abel, T., and Kandel, E. R. (1994). Requirement of a critical period of transcription for induction of a late phase of LTP. *Science* 265, 1104-1107.
- Nishiyama, M., Hong, K., Mikoshiba, K., Poo, M. M., and Kato, K. (2000). Calcium stores regulate the polarity and input specificity of synaptic modification. *Nature* 408, 584-588.
- Ostroff, L. E., Fiala, J. C., Allwardt, B., and Harris, K. M. (2002). Polyribosomes redistribute from dendritic shafts into spines with enlarged synapses during LTP in developing rat hippocampal slices. *Neuron* 35, 535-545.
- Otani, S., Marshall, C. J., Tate, W. P., Goddard, G. V., and Abraham, W. C. (1989). Maintenance of long-term potentiation in rat dentate gyrus requires protein synthesis but not messenger RNA synthesis immediately post-tetaniization. *Neuroscience* 28, 519-526.

- Otmakhova, N. A., and Lisman, J. E. (1996). D1/D5 dopamine receptor activation increases the magnitude of early long-term potentiation at CA1 hippocampal synapses. *J. Neurosci.* 16, 7478-7486.
- Otsu, N. (1979). A threshold selection method from gray-level histograms. *IEEE Trans. Sys., Man, and Cyb.* 9, 62-66.
- Ouyang, Y., Kantor, D. B., Harris, K. M., Schuman, E. M., and Kennedy, M. B. (1997). Visualization of the distribution of autophosphorylated calcium/calmodulin-dependent protein kinase II after tetanic stimulation in the CA1 area of the hippocampus. *J. Neurosci.* 17, 5416-5427.
- Ouyang, Y., Rosenstein, A., Kreiman, G., Schuman, E. M., and Kennedy, M. B. (1999). Tetanic stimulation leads to increased accumulation of ca(2+)/calmodulin-dependent protein kinase II via dendritic protein synthesis in hippocampal neurons. *J. Neurosci.* 19, 7823-7833.
- Patrick, G. N., Bingol, B., Weld, H. A., and Schuman, E. M. (2003). Ubiquitin-mediated proteasome activity is required for agonist-induced endocytosis of GluRs. *Curr. Biol.* 13, 2073-2081.
- Patrick, G. N., Zukerberg, L., Nikolic, M., de la Monte, S., Dikkes, P., and Tsai, L. H. (1999). Conversion of p35 to p25 deregulates cdk5 activity and promotes neurodegeneration. *Nature* 402, 615-622.
- Pei, L., Lee, F. J., Moszczynska, A., Vukusic, B., and Liu, F. (2004). Regulation of dopamine D1 receptor function by physical interaction with the NMDA receptors. *J. Neurosci.* 24, 1149-1158.
- Petralia, R. S., Esteban, J. A., Wang, Y. X., Partridge, J. G., Zhao, H. M., Wenthold, R. J., and Malinow, R. (1999). Selective acquisition of receptors over postnatal development suggests a molecular basis for silent synapses. *Nat. Neurosci.* 2, 31-36.
- Pierce, J. P., Mayer, T., and McCarthy, J. B. (2001). Evidence for a satellite secretory pathway in neuronal dendritic spines. *Curr. Biol.* 11, 351-355.
- Pinkstaff, J. K., Chappell, S. A., Mauro, V. P., Edelman, G. M., and Krushel, L. A. (2001). Internal initiation of translation of five dendritically localized neuronal mRNAs. *Proc. Natl. Acad. Sci. USA* 98, 2770-2775.
- Pittenger, C., Huang, Y. Y., Paletzki, R. F., Bourtchouladze, R., Scanlin, H., Vronskaya, S., and Kandel, E. R. (2002). Reversible inhibition of CREB/ATF transcription factors in region CA1 of the dorsal hippocampus disrupts hippocampus-dependent spatial memory. *Neuron* 34, 447-462.

- Rao, A., and Steward, O. (1991). Evidence that protein constituents of postsynaptic membrane are locally synthesized: Analysis of proteins synthesized within synaptosomes. *J. Neurosci.* 11, 2881-2895.
- Reymann, K. G., Malisch, R., Schulzeck, K., Brodemann, R., Ott, T., and Matthies, H. (1985). The duration of long-term potentiation in the CA1 region of the hippocampal slice preparation. *Brain. Res. Bull.* 15, 249-255.
- Richmond, S. A., Irving, A. J., Molnar, E., McIlhinney, R. A., Michelangeli, F., Henley, J. M., and Collingridge, G. L. (1996). Localization of the glutamate receptor subunit GluR1 on the surface of living and within cultured hippocampal neurons. *Neuroscience* 75, 69-82.
- Rook, M. S., Lu, M., and Kosik, K. S. (2000). CamkII-alpha 3' untranslated region-directed mRNA translocation in living neurons: Visualization by GFP linkage. *J. Neurosci.* 20, 6385-6393.
- Scheetz, A. J., Nairn, A. C., and Constantine-Paton, M. (2000). NMDA receptor-mediated control of protein synthesis at developing synapses. *Nat. Neurosci.* 3, 211-216.
- Schuman, E. M. (1999a). mRNA trafficking and local protein synthesis at the synapse. *Neuron* 23, 645-648.
- Schuman, E. M. (1999b). Neurotrophin regulation of synaptic transmission. *Curr. Opin. Neurobiol.* 9, 105-109.
- Schuman, E. M., and Madison, D. V. (1994). Locally distributed synaptic potentiation in the hippocampus. *Science* 263, 532-536.
- Shi, S., Hayashi, Y., Esteban, J. A., and Malinow, R. (2001). Subunit-specific rules governing AMPA receptor trafficking to synapses in hippocampal pyramidal neurons. *Cell* 105, 331-343.
- Shi, S. H., Hayashi, Y., Petralia, R. S., Zaman, S. H., Wenthold, R. J., Svoboda, K., and Malinow, R. (1999). Rapid spine delivery and redistribution of AMPA receptors after synaptic NMDA receptor activation. *Science* 284, 1811-1816.
- Srinivasan, Y., Guzikowski, A. P., Haugland, R. P., and Angelides, K. J. (1990). Distribution and lateral mobility of glycine receptors on cultured spinal cord neurons. *J. Neurosci.* 10, 985-995.
- St. Johnston, D., Beuchle, D., and Nusslein-Volhard, C. (1991). Staufen, a gene required to localize maternal RNAs in the *Drosophila* egg. *Cell* 66, 51-63.
- Stanton, P. K., and Sarvey, J. M. (1984). Blockade of long-term potentiation in rat hippocampal CA1 region by inhibitors of protein synthesis. *J. Neurosci.* 4, 3080-3084.

- Starck S. R., and Roberts R. W. (2002). Puromycin oligonucleotides reveal steric restrictions for ribosome entry and multiple modes of translation inhibition. *RNA* 8, 890-903.
- Staubli, U., and Lynch, G. (1987). Stable hippocampal long-term potentiation elicited by 'theta' pattern stimulation. *Brain Res.* 435, 227-234.
- Stebbins-Boaz, B., Cao, Q., de Moor, C. H., Mendez, R., and Richter, J. D. (1999). Maskin is a CPEB-associated factor that transiently interacts with eIF-4e. *Mol. Cell* 4, 1017-1027.
- Stevens, C. F. (1998). A million dollar question: Does LTP = memory? *Neuron* 20, 1-2.
- Steward, O. (1997). mRNA localization in neurons: A multipurpose mechanism? *Neuron* 18, 9-12.
- Steward, O., and Levy, W. B. (1982). Preferential localization of polyribosomes under the base of dendritic spines in granule cells of the dentate gyrus. *J. Neurosci.* 2, 284-291.
- Steward, O., and Reeves, T. M. (1988). Protein-synthesis machinery beneath postsynaptic sites on CNS neurons: Association between polyribosomes and other organelles at the synaptic site. *J. Neurosci.* 8, 176-184.
- Steward, O., and Schuman, E. M. (2001). Protein synthesis at synaptic sites on dendrites. *Ann. Rev. Neurosci.* 24, 299-325.
- Steward, O., and Schuman, E. M. (2003). Compartmentalized synthesis and degradation of proteins in neurons. *Neuron* 40, 347-359.
- Steward, O., Wallace, C. S., Lyford, G. L., and Worley, P. F. (1998). Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites. *Neuron* 21, 741-751.
- Swanson-Park, J. L., Coussens, C. M., Mason-Parker, S. E., Raymond, C. R., Hargreaves, E. L., Dragunow, M., Cohen, A. S., and Abraham, W. C. (1999). A double dissociation within the hippocampus of dopamine D1/D5 receptor and beta-adrenergic receptor contributions to the persistence of long-term potentiation. *Neuroscience* 92, 485-497.
- Tang, S. J., Meulemans, D., Vasquez, L., Colaco, N., and Schuman, E. M. (2001). A role for Staufen in the delivery of RNA to neuronal dendrites. *Neuron* 8, 463-475.
- Tang, S. J., Reis, G., Kang, H., Gingras, A. C., Sonenberg, N., and Schuman, E. M. (2002). A rapamycin-sensitive signaling pathway contributes to long-term synaptic plasticity. *Proc. Natl. Acad. Sci. USA* 99, 467-472.

- Tao, H. W., Zhang, L. I., Engert, F., and Poo, M. (2001). Emergence of input specificity of LTP during development of retinotectal connections in vivo. *Neuron* 31, 569-580.
- Terada, N., Patel, H. R., Takase, K., Kohno, K., Nairn, A. C., and Gelfand, E. W. (1994). Rapamycin selectively inhibits translation of mRNAs encoding elongation factors and ribosomal proteins. *Proc. Natl. Acad. Sci. USA* 91, 11477-11481.
- Tongiorgi, E., Righi, M., and Cattaneo, A. (1997). Activity-dependent dendritic targeting of BDNF and TRKB mRNAs in hippocampal neurons. *J. Neurosci.* 17, 9492-9505.
- Torre, E. R., and Steward, O. (1996). Protein synthesis within dendrites: Glycosylation of newly synthesized proteins in dendrites of hippocampal neurons in culture. *J. Neurosci.* 16, 5967-5978.
- Torre, E. R., and Steward, O. (1992). Demonstration of local protein synthesis within dendrites using a new cell culture system that permits the isolation of living axons and dendrites from their cell bodies. *J. Neurosci.* 12, 762-772.
- Twiss, J. L., Smith, D. S., Chang, B., and Shooter, E. M. (2000). Translational control of ribosomal protein l4 mRNA is required for rapid neurite regeneration. *Neurobiol. Dis.* 7, 416-428.
- Wallace, C. S., Lyford, G. L., Worley, P. F., and Steward, O. (1998). Differential intracellular sorting of immediate early gene mRNAs depends on signals in the mRNA sequence. *J. Neurosci.* 18, 26-35.
- Weiler, I. J., and Greenough, W. T. (1991). Potassium ion stimulation triggers protein translation in synaptoneurosomal polyribosomes. *Mol. Cell. Neurosci.* 2, 305-314.
- Weiler, I. J., and Greenough, W. T. (1993). Metabotropic glutamate receptors trigger postsynaptic protein synthesis. *Proc. Natl. Acad. Sci. USA* 90, 7168-7171.
- Weiler, I. J., Irwin, S. A., Klintsova, A. Y., Spencer, C. M., Brazelton, A. D., Miyashiro, K., Comery, T. A., Patel, B., Eberwine, J., and Greenough, W. T. (1997). Fragile X mental retardation protein is translated near synapses in response to neurotransmitter activation. *Proc. Natl. Acad. Sci. USA* 94, 5395-9400.
- Wells, D. G., Richter, J. D., and Fallon, J. R. (2000). Molecular mechanisms for activity-regulated protein synthesis in the synapto-dendritic compartment. *Curr. Opin. Neurobiol.* 10, 132-137.
- Wey, C. L., Cone R.A., and A., E. M. (1981). Lateral diffusion of rhodopsin in photoreceptor cells measured by fluorescence photobleaching and recovery. *Biophys. J.* 33, 225-232.

Wickham, L., Duchaine, T., Luo, M., Nabi, I. R., and DesGroseillers, L. (1999). Mammalian Staufen is a double-stranded RNA- and tubulin-binding protein which localizes to the rough endoplasmic reticulum. *Mol. Cell. Biol.* 19, 2220-2230.

Wu, L., Wells, D., Tay, J., Mendis, D., Abbott, M. A., Barnitt, A., Quinlan, E., Heynen, A., Fallon, J. R., and Richter, J. D. (1998). CPEB-mediated cytoplasmic polyadenylation and the regulation of experience-dependent translation of alpha-CamKII mRNA at synapses. *Neuron* 21, 1129-1139.

Zheng, J. Q., Kelly, T. K., Chang, B., Ryazantsev, S., Rajasekaran, A. K., Martin, K. C., and Twiss, J. L. (2001). A functional role for intra-axonal protein synthesis during axonal regeneration from adult sensory neurons. *J. Neurosci.* 21, 9291-9303.