Chapter 3

Modulating Hypoxia-Inducible Transcription by Disrupting the HIF-1-DNA Interface

Reproduced with permission from (Nickols, N.G., Jacobs, C.S., Farkas, M.E., & Dervan, P.B. (2007) *ACS Chem Biol* **2**(8), 561–571.) Copyright [2007] American Chemical Society.

3.1 Abstract

Transcription mediated by hypoxia inducible factor (HIF-1) contributes to tumor angiogenesis and metastasis but is also involved in activation of cell-death pathways and normal physiological processes. Given the complexity of HIF-1 signaling, it could be advantageous to target a subset of HIF-1 effectors rather than the entire pathway. We compare the genome-wide effects of three molecules that each interfere with the HIF-1-DNA interaction: a polyamide targeted to the hypoxia response element (HRE), siRNA targeted to HIF-1 α , and echinomycin, a DNA-binding natural product with a similar but less specific sequence preference than the polyamide. The polyamide affects a subset of hypoxia-induced genes consistent with its binding site preferences. For comparison, HIF-1 α siRNA and echinomycin each affect the expression of nearly every gene induced by hypoxia. Remarkably, the total number of genes affected by either polyamide or HIF-1 α siRNA over a range of thresholds is comparable. The data show that polyamides can be used to affect a subset of a pathway regulated by a transcription factor. In addition, this study offers a unique comparison of three complementary approaches towards exogenous control of endogenous gene expression.



3.2 Introduction

Exogenous chemical and biological methods to control directly expression of selected endogenous genes could have broad implications for human medicine. Towards this goal, a number of technological approaches are currently being investigated. Polydactyl zinc finger proteins are a programmable class of DNA binding proteins capable of sequencespecific binding (1, 2). These designed proteins have been used to inhibit expression of target genes (3), and transcriptional activator domain-zinc finger conjugates have been used to activate expression of target genes (4). The RNA-interference pathway can be used to regulate gene expression at the post-transcriptional level (5). siRNA and shRNA molecules enlist cellular machinery to degrade selected mRNA targets (6, 7). RNAi technology has been highly effective in achieving potent and specific knock-down of target mRNAs and is now widely used to probe target gene function (8). However, bioavailability and delivery of zinc finger proteins and siRNA to targets in humans could be an obstacle to their therapeutic application and continues to receive considerable attention (8). Cellpermeable small molecules that modulate protein-protein or protein-DNA interactions offer another approach to the control of endogenous gene regulation. Screening small molecule and natural product libraries for a desired effect can identify candidate molecules with high likelihoods of possessing drug-like bioavailability; drawbacks include the need to screen anew for each target protein-protein or protein-DNA interaction. Polyamides containing *N*-methylimidazole (Im) and *N*-methylpyrrole (Py) are a class of programmable DNA-binding small molecules previously used to disrupt protein-DNA interactions in a sequence-specific manner in cell culture (9, 10).

Controlling the transcriptional activity of hypoxia inducible factor (HIF-1), a heterodimer of HIF-1 α and HIF-1 β (ARNT), is a clinically relevant goal (*11–14*). HIF-1 is the principal mediator of the adaptive cellular response to hypoxia (*15*). Under normoxic conditions, HIF-1 α is specifically hydroxylated by an iron-dependent proline hydroxylase, ubiquitinated by the von Hippel-Lindau (VHL) tumor suppressor protein-ubiquitin ligase

protein complex, and degraded by the proteosome (16). Iron chelators, such as deferoxamine (DFO), can be used to mimic hypoxia in cell culture (16). Through interaction with coactivators p300/CBP, HIF-1 directly activates the expression of at least 100 genes involved in cellular and tissue adaptation to hypoxia (13), including pro-angiogenic factors such as vascular endothelial growth factor (VEGF), glycolytic enzymes, extra-cellular matrix remodeling enzymes, and genes involved in both pro-survival and death pathways (11). HIF-1 activation by the hypoxic microenvironment of solid tumors, or by deactivating mutations in VHL, contribute to an aggressive phenotype of increased cell proliferation, invasion, metastasis, and drug resistance (11). Given the complexity of HIF-1 signaling in cellular survival and death pathways, and its critical role in physiological processes in normal tissues, it could be advantageous to target a subset of HIF-1 effectors rather than the entire pathway (13).

In important proof-of-principle experiments, introduction of siRNA against HIF- 2α to VHL^{-/-} renal carcinoma cells was sufficient to abrogate tumor formation by these cells in mice (17). Screening for small molecules capable of disrupting the HIF-1-p300 interaction identified chetomin, a natural product that binds p300 that was shown to inhibit expression of HIF-1 regulated genes and exhibit anti-tumor activity in a mouse model (18). In an effort to inhibit directly HIF-1-DNA binding, a hairpin polyamide was designed to bind the sequence 5'-ATACGT-3' found in the VEGF HRE. This polyamide bound its target site and prevented HIF-1-DNA binding in a sequence specific manner, and inhibited hypoxia-induced expression of VEGF and several other HIF-1 regulated genes in cultured cells without the use of transfection agents (19, 20). Melillo and colleagues screened a library of small molecules for inhibition of HIF-1 mediated transcription in a cell-based assay and identified the natural product echinomycin, a DNA-binding bisintercalator (21). Echinomycin binds the four basepair sequence 5'-NCGN-3' found in the consensus HRE 5'-NACGTG-3' and inhibited expression of VEGF in cultured cells (22).

To establish a benchmark of comparison for the global effects of polyamides,

we compare the genome-wide effects on U251 cells induced with DFO of a polyamide targeted to the HRE sequence 5'-WTWCGW-3', echinomycin, and siRNA targeted against HIF-1 α . siRNA-mediated destruction of HIF-1 α mRNA establishes a maximum level of inhibition that can be achieved for HIF-1 target genes through disruption of the HIF-1-HRE interaction. Nearly all transcripts induced by DFO are inhibited by both HIF-1 α siRNA and echinomycin. Polyamide **1** inhibits only a subset of these genes, and shows a preference for genes containing HREs of the sequence 5'-(T/A)ACGTG-3', consistent with this molecule's predicted binding preferences. Remarkably, the total number of genes affected by either polyamide **1** or HIF-1 α siRNA over a range of thresholds is comparable. We show that HIF-1 occupancy at the HREs of two genes affected by polyamide **1** is reduced in the presence of the polyamide, while HIF-1 occupancy at the HREs of two unaffected genes is unchanged. We also show that a polyamide that binds a site immediately 5' to the VEGF HRE inhibits induced expression of VEGF but not of FLT1, a HIF-1 target gene lacking this flanking site.



Figure 3.1. Structures of molecules used in this study. a) Structures of polyamides 1–3 and echinomycin. Imidazole, pyrrole and chlorothiophene monomer units are represented, respectively, by closed circles, open circles and squares. b) Three approaches to inhibiting HIF-1 induced gene expression: sequence-specific small molecule binding to the HRE by a polyamide or echinomycin, and reduction in HIF-1 α mRNA using siRNA.

3.3 Results and Discussion

Binding of Polyamide 1, 2, 3 and Echinomycin at the VEGF and FLT1 HREs. Polyamide sequence specificity is programmed by side-by-side pairings of heterocyclic amino acids in the minor groove of DNA: Im/Py distinguishes G•C from C•G; Py/Py binds both A•T and T•A; 3-chlorothiophene/N-methylpyrrole (Ct/Py) prefers T•A at the N-terminus position (23–25). Polyamide 1 and echinomycin are expected to bind at the VEGF HRE sequence 5'-TACGTG-3'. Polyamide 2 is expected to bind the sequence 5'-AGTGCA-3' immediately 5' to the VEGF promoter HRE. HRE-mismatch control polyamide 3 targets the sequence 5'-WGGWCW-3', which is not found near the VEGF HRE. The DNA-binding affinities of

2, **3** and echinomycin for the VEGF HRE were measured by quantitative DNase I footprint titrations using a 5' ³²P-labeled PCR amplification product of the plasmid pGL2-VEGF-*Luc* containing the VEGF HRE (Figure 3.2A and 3.2B). Polyamide **1** was previously found to bind the VEGF HRE ($K_a = 2.6 (\pm 0.4) \times 10^{10} \text{ M}^{-1}$) (20). For ease of comparison, a footprinting gel of **1** is included. Polyamide **2** binds the site 5'-AGTGCA-3' immediately 5' to the VEGF HRE ($K_a = 3.2 (\pm 0.6) \times 10^9 \text{ M}^{-1}$). Echinomycin binds the VEGF HRE ($K_a = 8.4 (\pm 2.1) \times 10^6 \text{ M}^{-1}$). Polyamide **3** binds the VEGF HRE ($K_a = 8.0 (\pm 1.0) \times 10^8 \text{ M}^{-1}$), and approximately 35 basepairs 3' of the HRE, most likely two 5'-WGWCW-3' sites ($K_a = 7.6 (\pm 1.0) \times 10^8 \text{ M}^{-1}$).

The DNA-binding affinities of **1**, **2**, **3** and echinomycin for the HRE of FLT1 were measured by quantitative DNase I footprint titrations using a 5'³²P-labeled PCR amplification product of the plasmid pCSJ-FLT1 (Figure 3.2C and 3.2D). Although formally targeted to the sequence 5'-WTWCGW-3', polyamide **1** would be expected to bind 5'-CAACGT-3', albeit with a moderate decrease in affinity (*25*). The sequence preference of a Ct/Py pair for T•A is approximately 4-fold over A•T, but 50-fold over a G•C (*25*). Polyamide **1** binds the FLT1 HRE ($K_a = 2.7 (\pm 0.2) \times 10^9 \text{ M}^{-1}$). Polyamide **2** binds the FLT1 HRE ($K_a = 2.2 (\pm 0.8) \times 10^8 \text{ M}^{-1}$). Echinomycin binds the FLT1 HRE ($K_a = 2.9 (\pm 0.7) \times 10^7 \text{ M}^{-1}$). Polyamide **3** does not bind the FLT1 HRE with a measurable K_a , but was observed to bind a 5'-AGACA-3' site 16 basepairs 5' to the FLT1 HRE ($K_a = 2.7 (\pm 0.4) \times 10^9 \text{ M}^{-1}$). These K_a data are summarized in Table 3.1.

pGL2-VEGF-Luc	Ligand	Sequence	K (M ⁻¹)	
*	1	5'-ATACGT-3'	$2.6 (\pm 0.4) \times 10^{10}$	
	2	5'-AGTGCA-3'	3.2 (±0.6) x 10 ⁹	
	3	5'-TGGGCT-3'	8.0 (±1.0) x 10 ⁸	
	Е	5'-ATACGT-3'	8.4 (±2.1) x 10 ⁶	
pCSJ-FLT1	Ligand	Sequence	K (M ⁻¹)	
*	1	5'-CAACGT-3'	2.7 (±0.2) x 10 ⁹	
	2	5'-TGAGGA-3'	2.2 (±0.8) x 10 ⁸	
	3	5'-CAACGT-3'		
	E	5'-CAACGT-3'	$2.9 (\pm 0.7) \times 10^7$	

Table 3.1. K_a values (M⁻¹) for 1, 2, 3, and Echinomycin (E) at HRE on pGL2-VEGF-*Luc* and pCSJ-FLT1

K_a values reported are mean values of three DNase 1 footprint titration experiments; standard deviations in parentheses.



Figure 3.2. Quantitative DNase I footprint titration experiments for polyamides **1–3** and echinomycin, **E**. a) Illustration of pGL2-VEGF-*Luc* and partial sequence containing VEGF HRE and polyamides **1**, **2** and echinomycin putative binding sites. b) Quantitative DNase I footprint titration experiments for polyamides **1**, **2**, **3**, and **E**, on 5'-end-labeled PCR product of plasmid pGL2-VEGF-*Luc*. For **1**, **2**, and **3**: lanes 1–11, 100 nM, 30 nM, 10 nM, 3 nM, 1 nM, 300 pM, 100 pM, 30 pM, 10 pM, 3 pM and 1 pM polyamide, respectively; lane 12, DNAse I standard; lane 13, intact DNA; lane 14, A reaction; lane 15, G reaction. For **E**: lanes 1–11, 10 μ M, 3 μ M, 1 μ M, 300 nM, 100 nM, 30 nM, 10 nM, 3 nM, 1 nM, 300 pM, 100 pM echinomycin, respectively; lanes 12-15 as above. Polyamide **1** and **E** K_a = 2.6 (±0.4) × 10¹⁰ M⁻¹ and K_a = 8.4 (±2.1) × 10⁶ M⁻¹, respectively, at VEGF HRE. Polyamide **2** K_a = 3.2 (±0.6) × 10⁹ M⁻¹ for 5'-AGTGCA-3' site immediately 5' to VEGF HRE. Polyamide **3** K_a = 8.0 (±1.0) × 10⁸ M⁻¹ for VEGF HRE. c) Illustration of pCSJ-FLT1 and partial sequence containing FLT1 HRE and putative binding sites for **1** and **E**. d) Quantitative DNase I footprint titration experiments for **1**, **2**, **3**, and **E** on 5' end-labeled PCR product of plasmid pCSJ-FLT1. Lane assignments for gels in d) are as for b). Polyamide **1** and **E** K_a = 2.7 (±0.2) × 10⁹ M⁻¹ and K_a = 2.9 (±0.7) × 10⁷ M⁻¹, respectively, at FLT1 HRE. Polyamide **2** K_a = 2.2 (±0.8) × 10⁸ at this site. Polyamide **3** does not bind FLT1 HRE with a measurable K_a.

Suppression of induced VEGF and FLT1 expression. HIF-1 induces VEGF expression by binding to the 5'-TACGTG-3' HRE located approximately 950 base pairs upstream of the transcription start site (26, 27). The effect on induced VEGF expression by siRNA silencing of HIF-1 α mRNA establishes a theoretical maximum level of inhibition through disruption of HIF-1-DNA binding. HIF-1 α mRNA was reduced by approximately 95% in the presence of HIF-1 α siRNA, but was minimally affected by polyamides 1, 2 or echinomycin under induced conditions (Figure 3.3). A mismatch control siRNA did not reduce HIF-1 α mRNA. Polyamides 1 and 2 (1 μ M) and HIF-1 α siRNA had similar effects on induced VEGF expression; treatment inhibited most of the increase in VEGF mRNA following DFO treatment, but not to levels below that observed for non-induced controls (Figure 3.3A). As previously reported, 100 nM echinomycin potently inhibits VEGF expression to levels below the non-induced control (21). HRE-mismatch control polyamide 3, which binds the HRE with a much reduced affinity relative to 1, had a more modest effect on VEGF mRNA levels. It is also possible that the modest but measurable effect of polyamide 3 on VEGF expression could be due to interference with other protein-DNA interactions elsewhere in the promoter or enhancer of VEGF, or is due to secondary effects from direct effects on other genes. A mismatch control siRNA had a limited effect on VEGF mRNA levels.

Induction of FLT1 (VEFG receptor type 1) is mediated by HIF-1 binding to the 5'-AACGTG-3' HRE in the FLT1 promoter (28). Polyamide 1 and HIF-1 α siRNA both inhibited FLT1 expression following DFO induction (Figure 3.3B). Echinomycin reduced FLT1 expression to below that of the non-induced control. Polyamides 2 and 3 had minimal effect. A mismatch control siRNA also had a limited effect on FLT1 mRNA levels. Given the relative binding affinities of polyamide 1 and echinomycin, it may be surprising that 1 μ M of polyamide 1 is necessary to inhibit VEGF and FLT1 expression comparably to HIF-1 α siRNA, while 100 nM echinomycin reduces their expression to sub-basal levels. The structure of double-helical DNA is not greatly perturbed by minor groove-binding hairpin

polyamides (23); echinomycin-DNA binding causes local unwinding and lengthening of the DNA helix, which might account for its greater potency in these experiments (29, 30). Polyamide-intercalator conjugates have been shown to unwind DNA in a sequence-specific fashion and to improve the ability of a polyamide to inhibit binding of several DNA-binding proteins *in vitro* (31, 32). Attempts to use polyamide-intercalator conjugates to target the VEGF HRE have not been successful due to poor nuclear uptake.

The ability to target DNA sequences flanking critical protein-DNA binding sites while maintaining productive inhibition of protein-DNA binding expands the repertoire of such interactions amenable to inhibition by polyamides. In a similar approach, Kageyama *et al.* showed that polyamides targeted to sequences flanking the VEGF HRE could inhibit VEGF expression (*33*). Polyamides targeted to flanking sites have previously successfully inhibited protein-DNA binding in the cases of TATA-binding protein and LEF-1 (*34*). It should be noted that minor groove-binding polyamides and some major groove-binding proteins co-occupy DNA sequences in some cases (*35*).



Figure 3.3. Quantitative real-time PCR measurements. a) Induction of VEGF mRNA by deferoxamine (DFO) measured by quantitative real-time PCR: HIF-1 α siRNA, **R**; mismatch control siRNA, **R***; echinomycin (100 nM), **E**; and polyamides **1**, **2**, and **3** (1 μ M). Treatment with siRNA, **1**, or **2** decrease VEGF mRNA levels to near non-induced levels. Echinomycin potently inhibits VEGF mRNA to below non-induced levels. Polyamide **3** has a more modest effect. b) Induction of FLT1 mRNA by DFO measured by quantitative real-time PCR: HIF-1 α siRNA, **R**; mismatch control siRNA, **R***; echinomycin (100 nM), **E**; and polyamides **1**, **2**, and **3** (1 μ M). The siRNA, **E**, and **1** decrease FLT1 mRNA levels. Polyamides **2** and **3** have minimal or no effect. c) Measurement of HIF-1 α mRNA by quantitative real-time PCR: HIF-1 α siRNA, **R**; mismatch control siRNA, **R***; echinomycin (100 nM), **E**; and polyamides **1**, **3** (1 μ M). Treatment with siRNA decreases HIF-1 α mRNA by more than 95%.

Microarray analysis of gene expression. One potential limitation to the use of hairpinpolyamides for gene regulation is binding site size and specificity for match versus mismatch sites, which may result in prohibitively large numbers of affected genes. To examine this, the global effects of polyamide treatment on hypoxia-induced gene expression were measured using Affymetrix Human Genome U133 Plus 2.0 Arrays containing oligonucleotide sequences representing more than 50,000 transcripts. To establish a benchmark for comparison, the effects of HIF-1 α siRNA and echinomycin were also measured. Experiments were conducted in triplicate, and gene expression levels normalized to DFO-treated controls. Cells not treated with DFO were normalized to DFOtreated controls.

Polyamide 1 (1 μ M) affected expression of 2,284 transcripts by more than 2-fold (p \leq 0.01) (Table 3.2). At the same threshold, HIF-1 α siRNA affected 3,190 transcripts and echinomycin (100 nM) affected 10,906. In all cases, a majority of affected genes were down-regulated. For comparison, DFO treatment alone affected expression of 2,142 transcripts (4.6% of interrogated transcripts), with a majority up-regulated.

Treatment	-	R	\mathbf{E}	1
DFO	-	+	+	+
up-regulated (fold change ≥ 2.0)	662	1,380	3,480	709
down-regulated (fold change \leq -2.0)	1,480	1,810	7,426	1,575
up-regulatd (fold change ≥ 4.0)	62	122	413	57
down-regulated (fold change \leq -4.0)	296	356	4,133	336

Table 3.2. Number of transcripts affected ($p \le 0.01$). **R**, HIF-1 α siRNA; **E**, echinomycin; **1**, polyamide.

Clustering analysis was performed to identify similarities in the expression profiles between the different treatments (Figure 3.4). The expression profile of cells treated with HIF-1 α siRNA is similar to that of cells not treated with DFO under the conditions of the analysis; the expression profiles of echinomycin-treated and polyamide-treated cells are less similar to each other and to the other treatments. Analysis of transcripts affected by both **1** and HIF-1 α siRNA shows that 395 and 150 transcripts are commonly downand up-regulated, respectively, at least 2-fold ($p \le 0.01$). A similar analysis of transcripts affected by both 1 and echinomycin shows that 731 and 112 transcripts are commonly down- and up-regulated, respectively. Analysis of transcripts affected by both siRNA and echinomycin shows that 1140 and 443 transcripts are commonly down- and up-regulated, respectively. A side-by-side, genome-wide expression analysis of fluorescein-tagged analogs of polyamides 1 and 3 in DFO-induced cells was previously reported (*19*), and found that a majority of genes were uniquely affected by each polyamide, with a number of genes commonly affected, under the conditions of the experiments. It is not entirely unsurprising that there is some overlap in genes affected by polyamides targeted to different DNA sequences, given that we do not have a full understanding of all DNA sequences involved in the direct or indirect regulation of a given gene.



Figure 3.4. Microarray analysis of gene expression. a) Divisive clustering analysis over all interrogated transcripts for DFO-induced cells: HIF-1 α siRNA, **R**; echinomycin (100 nM), **E**; and polyamides **1** (1 μ M). b) Agglomerative clustering analysis over all 297 transcripts induced by DFO at least 4-fold (p ≤ 0.01). Of this transcript set, HIF-1 α siRNA inhibited 244, echinomycin inhibited 263, and polyamide **1** inhibited 69 by \geq 2-fold (p ≤ 0.01). c) Effects of the indicated treatments on a panel of genes previously characterized as direct targets of HIF-1 and also induced by DFO at least 1.5-fold (p ≤ 0.01) in this experiment. Treatments reported are an error-weighted average from three experiments.

We find that DFO induced the expression of a set of 297 transcripts by at least 4-fold ($p \le 0.01$) (Figure 3.4B). Of this set, 69 were inhibited by polyamide 1 by at least 2-fold ($p \le 0.01$). For comparison, HIF-1 α siRNA inhibited 244, and echinomycin 263 of the 297 DFO-induced transcripts. It is not known what proportion of these affected transcripts are direct HIF-1 targets. To examine more closely the effects of polyamide 1, HIF-1 α siRNA, and echinomycin on transcripts induced directly by HIF-1, we examined a limited set of 31 transcripts consisting of previously identified direct HIF-1 targets that were induced at least 1.5-fold ($p \le 0.01$) by DFO in this experiment (Figure 3.4C) (28, 36–45) Nearly all 31 transcripts in this set were down-regulated by HIF-1 α siRNA. In most cases, the expression was reduced to levels observed in cells untreated with DFO. Echinomycin treatment resulted in down-regulation of all 31 transcripts. For some genes, including VEGF, expression was reduced to levels far below those of the siRNA-treated cells and non-induced controls. Polyamide 1 inhibited the expression of 14 of these but displayed minimal effect on the others. Shown in Figure 3.5 is a Venn diagram representing transcripts commonly down- and up-regulated ($|fold-change| \ge 2.0, p \le 0.01$) by 1 and HIF- 1α siRNA, by 1 and echinomycin, and by HIF- 1α siRNA and echinomycin.

The HRE sequences for these genes, where known, are displayed in Table 3.3. Quantitative real-time RT–PCR was used to confirm the effects of polyamides **1**, **3**, echinomycin, and siRNA treatments on these 11 genes. RT-PCR confirms that siRNA and echinomycin reduced expression of all genes in this set. Polyamide **1** significantly affected four genes in this set. Polyamide **3** had a modest but measurable effect on VEGF expression, but little effect on the expression of all the others. Chromatin immunoprecipitation was used to measure HIF-1 occupancy at the VEGF and carbonic anhydrase IX (CA9) HRE, which were both affected by polyamide **1**, and PGK1, which was unaffected (Figure 3.6). HIF-1 α occupancy at the VEGF HRE was decreased by HIF-1 α siRNA, echinomycin, and polyamide **1**, but less so by HRE-mismatch polyamide **3**. HIF-1 occupancy at the CA9 locus was also decreased by HIF-1 α siRNA, echinomycin, and polyamide **1**, but was unaffected by **3**. HIF-1 occupancy at PGK1 was markedly decreased by siRNA, but minimally affected by polyamides **1** and **3**. Surprisingly, echinomycin did not appear to affect HIF-1 occupancy at this locus. It is interesting to note that all of the genes affected by polyamide **1** (Table 3.2) have HREs that fall within the sequence 5'-(T/A)ACGTG-3', consistent with the expected DNA binding preferences for **1**.

TFRC	agcg TACGT Gcctc	-2.0	-2.3	1.1	1.4
PKFB3	gcgg GACGTG acgc gacg CACGTG ggca	-5.5	-82.0	1.0	1.0
LDHA	ggcg GACGTG cggg ctca CACGTG ggtt	-1.7	-4.4	1.3	1.1
BNIP3	gccg CACGTG ccac	-9.4	-6.0	1.3	1.0
EGLN3	gggc TACGTG caga	-5.3	-33.6	1.0	-1.1
EGLN1	ggtg TACGTG caga	-3.4	-19.8	1.1	1.0
PGK1	gtga GACGTG cggc tgcc GACGTG cgct	-5.4	-3.3	-1.2	1.0
CA9	gctg TACGTG catt	-89.0	-9.4	-2.1	-1.1
VEGF	tgca TACGTG ggct	-3.4	-34.0	-2.0	-1.3
FLT1	gaac AACGTG gaat	-2.2	-4.7	-2.0	-1.1
EDN1	aggc AACGTG cagc	-3.5	-31.0	-2.5	1.3

Table 3.3. HIF-1 induced genes affected by HIF-1 α siRNA (R), echinomycin (E, 100 nm), 1 and 3 (1 μ M).HRE(s) (5' to 3')RE1

3.4 Discussion

In this experiment, polyamide 1 (1 μ M) affected expression of 2,284 transcripts by more than 2-fold (p \leq 0.01), which represents less than 5% of transcripts assayed. A search of the publicly available human genome for the sequence 5'-WTWCGW-3' finds 1,876,480 potential match sites for polyamide 1. This corresponds to an average of one binding site every 1,600 base pairs. The proportion of these sites accessible to polyamide binding in

the context of heterochromatin *in vivo* is currently unknown. Additionally, data from *in vitro* transcription experiments suggest that polyamides non-covalently bound within the coding region of a gene would not interfere with RNA polymerase activity at that locus (*34*). It would thus not be surprising if a significant fraction of polyamide-DNA binding events in a cell do not directly affect gene expression. In parallel to this, global analysis of transcription factor binding to chromatin *in vivo* has shown occupancy at promoters and enhancers without associated changes in gene expression at that locus (*46*).

Interestingly, polyamide 1 (1 μ M) affected the expression of fewer genes than HIF-1 α siRNA under the conditions of the experiment. A direct comparison in genomic specificity between polyamide and siRNA cannot be drawn from these limited data because a large proportion of the genes affected by siRNA are likely a result of silencing the target gene, HIF-1 α , rather than off-target effects involving post-transcriptional silencing of mRNA using the RNA interference pathway (47). If we eliminate from the total number of transcripts affected by the HIF-1 α siRNA (2-fold, p ≤ 0.01) all transcripts affected by treatment with DFO alone (1.5-fold, p ≤ 0.01) we are left with 1,523 affected transcripts. A similar treatment of the data for polyamide 1 results in 1,626 affected transcripts.



Figure 3.5: Venn diagrams representing transcripts commonly down- and up-regulated (|fold-change $| \ge 2.0$, p ≤ 0.01) by 1 and HIF-1 α siRNA, by 1 and echinomycin, and by HIF-1 α siRNA and echinomycin. Numbers inside intersections represent transcripts affected by both treatments.

It should also be noted that for most HIF-1 regulated genes affected by both polyamide and siRNA, inhibition by the polyamide was more modest than siRNA, suggesting incomplete abrogation of HIF-1 DNA binding by the polyamide. Approximately 23% of the 297 transcripts induced by DFO were inhibited by polyamide **1**. For genes where the functional HRE is known, the effects of treatment with polyamide **1** or echinomycin are, thus far, consistent with the expected binding preferences of these molecules.



Figure 3.6. Chromatin immunoprecipitation at three HREs. a) Chromatin immunoprecipitation of HIF-1 α at the VEGF HRE following DFO treatment: HIF-1 α siRNA, **R**; echinomycin (100 nM), **E**; and polyamides **1** and **3** (1 μ M). Occupancy is decreased in the presence of **R**, **E**, and **1**, but only modestly affected by **3**. Chromatin immunoprecipitation of HIF-1 α at the CA9 HRE (b), and PGK1 HRE (c).

HIF-1 is frequently overactive in tumors, and a number of direct targets in the HIF-1 pathway have become points of clinical intervention (48). Bevacizumab, an anti-VEGF antibody, and sorafenib and sunitinib, tyrosine kinase inhibitors with activity against the VEGF receptors, have shown some promise in clinical trials as cancer therapeutics (49– 51). Echinomycin had been previously brought to the clinic as a cancer therapeutic in phase I and II trials (52), based on observations that echinomycin exhibits potent antiproliferative effects on several tumor-derived cell lines (52, 53). However, survival benefit was not established (52). In light of recent work by Melillo and colleagues, re-examination of the clinical use of echinomycin in tumor types expected to be highly sensitive to HIF-1 activity may be justified (21).

The induction of pro-angiogenic, proliferative, metastatic, and glycolytic genes by HIF-1 are established as contributing to the cancer phenotype (*11*). Genes that promote

cell death, such as BNIP3 and NIX (BNIP3L), are also induced by hypoxia through HIF-1 (54). In this sense, HIF-1 plays dual roles in the survival and death pathways of tumor cells (12). A functional separation of these targets of HIF-1 at the level of HIF-1-DNA binding might have clinical relevance (12). Given sufficient knowledge of the particular regulatory sequences involved, one could, in principle, design a polyamide or cocktail of polyamides to affect a selected subset of target genes in the HIF-1 pathway, making use of the programmability of polyamide recognition for targeting particular HREs and flanking sequences. The utility of polyamides as regulators of hypoxia-induced gene expression awaits continued study in small animal models of HIF-1 activity.

3.5 Methods

Synthesis of Polyamides

Polyamides were synthesized by solid-phase methods on Kaiser oxime resin (Nova Biochem), cleaved from resin with 3,3'-diamino-*N*-methyl-dipropylamine and purified by reverse-phase HPLC (*55*). Isophthalic acid was activated with PyBOP (Nova Biochem) and conjugated to polyamides as previously described (*20*). Polyamide purity and identity assessed by HPLC, UV-visible spectroscopy, and MALDI-ToF MS (Figure 3.7).

Determination of DNA-binding affinities and sequence specificities. Quantitative DNase I footprint titration experiments were used to measure the binding affinities of polyamides **1**, **2**, **3**, and echinomycin on 5' ³²P-labeled fragments of pGL2-VEGF-*Luc and pCSJ-FLT1* containing promoter sequences containing the HREs of VEGF and FLT1, respectively. Quantitative DNase I footprint titration experiments were conducted as reported previously (*56*).

Measurement of hypoxia-induced gene expression. U251 cells were plated in 24-well plates at a density of $20-30 \times 10^3$ cells per well ($40-60 \times 10^3$ cells/ml) in RPMI (ATCC) supplemented with 5% FBS (Irvine Scientific). After 24 hours, polyamides were added to the adhered cells in cell media solutions at the appropriate concentration and incubated

with the cells for 48 hours. Hypoxic induction of gene expression was chemically induced by adding DFO to 300 µM for an additional 16 hours. When appropriate, echinomycin was added two hours prior to DFO stimulation. Isolation of RNA and subsequent cDNA synthesis were as previously described (19). When appropriate, HIF- 1α siRNA (HIF- 1α validated stealth duplex, Invitrogen) or mismatch control siRNA with similar GC content (Invitrogen) was transfected 48 hours prior to RNA isolation. Transfection of siRNA was achieved using Lipofectamine 2000 (Invitrogen) according to manufacturer's protocols. Quantitative real-time RT-PCR was performed using SYBR Green PCR Master Mix (Applied Biosystems) on an ABI 7300 instrument. Target gene mRNA was measured relative to ß-glucuronidase as an endogenous control. Primer employed were: VEGF, L 5'-AGGGCAGAATCATCACGAAG-3', R 5'-GGGTACTCC TGGAAGATGTCC-3'; ß-glucuronidase, L 5'-CTCATTTGGAATTTTGCCGATT-3', R 5'-CCGAGTGAAGATCCCCTTTTTA-3'; FLT1. L

5'-CAGCAACATGGGAAACAGAAT-3', R 5'-TAGAGTCAGCCACAACCAAGG-3'.

Chromatin Immunoprecipitation. U251 cells were plated in 15 cm diameter plates and left to attach overnight. Media, time course, DFO, polyamide, echinomycin, and siRNA treatments were as described above. After the 16 hour DFO treatment, cells were treated with 1% formaldehyde for 10 minutes. Chromatin was isolated and sheared. HIF-1α antibodies (Novus Biologicals) were used to immunoprecipitate HIF-1-bound DNA fragments. After crosslink reversal, PCRs using primers targeted to the regions of interest were used to assess enrichment of bound fragments as compared to mock-precipitated (no antibody) controls. PCRs were monitored either using SYBR Green PCR Master Mix (Applied Biosystems) on an ABI 7300 instrument, or directly visualized using gel electrophoresis. The following primers were used: VEGF, L 5'-CCTTTGGGTTTTGCCAGA-3', R 5'-CCAAGTTTGTGGAGCTGA-3'; CA9, L 5'-AAAAGGGCGCTCTGTGAGT-3', R 5'-GCTGACTGTGGGGTGTCC-3'; PGK1, L 5'-CCCTAAGTCGGGAAGGTT-3', R 5'-GTCCGTCTGCGAGGGTACTA-3'.

Analysis of Gene Expression with Oligonucleotide Microarrays. U251 cells were plated in 12-well plates at a density of 40–60 x 10³ cells per well. Media, time course, DFO, polyamide, echinomycin, and siRNA treatments were as described above. RNA was isolated as previously described. Further sample preparation for microarray experiments was carried out at the Millard and Muriel Jacobs Gene Expression Facility at Caltech. Labeled mRNA was hybridized to Affymetrix Human 133 arrays according to established protocols. Gene expression was analyzed using Resolver (Rosetta Biosoftware, Seattle).

Accession Codes. Data have been deposited in NCBIs Gene Expression Omnibus (GEO, http://www.ncbi.nlm.gov/geo/), accessible through GEO Series accession number GSE7535.



Figure 3.7: Characterization data for 3 A) Analytical HPLC trace and B) MALDI-ToF data.

References

1. Blancafort, P., Segal, D. J., and Barbas, C. F. (2004) Designing transcription factor architectures for drug discovery, *Mol. Pharmacol. 66*, 1361–1371.

2. Beerli, R. R., and Barbas, C. F. (2002) Engineering polydactyl zinc-finger transcription factors, *Nat. Biotechnol.* 20, 135–141.

3. Beerli, R. R., Segal, D. J., Dreier, B., and Barbas, C. F. (1998) Toward controlling gene expression at will: Specific regulation of the erbB-2/HER-2 promoter by using polydactyl zinc finger proteins constructed from modular building blocks, *Proc. Natl. Acad. Sci. U. S. A.* 95, 14628–14633.

4. Liu, P. Q., Rebar, E. J., Zhang, L., Liu, Q., Jamieson, A. C., Liang, Y. X., Qi, H., Li, P. X., Chen, B. L., et al. (2001) Regulation of an endogenous locus using a panel of designed zinc finger proteins targeted to accessible chromatin regions - Activation of vascular endothelial growth factor A, *J. Biol. Chem.* 276, 11323–11334.

5. Fire, A., Xu, S. Q., Montgomery, M. K., Kostas, S. A., Driver, S. E., and Mello, C. C. (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*, *Nature 391*, 806–811.

6. Meister, G., and Tuschl, T. (2004) Mechanisms of gene silencing by double-stranded RNA, *Nature 431*, 343–349.

 Mello, C. C., and Conte, D. (2004) Revealing the world of RNA interference, *Nature* 431, 338–342.

8. Hannon, G. J., and Rossi, J. J. (2004) Unlocking the potential of the human genome with RNA interference, *Nature 431*, 371–378.

9. Dervan, P. B., and Edelson, B. S. (2003) Recognition of the DNA minor groove by pyrrole-imidazole polyamides, *Curr. Opin. Struct. Biol.* 13, 284–299.

10. Nickols, N. G., and Dervan, P. B. (2007) Suppression of androgen receptor-mediated gene expression by a sequence-specific DNA-binding polyamide, *Proc. Natl. Acad. Sci. U. S. A. 104*, 10418–10423.

11. Semenza, G. L. (2003) Targeting HIF-1 for cancer therapy, *Nat. Rev. Cancer 3*, 721–732.

12. Zhou, J., Schmid, T., Schnitzer, S., and Brune, B. (2006) Tumor hypoxia and cancer progression, *Cancer Lett.* 237, 10–21.

13. Pouyssegur, J., Dayan, F., and Mazure, N. M. (2006) Hypoxia signalling in cancer and approaches to enforce tumour regression, *Nature 441*, 437–443.

14. Melillo, G. (2006) Inhibiting hypoxia-inducible factor 1 for cancer therapy, *Molecular Cancer Research 4*, 601–605.

15. Semenza, G. L. (2000) HIF-1 and human disease: one highly involved factor, *Genes Dev. 14*, 1983–1991.

Ivan, M., Kondo, K., Yang, H. F., Kim, W., Valiando, J., Ohh, M., Salic, A., Asara, J.
 M., Lane, W. S., et al. (2001) HIF alpha targeted for VHL-mediated destruction by proline hydroxylation: Implications for O-2 sensing, *Science 292*, 464–468.

17. Zimmer, M., Doucette, D., Siddiqui, N., and Iliopoulos, O. (2004) Inhibition of hypoxia-inducible factor is sufficient for growth suppression of VHL-/- tumors, *Molecular Cancer Research 2*, 89–95.

Kung, A. L., Zabludoff, S. D., France, D. S., Freedman, S. J., Tanner, E. A., Vieira,
 A., Cornell-Kennon, S., Lee, J., Wang, B. Q., et al. (2004) Small molecule blockade of transcriptional coactivation of the hypoxia-inducible factor pathway, *Cancer Cell 6*, 33–43.

Olenyuk, B. Z., Zhang, G. J., Klco, J. M., Nickols, N. G., Kaelin, W. G., and Dervan,
 P. B. (2004) Inhibition of vascular endothelial growth factor with a sequence-specific hypoxia response element antagonist, *Proc. Natl. Acad. Sci. U. S. A. 101*, 16768–16773.

20. Nickols, N. G., Jacobs, C. S., Farkas, M. E., and Dervan, P. B. (2007) Improved nuclear localization of DNA-binding polyamides, *Nucleic Acids Res. 35*, 363–370.

21. Kong, D. H., Park, E. J., Stephen, A. G., Calvani, M., Cardellina, J. H., Monks, A., Fisher, R. J., Shoemaker, R. H., and Melillo, G. (2005) Echinomycin, a small-molecule

inhibitor of hypoxia-inducible factor-1 DNA-binding activity, *Cancer Res.* 65, 9047–9055.

22. Van Dyke, M. M., and Dervan, P. B. (1984) Echinomycin Binding-Sites on DNA, *Science 225*, 1122–1127.

23. Kielkopf, C. L., Baird, E. E., Dervan, P. D., and Rees, D. C. (1998) Structural basis for G•C recognition in the DNA minor groove, *Nat. Struct. Biol. 5*, 104–109.

24. White, S., Szewczyk, J. W., Turner, J. M., Baird, E. E., and Dervan, P. B. (1998) Recognition of the four Watson-Crick base pairs in the DNA minor groove by synthetic ligands, *Nature 391*, 468–471.

25. Foister, S., Marques, M. A., Doss, R. M., and Dervan, P. B. (2003) Shape selective recognition of T•A base pairs by hairpin polyamides containing N-terminal 3-methoxy (and 3-chloro) thiophene residues, *Bioorg. Med. Chem. 11*, 4333–4340.

26. Liu, Y. X., Cox, S. R., Morita, T., and Kourembanas, S. (1995) Hypoxia Regulates Vascular Endothelial Growth-Factor Gene Expression in Endothelial Cells – Identification of a 5'-Enhancer, *Circ.Res.* 77, 638–643.

27. Forsythe, J. A., Jiang, B. H., Iyer, N. V., Agani, F., Leung, S. W., Koos, R. D., and Semenza, G. L. (1996) Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1, *Mol. Cell. Biol. 16*, 4604–4613.

28. Gerber, H. P., Condorelli, F., Park, J., and Ferrara, N. (1997) Differential transcriptional regulation of the two vascular endothelial growth factor receptor genes –Flt-1, but not Flk-1/KDR, is up-regulated by hypoxia, *J. Biol. Chem.* 272, 23659–23667.

29. Waring, M. J., and Wakelin, L. P. G. (1974) Echinomycin – Bifunctional Intercalating Antibiotic, *Nature 252*, 653–657.

30. Low, C. M. L., Drew, H. R., and Waring, M. J. (1984) Sequence-Specific Binding of Echinomycin to DNA – Evidence for Conformational-Changes Affecting Flanking Sequences, *Nucleic Acids Res. 12*, 4865–4879.

31. Fechter, E. J., and Dervan, P. B. (2003) Allosteric inhibition of protein-DNA

complexes by polyamide-intercalator conjugates, J. Am. Chem. Soc. 125, 8476–8485.

32. Fechter, E. J., Olenyuk, B., and Dervan, P. B. (2004) Design of a sequence-specific DNA bisintercalator, *Angew. Chem.-Int. Edit. 43*, 3591–3594.

33. Kageyama, Y., Sugiyama, H., Ayame, H., Iwai, A., Fujii, Y., Huang, L. E., Kizaka-Kondoh, S., Hiraoka, M., and Kihara, K. (2006) Suppression of VEGF transcription in renal cell carcinoma cells by pyrrole-imidazole hairpin polyamides targeting the hypoxia responsive element, *Acta Oncol.* 45, 317–324.

34. Dickinson, L. A., Gulizia, R. J., Trauger, J. W., Baird, E. E., Mosier, D. E., Gottesfeld,
J. M., and Dervan, P. B. (1998) Inhibition of RNA polymerase II transcription in human
cells by synthetic DNA-binding ligands, *Proc. Natl. Acad. Sci. U. S. A.* 95, 12890–12895.

35. Oakley, M. G., Mrksich, M., and Dervan, P. B. (1992) Evidence That a Minor Groove-Binding Peptide and a Major Groove-Binding Protein Can Simultaneously Occupy a Common Site on DNA, *Biochemistry 31*, 10969–10975.

36. Tacchini, L., Bianchi, L., Bernelli-Zazzera, A., and Cairo, G. (1999) Transferrin receptor induction by hypoxia – HIF-1-mediated transcriptional activation and cell-specific post-transcriptional regulation, *J. Biol. Chem.* 274, 24142–24146.

37. Lok, C. N., and Ponka, P. (1999) Identification of a functional hypoxia-response element in the transferrin-receptor gene, *Exp. Hematol.* 27, 91–91.

38. Fukasawa, M., Tsuchiya, T., Takayama, E., Shinomiya, N., Uyeda, K., Sakakibara, R., and Seki, S. (2004) Identification and characterization of the hypoxia-responsive element of the human placental 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene, *J. Biochem.* 136, 273–277.

39. Obach, M., Navarro-Sabate, A., Caro, J., Kong, X. G., Duran, J., Gomez, M., Perales, J. C., Ventura, F., Rosa, J. L., et al. (2004) 6-phosphofructo-2-kinase (pfkfb3) gene promoter contains hypoxia-inducible factor-1 binding sites necessary for transactivation in response to hypoxia, *J. Biol. Chem.* 279, 53562–53570.

40. Semenza, G. L., Jiang, B. H., Leung, S. W., Passantino, R., Concordet, J. P., Maire,

P., and Giallongo, A. (1996) Hypoxia response elements in the aldolase A, enolase 1, and lactate dehydrogenase A gene promoters contain essential binding sites for hypoxia-inducible factor 1, *J. Biol. Chem. 271*, 32529–32537.

41. Semenza, G. L., Roth, P. H., Fang, H. M., and Wang, G. L. (1994) Transcriptional Regulation of Genes Encoding Glycolytic-Enzymes by Hypoxia-Inducible Factor-1, *J. Biol. Chem.* 269, 23757–23763.

42. Pescador, N., Cuevas, Y., Naranjo, S., Alcaide, M., Villar, D., Landazuri, M., and Del Peso, L. (2005) Identification of a functional hypoxia-responsive element that regulates the expression of the egl nine homologue 3 (egln3/phd3) gene, *Biochem. J. 390*, 189–197.

43. Willam, C., Nicholls, L. G., Ratcliffe, P. J., Pugh, C. W., and Maxwell, P. H.(2004) The prolyl hydroxylase enzymes that act as oxygen sensors regulating destruction of hypoxia-inducible factor alpha, *Advances in Enzyme Regulation*, *44*, 75–92.

44. Grabmaier, K., de Weijert, M. C. A., Verhaegh, G. W., Schalken, J. A., and Oosterwijk, E. (2004) Strict regulation of CAIX(G250/MN) by HIF-1 alpha in clear cell renal cell carcinoma, *Oncogene 23*, 5624–5631.

45. Hu, J., Discher, D. J., Bishopric, N. H., and Webster, K. A. (1998) Hypoxia regulates expression of the endothelin-1 gene through a proximal hypoxia-inducible factor-1 binding site on the antisense strand, *Biochem. Biophys. Res. Commun. 245*, 894–899.

46. Zhang, X. M., Odom, D. T., Koo, S. H., Conkright, M. D., Canettieri, G., Best, J., Chen, H. M., Jenner, R., Herbolsheimer, E., et al. (2005) Genome-wide analysis of cAMP-response element binding protein occupancy, phosphorylation, and target gene activation in human tissues, *Proc. Natl. Acad. Sci. U. S. A. 102*, 4459–4464.

47. Jackson, A. L., Bartz, S. R., Schelter, J., Kobayashi, S. V., Burchard, J., Mao, M., Li, B., Cavet, G., and Linsley, P. S. (2003) Expression profiling reveals off-target gene regulation by RNAi, *Nat. Biotechnol.* 21, 635–637.

48. Cohen, H. T., and McGovern, F. J. (2005) Renal-cell carcinoma, *N. Engl. J. Med. 353*, 2477–2490.

49. Escudier, B., Eisen, T., Stadler, W. M., Szczylik, C., Oudard, S., Siebels, M., Negrier, S., Chevreau, C., Solska, E., et al. (2007) Sorafenib in advanced clear-cell renal-cell carcinoma, *N. Engl. J. Med.* 356, 125–134.

50. Motzer, R. J., Hutson, T. E., Tomczak, P., Michaelson, M. D., Bukowski, R. M., Rixe, O., Oudard, S., Negrier, S., Szczylik, C., et al. (2007) Sunitinib versus interferon alfa in metastatic renal-cell carcinoma, *N. Engl. J. Med.* 356, 115–124.

51. Yang, J. C., Haworth, L., Sherry, R. M., Hwu, P., Schwartzentruber, D. J., Topalian, S. L., Steinberg, S. M., Chen, H. X., and Rosenberg, S. A. (2003) A randomized trial of bevacizumab, an anti-vascular endothelial growth factor antibody, for metastatic renal cancer, *N. Engl. J. Med.* 349, 427–434.

 Foster, B. J., Clagettcarr, K., Shoemaker, D. D., Suffness, M., Plowman, J., Trissel,
 L. A., Grieshaber, C. K., and Leylandjones, B. (1985) Echinomycin – the 1st Bifunctional Intercalating Agent in Clinical Trials, *Invest. New Drugs 3*, 403–410.

53. May, L. G., Madine, M. A., and Waring, M. J. (2004) Echinomycin inhibits chromosomal DNA replication and embryonic development in vertebrates, *Nucleic Acids Res.* 32, 65–72.

54. Kothari, S., Cizeau, J., McMillan-Ward, E., Israels, S. J., Bailes, M., Ens, K., Kirshenbaum, L. A., and Gibson, S. B. (2003) BNIP3 plays a role in hypoxic cell death in human epithelial cells that is inhibited by growth factors EGF and IGF, *Oncogene 22*, 4734–4744.

55. Belitsky, J. M., Nguyen, D. H., Wurtz, N. R., and Dervan, P. B. (2002) Solid-phase synthesis of DNA binding polyamides on oxime resin, *Bioorg. Med. Chem.* 10, 2767–2774.

56. Trauger, J. W., and Dervan, P. B. (2001) Footprinting methods for analysis of pyrrole-imidazole polyamide/DNA complexes, *Methods Enzymol. 340*, 450–466.