

LIST OF FIGURES

Figure 2-1: Cartoon representation of the components implicated in electron transfer for Fe(II) oxidation by *Acidithiobacillus ferrooxidans* strain ATCC 33020. The product of the *iro* gene is not thought to play a role in this strain, but may in others32

Figure 2-2: Genes proposed to encode the components of Fe(II) oxidation in *Acidithiobacillus ferrooxidans* strain ATCC 3302035

Figure 3-1: Growth of TIE-1 and SW2 on 4 mM Fe(II)Cl₂·H₂O + varying concentrations of NTA. A. Data for TIE-1: ◆ - 0 mM NTA, ■ - 7.5 mM NTA, ▲ - 15 mM NTA, ● - 20 mM NTA, ✕ - Abiotic + 20 mM NTA. B. Data for SW2: ◆ - 0 mM NTA, ■ - 7.5 mM NTA, ● - 10 mM NTA, ▲ - 15 mM NTA, ✕ - Abiotic + 20 mM NTA. No growth was observed in cultures of TIE-1 or SW2 where only NTA and no Fe(II) was added, indicating that these strains cannot use NTA as a substrate for growth. The lower concentration of Fe(II) at time 0 in the cultures where NTA has been added as compared to the cultures with no NTA addition indicates there is a pool of Fe(II) we cannot measure with the *Ferrozine* assay. Error bars represent the error on duplicate cultures43

Figure 3-2: Concentrations of A. Fe(II) and B. NTA species in the phototrophic basal medium (pH 6.8). Concentrations are represented as % of total Fe(II) (4 mM Fe(II)Cl₂·H₂O) and NTA (20 mM Na₂NTA) as calculated with MINEQL⁺. See text for model parameters45

Figure 3-3: H₂ inhibits the Fe(II) oxidation activity of both TIE-1 and SW2 to varying degrees depending on the concentration of NaHCO₃. A. Data for TIE-1: ▲ - H₂ + 1 mM NaHCO₃ + 0.5 mM FeCl₂·H₂O; ● - N₂ + 1 mM NaHCO₃ + 0.5 mM FeCl₂·H₂O; ◆ - H₂ + 20 mM NaHCO₃ + 0.5 mM FeCl₂·H₂O; ■ - N₂ + 20 mM NaHCO₃ + 0.5 mM FeCl₂·H₂O. B. Data for SW2: ▲ - H₂ + 1 mM NaHCO₃ + 0.5 mM FeCl₂·H₂O; ● - N₂ + 1 mM NaHCO₃ + 0.5 mM FeCl₂·H₂O; ◆ - H₂ + 20 mM NaHCO₃ + 0.5 mM FeCl₂·H₂O; ■ - N₂ + 20 mM NaHCO₃ + 0.5 mM FeCl₂·H₂O. Data are representative of at least two independent experiments. The volume of the assay was 1 ml and the assay bottles were shook vigorously to ensure maximal H₂ saturation of the cell suspension solution. Error bars represent the error on duplicate cell suspension assays for TIE-1 and triplicate assays for SW2.....49

Figure 3-4: The Fe(II) oxidation activity of cell suspensions of TIE-1 and SW2 is completely light dependent. ◆ - H₂ pre-grown TIE-1 cells + N₂ + 20 mM NaHCO₃ + 1 mM FeCl₂·H₂O, incubated in the dark. ■ - H₂ pre-grown SW2 cells + N₂ + 20 mM NaHCO₃ + 2 mM FeCl₂·H₂O, incubated in the dark51

Figure 3-5: Fe(II) oxidation activity of cell suspensions of TIE-1 pre-grown photoautotrophically on different inorganic electron donors. ■ - TIE-1 pre-grown on 10 mM thiosulfate, ◆ - TIE-1 pre-grown on H₂, ▲ - TIE-1 pre-grown on 4 mM FeCl₂·H₂O + 10 mM NTA. Error bars represent the error on triplicate cell suspension assays53

Figure 3-6: The Fe(II) oxidation activity of H₂ pre-grown cells of TIE-1 decreases with increasing concentration of gentamicin. All assays here contain 1 mM FeCl₂·H₂O and 20 mM NaHCO₃. - (short dash) - no gentamicin added; ■ - 0.1 mg/ml gentamicin; ◆ - 0.2 mg/ml gentamicin; ▲ - 0.5 mg/ml gentamicin; ● - 1 mg/ml gentamicin; + - 2 mg/ml gentamicin; ✕ - 4 mg/ml gentamicin; — (long dash) abiotic control + 4 mg/ml gentamicin.....54

Figure 3-7: A cartoon representation of the flow of electrons from Fe(II) and H₂ to the photosynthetic electron transport chain and CO₂. For simplicity, Fe(II) oxidation is represented as occurring outside the cell. The red lines, associated with k₁, represent the pathway and the overall rate of electron flow from Fe(II) to the photosynthetic electron transport chain. The blue lines, associated with k₂, represent the pathway and the overall rate of electron flow from H₂ to the photosynthetic electron transport chain. OM: outer membrane; PERI: periplasm; CM: cytoplasmic membrane; ICM: intracytoplasmic membrane; CYT: cytoplasm.56

Figure 3-8: Hydrogenase and Fe(II) oxidation activity for TIE-1 as measured by benzyl viologen (BV) reduction and *Ferrozine*, respectively. A. Hydrogenase activity for TIE-1: ◆ - H₂ + 20 mM NaHCO₃ + 1 mM FeCl₂·H₂O + 5 mM BV; ● - H₂ + 20 mM NaHCO₃ + 5 mM BV. B. Fe(II) oxidation activity for TIE-1: ▲ H₂ + 20 mM NaHCO₃ + 1 mM FeCl₂·H₂O; ● - N₂ + 20 mM NaHCO₃ + 1 mM FeCl₂·H₂O. The volume of the assay was 1 ml and the assay bottles were shook vigorously to ensure maximal H₂ saturation of the cell suspension solution. Error bars represent the error on triplicate cell suspension assays.....59

Figure 4-1: Fe(II)-oxidation by the two enrichment cultures and *Thiodictyon* strain F4. ● - F4, ■ - enrichment 1, ▲ - enrichment 2, □ - uninoculated control, ◇ - medium inoculated with F4, incubated in the dark. All cultures, except the dark control, were incubated at 40 cm from the 40 W light source. The dark control is representative of dark controls performed with the two enrichment cultures. Iron contents for the uninoculated and dark controls are consistent over time within analytical errors. Data for the enrichment cultures and *Thiodictyon* strain F4 were collected in separate experiments105

Figure 4-2: Fe(II)-oxidation by cultures of *Thiodictyon* strain F4 inoculated with approximately the same number of cells and incubated at 40, 80, and 120 cm from the light source. ◆ - F4 incubated at 40 cm from the light, ■ - 80 cm, ▲ - 120 cm. The data shown are representative of duplicate cultures..... 106

Figure 4-3: Differential interference contrast (DIC) micrographs of the enrichments and *Thiodictyon* strain F4. A. A representative micrograph of the two enrichments growing photosynthetically on 10 mM Fe(II)_{aq} supplemented with 1 mM acetate. Three major cell morphologies are observed: approximately 1-1.5 μm by 4-5 μm, rod shaped cells with gas vacuoles (light areas within the cells) which tended to aggregate around the HFO precipitates (I), 1.5-2 μm by 3.5-4 μm rod shaped cells with no gas vesicles (II) and 0.5-1 μm by 1.5-2 μm rod shaped cells (III). B. DIC micrograph of *Thiodictyon* strain F4, growing photosynthetically on 10 mM Fe(II)_{aq}. Cells are approximately 1.5-2 μm by 5-7 μm and contain gas vacuoles. Note the similarity in size and shape between cells of *Thiodictyon* strain F4 and cells of type I in the enrichment culture. C. DGGE of the enrichments and *Thiodictyon* strain F4. From left to right lanes correspond to enrichment 1, enrichment 2 and *Thiodictyon* strain F4 107

Figure 4-4: The phylogenetic relationship of *Thiodictyon* strain F4 inferred from 16S rDNA sequences. The tree was constructed by the maximum-likelihood method using the ARB software package with 1250 positions considered. Bootstrap values above 50% from 100 bootstrap analyses are given at branch nodes. Anaerobic phototrophs able to oxidize Fe(II) are in blue to illustrate the evolutionary diversity of organisms capable of this form of metabolism. Aerobic phototrophs (cyanobacteria) and other organisms capable of oxidizing Fe(II) non-photosynthetically are also shown for phylogenetic comparison. Accession numbers are listed after the bacterium. PNSB – purple non sulfur bacteria, PSB – purple sulfur bacteria, GSB – green sulfur bacteria 109

Figure 4-5: Isotopic data for the two enrichments incubated at 40 cm from the light source and the uninoculated control. The δ⁵⁶Fe values for duplicate samples of the Fe(II)_{aq} and HFO fractions taken from single cultures are plotted as a function of time. A. Enrichment 1. ■ and □ - duplicate Fe(II)_{aq} fractions. ▲ and △ - duplicate HFO fractions. B. Enrichment 2. ■ and □ - duplicate Fe(II)_{aq} fractions. ▲ and △ - duplicate HFO fractions. C. The uninoculated control. ■ and □ - duplicate Fe(II)_{aq} fractions. The dashed line plots on graphs A., B., and C. are Fe(II)_{aq} concentrations (mM) as determined by *Ferrozine* assay and the error bars represent the error on triplicate assays for each time

point. The shaded box on each of the graphs illustrates the error on the isotopic measurements from the uninoculated control. In some cases the points are larger than the error 113

Figure 4-6: Isotopic data for *Thiodictyon* strain F4 incubated at 40, 80, and 120 cm from the light source and the uninoculated and dark controls. The $\delta^{56}\text{Fe}$ values for duplicate samples of the $\text{Fe(II)}_{\text{aq}}$ and HFO fractions taken from single cultures are plotted as a function of time. A. F4 incubated at 40 cm from the light. ■ and □ - duplicate $\text{Fe(II)}_{\text{aq}}$ fractions. ▲ and △ - duplicate HFO fractions. B. F4 incubated at 80 cm from the light. ■ and □ - duplicate $\text{Fe(II)}_{\text{aq}}$ fractions. ▲ and △ - duplicate HFO fractions. C. F4 incubated at 120 cm from the light. ■ and □ - duplicate $\text{Fe(II)}_{\text{aq}}$ fractions. ▲ and △ - duplicate HFO fractions. D and E. The uninoculated and dark controls, respectively. ■ and □ - duplicate $\text{Fe(II)}_{\text{aq}}$ fractions. The dashed line plots on graphs A., B., and C. are $\text{Fe(II)}_{\text{aq}}$ concentrations (mM) as determined by *Ferrozine* assay and the error bars represent the error on triplicate assays for each time point. The shaded box on each of the graphs illustrates the error on the isotopic measurements from the uninoculated and dark controls. In some cases the plotted points are larger than the error..... 115

Figure 4-7: Fe isotope fractionations between $\text{Fe(II)}_{\text{aq}}$ and HFO in the enrichments and *Thiodictyon* strain F4 cultures. $\Delta_{\text{Fe(II)}_{\text{aq}}-\text{HFO}}$ values are plotted as a function of "F", defined as the fraction toward complete oxidation of initial $\text{Fe(II)}_{\text{aq}}$. Note that the true isotopic fractionation factor (assuming it is constant over the reaction progress) is most closely constrained at low F values. Open and closed symbols of the same type represent the difference between the $\delta^{56}\text{Fe}$ values of $\text{Fe(II)}_{\text{aq}}$ and HFO samplings, in duplicate, for a particular culture. A. The enrichments. ■ and □ - enrichment 1. ▲ and △ - enrichment 2. B. *Thiodictyon* strain F4. ■ and □ - the 40 cm culture. ▲ and △ - the 80 cm culture. ◆ and ◇ - the 120 cm culture. Rayleigh (solid curved line) and closed-system (dashed straight line) equilibrium models are shown for comparison 117

Figure 5-1: Mutants 76H3 and A2 are specifically defective in Fe(II) oxidation. A. Normal growth of mutant 76H3 and A2 with H_2 as the electron donor. Data are representative of two independent cultures. B. Defects in phototrophic growth on Fe(II) for mutants 76H3 and A2 compared to wild type. Growth was stimulated with H_2 present in the headspace initially. Data are representative of duplicate cultures. C. Mutant 76H3 and A2 carrying plasmids pT198 and pT498, respectively, show 80% of Fe(II) oxidation compared to the wild type in the cell suspension assay. D. Organization of the genomic regions surrounding the mutated genes in mutants 76H3 and A2. The black

arrows indicate the disrupted genes and the transposon insertion sites are marked by the open triangles. The numbers provided below the open reading frames (all arrows) are consistent with the numbers given for the identical regions from the CGA009 genome 145

Figure 5-2: Restriction digests of ten randomly selected cosmids with *EcoRI* reveal that the cosmid genomic library of SW2 is representative of the SW2 genome and that the average insert size is approximately 23.5 kb. Lane 1 – λ *HindIII* molecular weight marker; lanes 2-12 – ten randomly selected cosmids digested with *EcoRI*; Lane 13 – 1 kb molecular weight marker (Bio-Rad)..... 149

Figure 5-3: Strains of *Rhodobacter capsulatus* SB1003 containing cosmid clones p2B3, p9E12, p11D3 and p12D4 show a decrease in Fe(II) that is 99% greater than the negative control, 1003 + pLAFR5. Error bars represent the error for cell suspension assays of 24, 14, 24, 10, and 9 independent colonies of 1003 containing p2B3, p9E12, p11D3, p12D4 and pLAFR5 respectively, and 4 independent cultures of SW2. The inset shows the color difference between the positive control, SW2 and the abiotic control after the addition of *Ferrozine* during a 96 well plate format cell suspension assay 150

Figure 5-4: In our 96 well format cell suspension assay, *Rhodobacter capsulatus* SB1003 shows a decrease in Fe(II), equivalent to approximately 73% of the total Fe(II) added. This decrease, however is less than that observed for the p2B3, p9E12, p11B3, and p12D4 containing 1003 strains..... 151

Figure 5-5: Fe(II) oxidation by 1003 + p2B3 (A), p9E12 (B), p11D3 (C), and p12D4 (D) in comparison to SW2 and 1003. On all graphs, ■ – Fe total for 1003 + cosmid; ▲ – Fe total for SW2; ◆ – Fe total for 1003; □ – Fe(II) for 1003 + cosmid strain; ◇ – Fe(II) for SW2; △ – Fe(II) for 1003. When both the concentration of Fe(II) and total Fe were followed in cell suspension assays over time, a decrease in the amount of Fe(II) was observed while the total amount of Fe stayed constant. This shows that Fe(II) is being oxidized by these cosmid containing strains rather than being sequestered within the cells or chelated by cell surface components or other cell produced molecules. Assays were normalized for cell number and the error bars represent the error on triplicate *Ferrozine* measurements. In these assays, it seem as though Fe(II) is not oxidized to completion by SW2 or the cosmid strains. This, however, is due to OD₅₇₀ absorbance by the high number of cells in the sample taken for the [Fe] measurement..... 153

Figure 5-6: Fe(II) oxidation conferring cosmids p2B3, p9E12, p11D3 and p12D4 do not confer the ability to grow photoautotrophically on

Fe(II) to 1003. ◆ – p2B3; ■ – p9E12; ▲ - p11D3; X – p12D4; ✕ – 1003;
● – SW2 154

Figure 5-7: Digestion of p2B3, p9E12, p11D3 and p12D4 with *EcoRI*, *HindIII* and *PstI* reveals common restriction fragments among three or more of these cosmids. The common fragments are highlighted by the white arrows. Lanes 1 and 14 – λ *HindIII* molecular weight marker; lanes 2-5 – p2B3, p9E12, p11D3 and p12D4 digested with *EcoRI*, respectively; lanes 6-9 – p2B3, p9E12, p11D3 and p12D4 digested with *HindIII*, respectively; lanes 10-13 – p2B3, p9E12, p11D3 and p12D4 digested with *PstI*, respectively. Some small pieces are missing from this gel. These include *HindIII* fragments of p2B3 (~0.5 kb) and p11D3 (~0.5 kb) and *PstI* fragments of p2B3 (~0.6 and 0.2 kb), p9E12 (~0.2 kb), p11D3 (~0.2 kb) and p12D4 (~0.6 and 0.7 kb)..... 155

Figure 5-8: Restriction map of p2B3, p9E12, p11B3 and p12D4 digested with *PstI*. The red bars represent fragments in the *PstI* digest common among p2B3, p9E12 and p11D3. The position of fragment 3 in the p12D4 *PstI* digest is inferred from the sizes of the other fragments in this digest relative to those in the digests of p2B3 and p9E12. This fragment likely represents a smaller piece of the common fragment in red due to partial digest conditions. The green bars represent the positions of the pH5 and pH6 inserts on fragment 3 of the p9E12 *PstI* digest inferred from sequence data. Fragment 6 in the *HindIII* digest of p9E12 was common among all the cosmids. From sequence data, fragments 7 and 3 in the p9E12 *PstI* digest are contiguous as are fragments 5, 6, and 2, and 1, although, the position of *PstI* fragment 7 with respect to fragment 3 is arbitrary (*i.e.*, fragment 7 could be on the other side of fragment 3). The positioning, however, is likely accurate given the nature of the sequence at the end of the P3 T7 and the sequence on fragment 4 (P4, 100% identical at the DNA and protein level to *E. coli*) in comparison to the sequence at the beginning of P3 T7 (97% identical at the protein level to *Rhodobacter sphaeroides* 2.4.1. The positions of p9E12 *PstI* fragments 4, 8 and 9 are inferred from comparison to the digests of the other cosmids 158

Figure 5-9: The 9.4 kb *PstI* fragment of p9E12 confers Fe(II) oxidation activity to *Rhodobacter capsulatus* SB1003 when cloned into pBBR1MCS5(Gm^R). This activity is independent of the orientation in which the fragment is cloned suggesting that the gene(s) responsible for the observed phenotype is cloned with its endogenous promoter 176

Figure 5-10: Sequence region of the pP3 insert that contains genes predicted to encode a putative PQQ containing protein (bold red), a permease of the drug/metabolite transporter superfamily (bold blue) an acetyltransferase of the GNAT family (bold pink), and a permease with

no homolog in *Rhodobacter* (bold green). Candidate promoters for these genes are highlighted in the color representative of the gene sequence177

Figure 6-1: A. Heme stain of soluble (S) and membrane (M) proteins of SW2 cells grown phototrophically on Fe(II), H₂, and acetate, and *Rhodobacter capsulatus* SB1003 grown photoheterotrophically on RCV. The arrow highlights a *c*-type cytochrome of approximately 15 kDa that appears unique to the membrane fraction of Fe(II)-grown SW2 cells. B. Total soluble and membrane protein profiles of SW2 grown on Fe(II), H₂, and acetate. The red dots highlight proteins that appear unique to the membrane fraction of Fe(II)-grown cells and the arrow identifies the protein that corresponds to the heme in part A..... 193

Figure 6-2: A. Heme stain of soluble (S) and membrane (M) proteins of TIE-1 cells grown on Fe(II), H₂, and thiosulfate, separated by SDS-PAGE. The black arrows highlight *c*-type cytochromes of approximately 45 kDa that are much more highly expressed in the crude and soluble fractions of Fe(II)-grown cells. In addition, there is a *c*-type cytochrome of approximately 90 kDa that is more highly expressed in the membrane fraction (indicted by the red arrow). B. Total soluble and membrane protein profiles of TIE-1 grown on Fe(II), H₂, and acetate. The red dots highlight proteins that appear unique to the soluble fraction of Fe(II)-grown cells 195

Figure 6-3: Gene cluster in *Rhodopseudomonas palustris* GCA009 with homolog in TIE-1, containing genes encoding proteins with possible function in photoautotrophic Fe(II) oxidation. 1: A gene predicted to encode a high redox potential Fe-S protein that is homologous to the *iro* (possible iron oxidase) in *Acidithiobacillus ferrooxidans*. 2: A gene predicted to encode an outer membrane protein, homologous to MtrA, (an outer membrane protein involved in Fe(III) respiration in *Shewanella oneidensis*. 3: The predicted product of this gene has an a multi-copper oxidase copper binding motif at the N-terminal and a C-terminal sequence is homologous to a deca-heme cytochrome *c* in *Shewanella*..... 198