PROTEIN ENGINEERING VIA SITE-SPECIFIC INCORPORATION OF NONNATURAL AMINO ACIDS

Thesis by

Inchan Kwon

In Partial Fulfillment of the Requirements

for the Degree of

Doctor of Philosophy

California Institute of Technology

Pasadena, California

2007

(Defended January 19, 2007)

© 2007

Inchan Kwon

All Rights Reserved

Acknowledgments

I would like to acknowledge my thesis advisor David Tirrell. I still remembered Dave's research presentation in my first year at Caltech. I was captivated by his clear presentation and fascinating research area, which influenced me to join his lab. During my days in his lab, I was deeply impressed by his endless curiosity in science and his systematic approach to solve problems with deep knowledge and strong enthusiasm. Without his timely advice and guidance throughout my PhD program, most of my projects would not have been successful. Dave was very supportive of my future career in all aspects.

I also would like to acknowledge other committee members, Prof. Frances Arnold, Prof. Dennis Dougherty, and Prof. Christina Smolke. They also served as my candidacy committee and provided me with many comments and much help since then. I appreciate their time and efforts to review my thesis as well as their great support for my career development.

I am grateful to all former and current Tirrell laboratory members. I was very lucky to work with such wonderful colleagues. In particular, Pin, Jamie, and Rebecca helped me a lot to settle down when I joined Tirrell laboratory. They knew everything I needed and were always willing to assist me. Three postdoctoral fellows, Kent, Jinsang, and Jin impressed me deeply with their commitment and enthusiasm to pursue science. They were also willing to share their experience and helped me to go through many non-scientific issues. They were invaluable role models as young scientists. I thank Caglar and Kimberly for their efforts and time reviewing my writing many times as well as for their helpful

... 111 discussions on diverse topics. I appreciate Julie and Jamie for sharing their experience in career development.

Finally, I wish to thank my wife Junghi and my son Hyungju for their continual support and encouragement throughout my PhD program. My daughter Suah, who was born recently, has added great pleasure to my life.

Abstract

Nonnatural amino acid incorporation has been one of the most important protein engineering techniques. Particularly site-specific incorporation of nonnatural amino acids would allow design of artificial proteins containing a nonnatural amino acid with minimal perturbation of their native properties. Site-specific incorporation of phenylalanine (Phe) analogs and tryptophan (Trp) analogs, such as *p*-bromophenylalanine (pBrF), *p*iodophenylalanine, p-azidophenylalanine, 6-chlorotrytophan, 6-bromotryptophan, 5bromotrytophan, and benzothienylalanine, into proteins in Escherichia coli has been realized by E. coli strains outfitted with a mutant yeast phenylalanyl-tRNA suppressor (ytRNA^{Phe}_{CUA}) and a mutant yeast phenylalanyl-tRNA synthetase (yPheRS (T415G)) with a point mutation in the active site of the enzyme. In order to reduce Trp and lysine (Lys) misincorporation at an amber codon, the ytRNA^{Phe}_{CUA_UG} containing the optimized sequence and the yPheRS (T415A) showing higher specificity toward pBrF were developed. Combining ytRNA^{Phe}_{CUA UG} and yPheRS (T415A) allowed incorporation of pBrF into murine dihydrofolate reductase in response to an amber codon with at least 98% fidelity.

Re-assignment of degenerate sense codons offers the prospect of a substantially expanded genetic code and a correspondingly enriched set of building blocks for natural and artificial proteins. Here we describe the use of a mutant yeast phenylalanine transfer RNA (ytRNA^{Phe}_{AAA}) containing a modified anticodon to break the degeneracy of the genetic code in *E. coli*. By using an *E. coli* strain co-transformed with ytRNA^{Phe}_{AAA} and a mutant yPheRS (T415G), we demonstrated efficient replacement of Phe by L-3-(2-naphthyl)alanine (2Nal) at UUU, but not at UUC codons.

However, this method had two limitations. First, the yPheRS (T415G) also activated Trp, which led to Trp misincorporation. Second, 2Nal was misincorporated at UUC Phe codons, due to the relaxed codon recognition of AAA anticodon in the ytRNA^{Phe}_{AAA}. High-throughput screening of a yPheRS library led to a more selective yPheRS variant (yPheRS_naph). The rationally designed ytRNA^{Phe}_{CAA}, which has the CAA anticodon recognizing only a UUG (Leu) codon, allowed incorporation of 2Nal only at UUG codon. Combined use of yPheRS_naph and ytRNA^{Phe}_{CAA} achieved multiple-site-specific incorporation of 2Nal into proteins. These results illustrate a general method for increasing the number of distinct, genetically-encoded amino acids available for protein engineering.

Table of Contents

Acknowled	gments	iii
Abstract		v
Table of Co	ontents	vii
List of Tab	les	xii
List of Figu	res	xiii
Chapter 1	Expansion of the Number of the Genetically Encoded Amino Acids in <i>E. coli</i>	1
Abstr	act	2
Introd	luction	2
Site-S	Specific Incorporation	3
Choic	e of a Nonnatural Amino Acid	4
Ortho	ogonal Pairs	5
1	Modification of Endogenous Orthogonal Pairs	5
(Orthogonal Pairs Derived from Yeast	6
(Orthogonal Pairs Derived from Archaebacteria	7
Substrate Specificity Change		9
Rational Design of the Active Sites of Aminoacyl-tRNA Synthetases		9
Ι	Directed Evolution of Aminoacyl-tRNA Synthetase Substrate	12
New	Codon-Anticodon Interactions	14
1101	Nonsense Suppression	15
I	Breaking the Degeneracy of the Genetic Code	15
I	Frameshift Suppression	17
Conc	lusions	18
Refer	ences	19
Chapter 2	Site-Specific Incorporation of Phenylalanine Analogs into Proteins in Vivo by an Engineered Yeast Phenylalanyl-tRNA Synthetase	36
Abstr	act	37
Introd	luction	38
Materials and Methods		40
Ν	Materials	40
ł	Plasmid Construction for Synthetase Expression	40

	4.1
Synthetase Expression and Purification	41
Amino Acid Activation Assay	42
Plasmid and Strain Construction for in Vivo Incorporation Assay	42
Analog Incorporation Assay in Vivo	43
Composition Analysis of Mutant mDHFR	44
Results and Discussion	45
Rationales for Engineering the New Synthetase Specificity	
Synthetase Expression and Purification	46
Amino Acid Specificity of Mutant Yeast PheRS	47
Site-Specific Incorporation of Phe Analogs into mDHFR	48
in Vivo Using a Phenylalanine Auxotrophic Strain	
Site-specific Incorporation of Phe Analogs into mDHFR	50
in Vivo Using a Phe/Trp Double Auxotrophic Strain	
Site-specific Incorporation of Phe Analogs into mDHFR	51
in Vivo Using a Phe/Trp/Lys Triple Auxotrophic Strain.	-
Conclusions	52
Acknowledgments	53
References	53
	00
Chanter 3 Design of a Bacterial Host for Site-Specific Incorporation of	68
<i>n</i> -Bromonhenvlalanine into Recombinant Proteins	00
<i>p</i> -bromophenylatannie into recombinant i roteiny	
Abstract	69
Introduction	69
Materials and Methods	
Materials	
Plasmid Construction for Synthetase Expression	72
Synthetase Expression and Purification	73
Amino Acid Activation Assav	73
Plasmid Construction for vtRNA ^{Phe} Expression	74
In Vitro Transcription	75
Aminoacylation Assay	76
Strain and Plasmid Construction for in Vivo Incorporation of pBrF	77
In Vivo Incorporation of pBrF	77
Quantitative Analysis of pBrF Incorporation by Liquid	78
Chromatography Mass Spectrometry (I C-MS)	70
Results and Discussion	79
A minoacylation of vtRNA ^{Phe} and vtRNA ^{Phe} we with I vs	79
$\Omega_{\text{ccupancy of the Amber Site}}$	80
Reduced Lys Misincorporation by $ytPNA^{Phe}$	81
Padasian of Donulalanul tPNA Sunthatasa	01 Q1
ATD DDi Eystenge and Amineseylation of utDNA ^{Phe} Catalyzed	01
ATP-PPI Exchange and Annhoacylation of ytkinA Catalyzed	× /
by yDboDS Vorients	02
by yPheRS Variants	02
by yPheRS Variants High Fidelity Site-Specific Incorporation of pBrF	83

Acknowledgments	
References	84
Chapter 4 Site-specific Incorporation of Tryptophan Analogs in	to
Recombinant Proteins in Vivo	96
Abstract	97
Introduction	98
Materials and Methods	102
Materials	102
Amino Acid Activation Assay	102
Strain and Plasmid Construction for in Vivo Incorporation	Assays 103
In Vivo Analog Incorporation Assays	104
Quantitative Analysis of Analog Incorporation by Liquid Chromatography Mass Spectrometry (LC-MS)	105
Fluorescent Protein Isolation	105
Characterization of Spectral Properties	106
Results and Discussion	107
Rational Design of yPheRS and Choice of Trp Analogs	107
Activation of Trp Analogs by yPheRS (T415G)	108
Site-Specific Incorporation of Trp Analogs into mDHFR i	n Vivo 109
Spectral Analysis of Fluorescent Proteins	111
Conclusions	113
Acknowledgments	113
References	114
Chapter 5 Breaking the Degeneracy of the Genetic Code	127
Abstract	128
Introduction	128
Materials and Methods	129
Materials	129
Strains and Plasmids	129
Construction of Plasmid Carrying Mutant Yeast tRNA	129
Construction of Plasmid Carrying mDHFR and Mutant Ye Phenylalanyl-tRNA Synthetase	east 130
Protein Expression and Purification	130
Amino Acid Analysis	131
MALDI-MS Analysis	131
Results and Discussion	132
Conclusions	137
Acknowledgments	137
References	137

ix

Chapter 6	Breaking the Degeneracy of the Leucine Codons in <i>Escherichia coli</i>	14
Abstr	act	14
Intro	duction	14
Mate	rials and Methods	15
]	Materials	15
]	Preparation of <i>E. coli</i> hosts	15
1	Plasmid Construction for Reporter Gene Expression	15
(Construction of vPheRS Library	15
(Construction of Expression Library	15
S	Screening of vPheRS Library	15
(Characterization of the Isolated Clones	15
	Amino Acid Activation Assav	15
(Construction of Plasmids and Expression Hosts for Incorporation	15
1	Residue- and Site-Specific Incorporation of 2Nal into	15
(Construction of Plasmids and Expression Hosts for 2Nal	15
-	Incorporation at Leu Codons	
l	Expression of mDHFR Variants and GFP Variants in Vivo	16
	Quantitative Analysis of Codon Occupancy	16
Resu	Its and Discussion	16
I	to UUU Codons	16
(Construction of yPheRS Library	16
l	High-throughput Screening of yPheRS library	16
(Characterization of the Selective yPheRS Variant	16
1	Amino Acid Activation Analysis by yPheRS_naph	16
]	Elimination of Trp Misincorporation at UUU Codons by vPheRS naph in Vivo	16
1	Residue- and Single-Site-Specific Incorporation of 2Nal in Vivo	16
	Single-Site-Specific Incorporation of 2Nal into GFP in Vivo	16
Ī	Misincorporation of 2Nal at Unwanted Sites (UUC Codons) in Vivo	16
	Breaking the Degeneracy of the Leucine Codons	16
	UUG Codon-Specific Incorporation of 2Nal	17
ו	Minimal Perturbation of Native Properties of GFP upon	17
1	Incorporation of 2Nal	17
Conc	lusions	17
Ackn	owledgments	17
Refer	rences	17
Appendix A	A Plasmid Information	18
		~

Appendix C Names and Sequences of PCR Primers

215

List of Tables		Page	
Table 2.1	Kinetic parameters for ATP-PPi exchange of amino acids (1-7) by the mutant yeast PheRS	61	
Table 3.1	Occupancy of amber sites and expression yields	93	
Table 3.2	Kinetic parameters for ATP-PPi exchange of amino acids by the wild-type yPheRS, yPheRS (T415G), and yPheRS (T415A) variant	94	
Table 4.1	Kinetic parameters for ATP-PPi exchange by the yPheRS (T415G)	124	
Table 4.2	Occupancy of amber sites and mDHFR expression yields	125	
Table 4.3	Spectral properties of CFP6 variants	126	
Table 5.1	Molar masses and numbers of 2Nal residues observed for mDHFR samples prepared under various conditions	140	
Table 6.1	Kinetic parameters for ATP-PPi exchange of amino acids by the yPheRS (T415G) and yPheRS_naph variants	187	
Table 6.2	Occupancy of UUU and UAG codons by various amino acids	188	

List of Figures		Page	
Figure 1.1	Nonnatural amino acids that have been genetically incorporated into proteins in bacteria, yeast or mammalian cells.	28	
Figure 1.2	Structures of yeast tRNA ^{Phe} and <i>E. coli</i> tRNA ^{Phe} .	29	
Figure 1.3	Anticodon-loop tRNA library, all-loop tRNA library derived from <i>M. jannaschii</i> tRNA ^{Tyr} _{CUA} and a general selection scheme for suppressor tRNAs that are orthogonal to <i>E. coli</i> endogenous systems and charged efficiently by a cognate synthetase.	30	
Figure 1.4	Crystal structure of the <i>T. thermophilus</i> PheRS (<i>t</i> PheRS, pdb 1B70) in ribbon model and sequence alignment of PheRS from 21 different organisms.	31	
Figure 1.5	A Phe substrate and four residues within 7 Å of the <i>para</i> -position of phenyl ring of a Phe substrate inside the binding pocket of a homology model of yPheRS structure, and a screening scheme of yPheRS library.	33	
Figure 1.6	A schematic diagram describing the concept of breaking the degeneracy of the genetic code.	35	
Figure 2.1	Sequence alignment of PheRS variants from <i>Thermus thermophilus Escherichia coli</i> , <i>Saccharomyces cerevisiae</i> and stereoview of active site of PheRS from <i>T. thermophilus</i> .	s, 59	
Figure 2.2	 Amino acids involved in this study. (1) phenylalanine; (2) <i>p</i>-bromophenylalanine; (3) <i>p</i>-iodophenylalanine; (4) <i>p</i>-azidophenylalanine; (5) tyrosine; (6) trytophan; (7) 3-(2-naphthyl)alanine; (8) L-lysine. 	60	
Figure 2.3	Amino acid sequence of marker protein mDHFR (38Am).	62	
Figure 2.4	MALDI-MS of proteolytic peptide fragments derived from mDHFR (38Am) expressed in media supplemented with 7 or 2.	63	
Figure 2.5	The tandem mass spectrum of Peptide A (NGDLPWPPLRNE Z K) derived from mDHFR (38Am) expressed in AF strain in media supplemented with 7 .	65	
Figure 2.6	MALDI-MS of proteolytic peptide fragments of mDHFR (38Am) expressed in AFWK strain in media supplemented with 2 or 3 .	66	

xiii

Figure 3.1	Aminoacylation of $ytRNA_{CUA}^{Phe}$ and $ytRNA_{CUA_UG}^{Phe}$ with Lys by eLysRS.	88
Figure 3.2	LC-MS chromatograms of tryptic digests of mDHFR.	89
Figure 3.3	Charging of Phe and Trp by wild-type yPheRS, yPheRS (T415G), and yPheRS (T415A).	91
Figure 4.1	 Amino acids involved in this study: (1) tryptophan, (2) phenylalanine, (3) 6-chlorotryptophan, (4) 6-bromotryptophan, (5) 5-bromotryptophan, (6) benzothienylalanine. 	118
Figure 4.2	A hypothetical model for the adaptability of Trp analogs to the binding pocket of eTrpRS.	119
Figure 4.3	SDS-PAGE analysis of mDHFR_38Am expression.	120
Figure 4.4	Tandem mass spectrum of the peptide (NGDLPWPPLRNEZK).	121
Figure 4.5	Crystal structure of enhanced cyan fluorescent protein and chromophores of CFP6 variants containing 1, 3, 4, and 6.	122
Figure 4.6	Absorption and fluorescence emission spectra for CFP6 variants containing different Trp analogs at the Trp66 position.	123
Figure 5.1	A strategy for multi-site incorporation of 2Nal into recombinant proteins by breaking the degeneracy of the phenylalanine codons.	141
Figure 5.2	SDS-PAGE analysis of mDHFR prepared in minimal media supplemented with 3 mM 2Nal and free of exogenous Phe.	142
Figure 5.3	Replacement of Phe by 2Nal can be detected in MALDI-MS spectra of tryptic fragments of mDHFR.	143
Figure 5.4	Tandem mass spectrum of Peptide 1 _{UUU} (Nal) YKF*EVYEK.	145
Figure 6.1	LC-MS chromatogram of tryptic digests of mDHFR.	176
Figure 6.2	Phe substrate and four residues within 7 Å of <i>para</i> -position of the phenyl ring of the substrate inside the binding pocket of a homology model of yPheRS structure.	178
Figure 6.3	A screening scheme for yPheRS library.	179

Figure 6.4	Fluorescence intensities of cells containing GFP6 expressed in minimal medium without or with 2Nal.	180
Figure 6.5	Fluorescence intensities of cells under uninduced conditions.	181
Figure 6.6	UUC and UUU codon occupancy by Phe and 2Nal.	182
Figure 6.7	Fluorescence intensities of cells expressing GFP3_WC and GFP3 expressed in minimal medium without or with 2Nal.	183
Figure 6.8	LC-MS chromatogram of tryptic digests of mDHFR.	184
Figure 6.9	Fluorescence intensities of cells containing GFP3 (158UUG) expressed in minimal medium without or with 2Nal.	186

Chapter 1

Expansion of the Number of the Genetically Encoded Amino Acids in E. coli

Abstract

Nonnatural amino acid incorporation has been one of the most important protein engineering platforms. In particular, site-specific incorporation of nonnatural amino acids allows design of proteins containing a nonnatural amino acid with minimal perturbation of native properties. Until now more than 30 nonnatural amino acids have been introduced into proteins in *E. coli* hosts outfitted with orthogonal pairs of cognate tRNAs and aminoacyl-tRNA synthetases (aaRS). Substrate specificity of aaRS has been altered to recognize various nonnatural amino acids through either rational design of the active site of aaRS or high-throughput screening of aaRS library. In this approach, site-specificity has been achieved by assigning a nonnatural amino acid to a stop codon, a degenerate codon, or a frameshift codon for efficient discrimination of the programmed sites from unwanted sites.

Introduction

Genetically encoded nonnatural amino acids have been used to endow recombinant proteins with novel chemical, physical, or biological properties. Several distinct methods have been developed to introduce nonnatural amino acids into recombinant proteins at programmed sites in vivo. Residue-specific incorporation involves the global replacement of a particular natural amino acid with a nonnatural amino acid in a target protein using auxotrophic hosts.¹⁻¹⁴ The strength of this technique lies in efficient protein translation and multi-site incorporation of a nonnatural amino acid, because sense codons are reassigned for a nonnatural amino acid. Residue-specific incorporation of 5',5',5'-trifluoroleucine (TFL) into a leucine-zipper protein enhanced its thermal stability.¹⁵ A reactive phenylalanine analog, *para*-ethynylphenylalanine, has been used for selective dye-labeling of newly synthesized protein in *E. coli*.¹⁶ Newly synthesized proteins in mammalin cells have been selectively indentified using bioorthogonal noncanonical amino acid tagging.¹⁷ Trp analogs containing selenophene and thienyl functional groups have been used in the X-ray crystallography of proteins.^{18,19} Aminotryptophan was used to design protein-based pH sensors³ and greatly changed the spectral properties of fluorescent proteins.¹ Regardless of many successful applications of residue-specific incorporation, its further application to the design of biologically-active macromolecules could be restricted by two limitations. First, reassignment of sense codons for a nonnatural amino acid usually leads to exclusion of one natural amino acid, which is encoded by the sense codons, in protein translation. Second, native properties of a target protein may be impaired by incorporation of a nonnatural amino acid at non-permissive sites of a target protein,^{20,21} though the Tirrell laboratory recently showed that the impaired properties may be rescued through directed evolution of the target protein.²¹

Site-Specific Incorporation

Site-specific incorporation involves introduction of a nonnatural amino acid into a target protein at any position selected. Access to all 20 natural amino acids and site-specificity of the method may allow design of proteins containing a nonnatural amino acid with minimal perturbation of native properties. Site-specific incorporation requires outfitting the cell with an "orthogonal pair" comprising a suppressor tRNA and a cognate aminoacyl-tRNA synthetase (aaRS) that operate independently of the endogenous synthetase-tRNA pairs in *E. coli*. The site-specific incorporation of a nonnatural amino acid into recombinant proteins via this strategy was reported by Furter in 1998²² and the Schultz

laboratory has developed powerful selection methods to identify heterologous synthetases and tRNAs.²³⁻²⁶ A general strategy for site-specific incorporation consists of four steps and each step will be discussed in more detail in the following sections.

Choice of a Nonnatural Amino Acid

First of all, it is necessary to choose a nonnatural amino acid suitable for goals. Until now more than 30 nonnatural amino acids have been successfully introduced to proteins in a site-specific manner^{27,28} (Figure 1) including reactive nonnatural amino acids, such as *para*-acetylphenylalanine 1, *meta*-acetylphenylalanine 2, *para*-(3-oxobutanoyl)-Lphenylalanine 3. para-(2-amino-3-hydroxylenthynyl)phenylalanine 4. paraehthynylthiocarbonyl-phenylalanine 5, *para*-propargyloxyphenylalanine 6, paraazidophenylalanine 7, para-ethynylphenylalanine 32, para-iodophenylalanine 20, and para-bromophenylalanine 31. These reactive nonnatural amino acids can be used to modify proteins through bio-orthogonal chemical transformations. For example, the keto and β -diketo functional groups (1-3) can selectively react with both hydrazides and alkoxyamines.²⁹⁻³¹ Azide functional groups (7) can be ligated to alkynes through coppercatalyzed [3+2] cycloaddition.^{16,32,33} Aryl halides (20 and 31) can be conjugated to terminal alkynes or alkenes via palladium-catalyzed coupling reactions.³⁴⁻³⁹ Nonnatural amino acids containing photoactive side chains including *para*-azidophenylalanine $7^{40,41}$ parabenzoylphenylalanine 9, 42,43 O-(2-nitrobenzyl)tyrosine 12, 44 and S-(2-nitrobenzyl)cysteine 10^{45} have been inserted into proteins.

Genetically-encoded nonnatural amino acids can be used as probes of various kinds. First, nonnatural amino acids containing heavy atoms (I and Se) have been used for X-ray crystallographic studies of protein structure.^{18,46} Second, fluorinated phenylalanine and ¹⁵Nlabelled methoxy phenylalanine have been used to label proteins for NMR study.^{22,47} Third, para-L-cyanophenylalanine 22 containing nitrile group has been introduced to proteins as an infrared probe to investigate changes in local environment.⁴⁸ Fourth, introduction of a fluorophore into proteins at defined sites can greatly facilitate study of protein function and structure. For example, the two fluorescent amino acids. 2-amino-3-(5-(dimethylamino)naphthalene-1-sulfonamide)propanoic acid (dansylalanine) 18 and L-(7hydroxycoumarin-4-yl)ethylglycine 17, have been introduced to proteins to monitor unfolding of proteins.^{49,50}

Orthogonal Pairs

Activation of an amino acid followed by charging the activated amino acid into a cognate tRNA by a cognate aminoacyl-tRNA synthetase (aaRS) is a key step to ensure high fidelity in a protein translation process. One orthogonal pair of aaRS/tRNA is assigned to each natural amino acid. Therefore, a new orthogonal aaRS/tRNA pair is necessary for a nonnatural amino acid to be utilized in protein translation machinery. Therefore, the development of orthogonal pairs to each living organism is one of the key steps in achieving site-specific incorporation of a nonnatural amino acid into proteins in vivo.

Modification of Endogenous Orthogonal Pairs. Initially Schultz and colleagues tried to modify an *E. coli* endogenous RNA/aaRS pair to obtain a new orthogonal pair. Three sites in $tRNA^{Gln}_{2}$ sequence, which are critical for binding with *E. coli* glutamyl-tRNA synthetase (eGlnRS), were mutated to reduce affinity to eGlnRS. Then eGlnRS was evolved to

5

recognize the mutant $tRNA^{Gln}_2$ and retain its orthogonality with the endogenous tRNAs in *E. coli*. However, the approach was unsuccessful due to poor discrimination of the mutant $tRNA^{Gln}_2$ against wild-type $tRNA^{Gln}_2$ by even the best eGlnRS mutant.⁵¹

Orthogonal Pairs Derived from Yeast. A second approach to generate an orthogonal pair was the use of a heterologous tRNA/aaRS pair obtained from another organism in E. coli. The yeast phenylalanyl-tRNA/yeast phenylalanyl-tRNA synthetase pair was known to be orthogonal to the E. coli endogenous system. In vitro aminoacylation studies indicated that there was little cross-talk between E. coli and yeast phenylalanyl-tRNA/synthetase pairs.^{52,53} Aminoacylation activity of yeast tRNA^{Phe} by *E. coli* PheRS is 10-fold lower than that of E. coli tRNA^{Phe. 22} Aminoacylation activity of E. coli tRNA^{Phe} by yeast PheRS is 24or 34-fold lower than that of yeast tRNA^{Phe},^{54,55} because U20 in *E. coli* tRNA^{Phe} replaces G20, one of the identity elements of yeast tRNA^{Phe} recognized by yPheRS (Figure 2).⁵⁴ A yeast amber suppressor derived from tRNA^{Phe}_{GAA} (ytRNA^{Phe}_{CUA wt}) has been widely used to study suppression of amber codons, though E. coli lysyl-tRNA synthetase (eLysRS) shows a weak activity to aminoacylate ytRNA^{Phe}_{CUA_wt} with Lys.²² A wild-type tRNA sequence can be engineered to eliminate the cross-reactivity with E.coli synthetase but preserve efficient recognition by the cognate aaRS. When ytRNA^{Phe}_{CUA wt} was coexpressed with wild-type yPheRS in an E. coli host, 60% of the amber codon was occupied by Lys.²² The mischarging by Lys was likely due to the presence of U36 and A73 in the ytRNA^{Phe}_{CUA wt}, which are the main identity elements for eLysRS.⁵⁶ However, when G37 was replaced by A37 and ytRNA^{Phe}_{CUA} was co-expressed with wild-type yPheRS, only 5% of the amber codon was occupied by Lys. The remaining 5% of Lys misincorporation was

eliminated by rational design of ytRNA sequence. Disruption of a Watson-Crick base pair between nucleotides 30 and 40 in ytRNA^{Phe}_{CUA} reduced mischarging by the eLysRS. As an alternative orthogonal tRNA/synthetase pair, an amber suppressor derived from yeast tyrosyl-tRNA (ytRNA^{Tyr}) and yeast tyrosyl-tRNA synthetase (yTyrRS) pair was examined.^{57,58} RajBhandary and colleagues generated yTyrRS variants using error-prone PCR to enhance discrimination of the suppressor tRNA over E. coli proline tRNA by factors of 2.2- to 6.8-fold. Nishikawa and colleagues optimized the sequence of yeast amber suppressor tRNA^{Tyr} to minimize Lys charging by eLysRS.⁵⁹ This optimized yeast tRNA^{Tyr}, together with yeast TyrRS, could be used as an orthogonal pair for incorporation of nonnatural amino acids into proteins. Schultz and colleagues developed two orthogonal pairs derived from yeast. An amber suppressor derived from yeast glutamyl-tRNAs (ytRNA^{Gln}_{2 CUA}) is not a substrate for any *E. coli* endogenous synthetase. In turn, yeast glutamyl-tRNA synthetase (yGlnRS) aminoacylated ytRNA^{Gln}₂ in *E. coli* but did not charge E. coli tRNA^{Gln 60}. An amber suppressor derived from yeast aspartidyl-tRNA (ytRNA^{Asp}_{CUA}) is not recognized by *E. coli* synthetases. Introduction of E188K mutation in yeast aspartidyl-tRNA synthetase (yAspRS) led to reduction in aminoacylation activity *coli* tRNA^{Asp 61} Therefore, both ytRNA^{Gln}_{2 CUA}/yGlnRS pair with E. and ytRNA^{Asp}_{CUA}/yAspRS (E188K) form orthogonal pairs in *E. coli* suitable for the incorporation of nonnatural amino acids into proteins in vivo.

Orthogonal Pairs Derived from Archaebacteria. Schultz and colleagues developed an orthogonal tyrosyl-tRNA/synthetase pair obtained from archaea, since archeal aminoacyl-tRNA synthetases are more similar to eukaryotic than prokaryotic counterparts. The amber

suppressor tyrosyl-tRNA (MjtRNA^{Tyr})/synthetase (MjTyrRS) pair derived from Methanococcus jannaschii was selected and evolved to be orthogonal to the E. coli endogenous system.^{25,62} The amber suppressor *MitRNA*^{Tyr}_{CUA} was efficiently aminoacylated by MiTyrRS in E. coli. However, E. coli endogenous synthetases showed weak activity towards M_{jt} RNA^{Tyr}_{CUA}. Therefore, a selection method to obtain orthogonal tRNAs in E. coli has been developed to enhance orthogonality with respect to E. coli endogenous synthetases and retain recognition by the cognate synthetase and the protein translational machinery (Figure 3).²⁵ This method includes positive and negative selection of a mutant tRNA library. In the negative selection, amber codons were introduced into a toxic barnase protein at permissive sites that allow replacement of amino acids. Eleven nucleotides of *MitRNA*^{Tyr}_{CUA} that do not bind to *Mi*TyrRS were saturated with all four bases to generate a tRNA library. When the tRNA library was expressed in minimal medium without M/TyrRS, cells containing tRNA variants aminoacylated by E. coli endogenous synthetases resulted in cell death. The plasmids were isolated from the cells survived, and co-transformed with a plasmid harboring both a M_I TyrRS gene and a β lactamase gene containing an amber codon at a permissive site into E. coli hosts. When cells were cultured in liquid medium in the presence of ampicillin, cells harboring tRNA variants aminoacylated by MiTyrRS survived. After the negative and positive screening of tRNA library, the tRNAs were orthogonal to the E. coli endogenous synthetases and retained their affinity to the cognate synthetase. Schultz and colleagues developed another orthogonal pair obtained from Archea. An amber suppressor leucyl-tRNA and a leucyltRNA synthetase were derived from Halobacterium sp. NRS-1 and Methanobacterium thermoautotrophicum, respectively. Aminoacylation activity of the suppressor leucyltRNA by the leucyl-tRNA synthetase and its orthogonality with *E. coli* synthetases were significantly enhanced by extensive mutations in the anticodon loop and acceptor stem.²³

Substrate Specificity Change

Another issue to achieve incorporation of a nonnatural amino acid into protein in vivo is alteration of substrate specificity of an aminoacyl-tRNA synthetase toward the nonnatural amino acid. Aminoacyl-tRNA synthetases ensure the high fidelity of transforming genetic code sequences into biologically functional proteins through a twostep aminoacylation reaction.⁶³⁻⁶⁷ In the first step, the cognate amino acid is activated by aaRS in the presence of ATP to form the amino acid adenylate. Subsequently aaRS catalyzes the esterification reaction to join the amino acid to the 2'- or 3'-OH of the terminal ribonucleotide of its cognate tRNA. Manipulation of such reactions could potentially alter the genetic code to allow incorporation of novel amino acid into proteins in vivo.⁶⁸ Some nonnatural amino acids including TFL, azidohomoalanine, and pararecognized fluorophenylalanine, are by wild-type cognate aminoacyl-tRNA synthetases.^{5,15,22} However, the majority of nonnatural amino acids are very poor substrates for natural aminoacyl-tRNA synthetases. Engineering the new synthetase activity is almost inevitable when one wants to incorporate an amino acid that is not recognized by wild-type enzymes.

Rational Design of the Active Sites of Aminoacyl-tRNA Synthetases. Alteration of substrate specificity of aminoacyl-tRNA synthetases has been achieved in several labs by the rational design of the binding pocket to accommodate nonnatural amino acid. Tirrell

and colleagues have been interested in exploring the possibility of incorporating substituted *p*-bromophenylalanine, phenylalanines, such *p*-iodophenylalanine, as and pazidophenylalanine. On the basis of the crystal structure of the PheRS (tPheRS) from Thermus thermophilus (Figure 4a and b), Safro and colleagues proposed that Val 261 and Ala 314 in the amino acid binding pocket of α -subunit of the tPheRS are critical in the discrimination of Phe from its amino acid competitors. Sequence alignment indicates that Ala 314 in α -subunit of the tPheRS corresponds to Ala 294 in α -subunit of the ePheRS (Figure 4c), and the Hennecke group showed that the substrate specificity of the ePheRS can be relaxed by a mutation at Ala 294. The A294G mutant was shown to enable incorporation of *para*-chlorophenylalanine into recombinant proteins.⁶⁹ A subsequent computational simulation, consistent with the Safro prediction, identified two cavityforming mutations (T251G and A294G) in ePheRS binding pocket. These two mutations led to relaxed substrate specificity and efficient in vivo replacement of Phe by paraacetylphenylalanine (pAcF).⁴ Sequence alignment showed that Thr 415 in yPheRS is equivalent to Thr 251 in ePheRS. Therefore, a T415G mutation was introduced into the yPheRS to enlarge the active site. The yPheRS (T415G) variant showed activity for pbromophenylalanine, p-iodophenylalanine, and p-azidophenylalanine in ATP-PPi exchange assays in vitro (see Chapters 2 and 4). There was an interesting finding that the yPheRS (T415G) efficiently activated Trp and 2-naphthylalanine as well as Phe analogs. Considering the spectrum of nonnatural amino acids recognized by yPheRS (T415G), Tirrell and colleagues explored whether the yPheRS (T415G) can activate even Trp analogs. In vitro activation studies showed that the vPheRS (T415G) efficiently activated four Trp analogs, 6-chlorotryptophan (6ClW), 6-bromotryptophan (6BrW), and 5bromotrytophan (5BrW), which were not utilized by the endogenous *E. coli* translational system (see Chapter 3). A phenylalanine auxotrophic *E. coli* strain transformed with this yPheRS (T415G) and cognate suppressor tRNA enabled the assignment of an amber nonsense codon to Phe analogs and Trp analogs in vivo (see Chapters 2-4). The cavity formed due to the T415G mutation in the binding pocket of yPheRS led to the activation of Trp as well as nonnatural amino acids, which resulted in misincorporation of Trp at the programmed sites. Therefore, the binding site of yPheRS was re-designed to enhance specificity for pBrF. Specifically, Tirrell and colleagues used the T415A variant, which exhibits 5-fold higher activity towards pBrF as compared to Trp in ATP-PPi exchange assays. Use of the yPheRS (T415A) eliminated misincorporation of Trp at programmed sites in proteins.⁷⁰

Nishikawa and colleagues have changed substrate specificity of yeast tyrosyl-tRNA synthetase (yTyrRS) to accommodate tyrosine analogs.⁷¹ Based on the crystal structure of TyrRS obtained from *Bacillus stearothermophilus* TyrRS and sequence alignment of two homologous TyrRSs, active site residues in yTyrRS have been mutated to generate a series of yTyrRS variants. Among the yTyrRS variants, one containing the Tyr43Gly mutation (yTyrRS (Y43G)) was found to show activity for several 3-substituted tyrosine analogs in aminoacylation assays in vitro. yTyrRS (Y43G) efficiently charged ytRNA^{Tyr} with 3-iodotyrosine, which was not utilized by wild-type yTyrRS. The yTyrRS (Y43G) showed 400-fold lower activity towards tyrosine than wild-type yTyrRS. Similarly, Yokoyama and colleagues⁷² screened *E. coli* TyrRS variants recognizing 3-iodotyrosine via in vitro biochemical assays. The *E. coli* TyrRS variant that has two mutations (Y37V and Q195C) in the active site activated 3-iodotyrosine 10-fold more efficiently than tyrosine.⁷²

Combined use of *E. coli* TyrRS (V37C195) and a suppressor tRNA derived from *Bacillus stearothermophilus* allowed incorporation of 3-iodotyrosine into proteins in mammalian cells.⁷³

Directed Evolution of Aminoacyl-tRNA Synthetase Substrate Specificity. AminoacyltRNA synthetases are known to be readily evolvable. Schultz and colleagues have developed powerful selection methods to change the substrate specificity of tyrosyl-tRNA synthetase (M/TyrRS) derived from Methanococcus jannaschii toward nonnatural amino acids.^{23,24,26,74} The selection method consists of a series of positive and negative selections.⁶⁸ Based on the crystal structure of the homologous TyrRS from *Bacillus* stearothermophilus, five residues in the active site of MiTyrRS were selected and randomized to generate a M_i TyrRS mutant library. In the positive selection, an amber codon was inserted at permissive sites in a chloramphenicol acetyltransferase (CAT) gene. The library was cultured in medium containing chloramphenicol in the presence of a nonnatural amino acid. Only the cells containing *Mi*TyrRS variants allowing suppression of an amber codon in CAT gene survived. Surviving cells were subjected to the negative selection. In the negative selection, the cells were regrown in medium containing chloramphenicol in the absence of the nonnatural amino acid. Those cells that could not survive were isolated from the replica plates containing the nonnatural amino acid. Using this method, substrate specificity of *Mi*TyrRS was changed to selectively recognize *O*methyl-*L*-tyrosine.⁶⁸

Alternative selection methods for evolving substrate specificity of *Mj*TyrRS towards nonnatural amino acids have been developed. One of them involves a positive

selection similar to the previous one, but a different negative selection.⁶⁰ The negative selection is based on suppression of amber codons inserted at permissive sites in the barnase gene that is lethal to E. coli. When the cells survived from the positive selection were cultured in the absence of nonnatural amino acids, the cells containing M/TyrRS variants recognizing natural amino acids resulted in cell death due to expression of the barnase protein. Another method for evolving *Mi*TyrRS specificity to nonnatural amino acids involves an amplifiable fluorescent reporter, a green fluorescent protein.²⁴ Amber codons were introduced at various sites in a T7 RNA polymerase gene that controlled expression of a green fluorescent protein. In the positive screening, the M/TyrRS library cells were cultured in the presence of a nonnatural amino acid. The cells containing M_i TyrRS variants that recognized the nonnatural amino acid and suppressed the amber codons in T7 RNA polymerase gene became fluorescent and were collected with a fluorescence-activated cell sorter (FACS). In the negative screening, the cells collected from the positive screening were cultured in the absence of the nonnatural amino acid; the cells containing MiTyrRS variants that did not recognize any natural amino acid were not fluorescent and collected with a FACS. Therefore, the cells collected from the positive and negative screenings contained the M_i TyrRS variants that can selectively charge the nonnatural amino acid into the cognate tRNA. One advantage of the latter method is that both reporter genes are encoded in a single plasmid, which eliminates the need for DNA isolation after each selection step. However, the method requires a FACS for highthroughput screening.

Most attempts to evolve aminoacyl-tRNA synthetase substrate specificity have been limited to prokaryotic or archaeal aminoacyl-tRNA synthetases. Recently, Tirrell and colleagues have evolved a eukaryotic aminoacyl-tRNA synthetase, yPheRS, to change its substrate specificity towards a hydrophobic phenylalanine analog, 2-naphthylalanine (see Chapter 6). The yPheRS library was constructed by randomizing four residues (N412, T415, S418, and S437) that contact the Phe substrate inside the binding pocket (Figure 5a). The screening consisted of a series of positive and negative screenings (Figure 5b), and involved expression of a GFP reporting folding status. The active synthetases would allow incorporation of either 2Nal or other natural amino acids at non-permissive sites of a GFP, which resulted in poorly-folded GFP showing weak fluorescence. Therefore, in the positive screening, the library cells were cultured in the presence of 2Nal and then the cells containing active synthetases were collected by high-throughput screening with a FACS. The yPheRS variants with very low activity toward natural amino acids will not misincorporate any natural amino acid at Phe sites in GFP and so GFP would retain full intensity of fluorescence. In the negative screening, the cells obtained from the positive screening were cultured in the absence of 2Nal and then the cells containing selective synthetases were collected with a FACS.

New Codon-Anticodon Interactions

Among the sixty-four codons, sixty-one (sense codons) are assigned to one of the twenty natural amino acids. The other three codons (nonsense codons) are stop codons that terminate protein translation. In order to incorporate a nonnatural amino acid into proteins in vivo, we need to either re-assign an existing codon or generate a new codon for the nonnatural amino acid. Until now three distinct approaches have been explored, reassignment of a stop codon, reassignment of a degenerate codon, and assignment of a frameshift codon.

Nonsense Suppression. In 1989 incorporation of a nonnatural amino acid into proteins in vitro was demonstrated by two independent groups, the Chamberlin group⁷⁵ and the Schultz group.⁷⁶ Nonsense suppression means that one of the stop codons (nonsense codons) can be read by a suppressor charged with an amino acid. Otherwise protein translation will be terminated at the stop codon. Three stop codons, amber (UAG),^{46,68,70,74,77} ochre (UAA),^{23,78,79} and opal (UGA) stop codons,^{23,78} have been reassigned for nonnatural amino acids. The amber codon has been used most widely,^{22,40,42,43,48,70,74,76,80-83} because it is the least common stop codon in E. coli, and because several naturally occurring suppressor tRNAs recognize it efficiently.^{84,85} Use of the amber codon to encode nonnatural amino acids requires outfitting the cell with an "orthogonal pair" comprising a suppressor tRNA and a cognate aminoacyl-tRNA synthetase (aaRS) that operate independently of the endogenous synthetase-tRNA pairs in E. coli as described in previous sections. To date, more than thirty nonnatural amino acids have been incorporated into proteins in response to the amber stop codon with high fidelity.27,28,70

Breaking the Degeneracy of the Genetic Code. Suppression of the amber stop codon has worked well for single-site insertion of novel amino acids. However, the utility of this approach is limited by moderate suppression efficiency. Competition between elongation and termination produces mixtures of full-length and truncated chains, and protein yields

are reduced accordingly. Efficient multi-site incorporation has been accomplished by replacement of natural amino acids in auxotrophic *E. coli* strains,^{1,6,9-11,86,87} and by using aminoacyl-tRNA synthetases with relaxed substrate specificity⁴ or attenuated editing activity.¹¹ Although this method provides efficient incorporation of analogs at multiple sites, it suffers from the limitation that one of the natural amino acids must be excluded from the engineered protein. Efficient multiple-site-specific incorporation can be achieved by re-assignment of a sense codon, if the sense codon can be efficiently distinguished from other sense codons. Tirrell and colleagues investigated re-assignment of a degenerate codon for a nonnatural amino acid. As a test case for establishing the feasibility of breaking the degeneracy of the code, phenylalanine codons were investigated (see Chapter 5).

Although 2Nal incorporation was biased to UUU codons in the first study, misincorporation of 2Nal at unwanted sites (UUC codons) could not be avoided due to ambiguity in codon recognition by the AAA anticodon of ytRNA^{Phe}_{AAA}. Misincorporation of a nonnatural amino acid at unwanted sites might cause perturbation or loss of function of a target protein.^{20,21} Based on experimental findings⁸⁸⁻⁹⁵ and stereo-chemical modeling⁹⁶⁻⁹⁹ of codon-anticodon interactions, an expanded wobble rule was proposed by Lim and Curran.⁹⁰ The expanded wobble rule indicates that A in the first position of the anticodon can recognize all four bases in the third position of the codon, although its affinity to C is lower than that to U. The observed incomplete discrimination of UUU from UUC codons by the AAA anticodon of ytRNA^{Phe} motivated us to explore a different set of degenerate codons. Several considerations recommend the UUG codon. First, Leu is encoded by six codons: UUA, UUG, CUA, CUG, CUU, and CUC. Discrimination of UUG from CUN (N = A/U/G/C) codons might be highly efficient due to discrimination at the first position in

the codon. Second, our existing yeast orthogonal pair should be readily adapted to the incorporation of Phe analogs in response to UUG codons. Third, according to the expanded wobble rules, C in the first position of the anticodon can recognize only G in the third position of the codon (see Chapter 6).

Frameshift Suppression. The utility of stop or degenerate codons might be restricted by limited number of codons. In order to overcome this limitation, extended codons have been investigated. Although three-base codons are universal for most living organisms, frameshift suppression of a four-base codon occurs is a regulatory mechanism in viruses and bacteria, and is mediated either by slipping of normal tRNAs at the ribosome or by natural frameshift suppressor tRNAs with four-base anticodons. Sisido and colleagues have pioneered frameshift suppression for incorporation of nonnatural amino acids into proteins in E. coli.¹⁰⁰⁻¹⁰³ In this approach, four-base codons were inserted into a target protein at programmed sites. When the four-base codons are read by a modified tRNA that is charged with nonnatural amino acid and contains the corresponding anticodon, the full length target protein can be synthesized. If the four-base codons are read as three-base codons, truncated proteins will be formed due to early termination of protein translation. Four-base codons were usually derived from rarely used codons to minimize reading as three-base codons by the endogenous tRNAs. The translation efficiencies of some four-base codons were higher than those of three-base stop codons. In particular, a GGGU codon showed 86% translation efficiency,¹⁰⁴ while the suppression efficiencies of an amber codon were ca. 20-60%.^{75,105} A number of four-base codons; CGGG, GGGU, CGGU, AGGU, CCCU, and CUCU have been generated.^{101-103,106-108} The four-base codon strategy has also been applied to

incorporate two different nonnatural amino acids into two programmed sites in a protein, which showed orthogonality of each four-base codon to other four-base codons as well as three-base codons.^{103,104,109} The frameshift suppression method has been extended to five-base codon to incorporate nonnatural amino acids.¹¹⁰ Recently, Sisido and colleagues have demonstrated frameshift suppression of a four-base codon (UAGN) in mammalian cells using tRNA containing NCUA anticodon.¹⁰⁸

Schultz and colleagues developed a combinatorial approach to select efficient suppressors of four-base codons; AGGA, UAGA, CCCU, and CUAG.^{23,111} This approach has been extended to five- and six-base codons with tRNAs containing 6-10 nucleotides in the anticodon loops.¹¹² They identified suppressors of several five-base codons, such as AGGAU, CUACU, and CUAGU. However, translation efficiency of five-base codons was ca. 10%, which was lower than that of four-base codons. Suppression of six-base codons was too weak to be detected.¹¹² Recently, Schultz and colleagues have shown that a four-base codon can be used to incorporate a nonnatural amino acid into proteins in *E. coli* outfitted with a heterologous tRNA/synthetase orthogonal pair. In particular, L-homoglutamine was charged into archeal tRNA^{Lys} by the evolved archeal lysyl-tRNA synthetase, and then was inserted into proteins in response to an AGGA codon in *E. coli*.¹¹³

Conclusions

In 1989 the concept of site-specific incorporation in vitro was independently demonstrated by the Chamberlin group and the Schultz group. Recently, several groups achieved site-specific incorporation of nonnatural amino acids into proteins in vivo using *E. coli* hosts outfitted with a twenty-first orthogonal pair of tRNA/aaRS. Schultz and

colleagues altered the substrate specificity of TyrRS, which was derived from *M. jannaschii*, to various nonnatural amino acids *via* high-throughput screening of a *Mj*TyrRS library. Tirrell and colleagues altered yPheRS substrate specificity toward Phe analogs, which led to site-specific incorporation of Phe analogs in *E. coli*. The moderate suppression efficiency of amber codons has motivated exploration of alternative codons that encode nonnatural amino acids. Breaking the degeneracy of the genetic code or frameshift suppression greatly expands the theoretical number of codons that can be assigned to nonnatural amino acids.

References

- Bae, J. H.; Rubini, M.; Jung, G.; Wiegand, G.; Seifert, M. H. J.; Azim, M. K.; Kim, J. S.; Zumbusch, A.; Holak, T. A.; Moroder, L.; Huber, R.; Budisa, N. *J. Mol. Biol.* 2003, *328*, 1071-1081.
- Budisa, N.; Minks, C.; Medrano, F. J.; Lutz, J.; Huber, R.; Moroder, L. Proc. Natl. Acad. Sci. U. S. A. 1998, 95, 455-459.
- Budisa, N.; Rubini, M.; Bae, J. H.; Weyher, E.; Wenger, W.; Golbik, R.; Huber, R.; Moroder, L. Angew. Chem. Int. Ed. 2002, 41, 4066-4069.
- Datta, D.; Wang, P.; Carrico, I. S.; Mayo, S. L.; Tirrell, D. A. J. Am. Chem. Soc. 2002, 124, 5652-5653.
- Kiick, K. L.; Saxon, E.; Tirrell, D. A.; Bertozzi, C. R. Proc. Natl. Acad. Sci. U. S. A.
 2002, 99, 19-24.
- 6. Kirshenbaum, K.; Carrico, I. S.; Tirrell, D. A. ChemBioChem 2002, 3, 235-237.

- Link, A. J.; Vink, M. K. S.; Agard, N. J.; Prescher, J. A.; Bertozzi, C. R.; Tirrell, D. A. Proc. Natl. Acad. Sci. U. S. A. 2006, 103, 10180-10185.
- 8. Pratt, E. A.; Ho, C. Fed. Proc. 1974, 33, 1463-1463.
- 9. Sharma, N.; Furter, R.; Kast, P.; Tirrell, D. A. FEBS Lett. 2000, 467, 37-40.
- 10. Tang, Y.; Tirrell, D. A. J. Am. Chem. Soc. 2001, 123, 11089-11090.
- 11. Tang, Y.; Tirrell, D. A. Biochemistry 2002, 41, 10635-10645.
- 12. Wang, P.; Tang, Y.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 6900-6906.
- 13. Link, A. J.; Mock, M. L.; Tirrell, D. A. Curr. Opin. Biotechnol. 2003, 14, 603-609.
- 14. Budisa, N. Angew. Chem. Int. Ed. 2004, 43, 6426-6463.
- Tang, Y.; Ghirlanda, G.; Petka, W. A.; Nakajima, T.; DeGrado, W. F.; Tirrell, D. A. Angew. Chem.Int. Ed. 2001, 40, 1494-1496.
- Beatty, K. E.; Xie, F.; Wang, Q.; Tirrell, D. A. J. Am. Chem. Soc. 2005, 127, 14150-14151.
- Dieterich, D. C.; Link, A. J.; Graumann, J.; Tirrell, D. A.; Schuman, E. M. Proc. Natl. Acad. Sci. U. S. A. 2006, 103, 9482-9487.
- Bae, J. H.; Alefelder, S.; Kaiser, J. T.; Friedrich, R.; Moroder, L.; Huber, R.; Budisa, N. J. Mol. Biol. 2001, 309, 925-936.
- Budisa, N.; Pal, P. P.; Alefelder, S.; Birle, P.; Krywcun, T.; Rubini, M.; Wenger, W.;
 Bae, J. H.; Steiner, T. *Biol. Chem.* 2004, *385*, 191-202.
- 20. Budisa, N.; Pal, P. P. Biol. Chem. 2004, 385, 893-904.
- 21. Montclare, J. K.; Tirrell, D. A. Angew. Chem. Int. Ed. 2006, 45, 4518-4521.
- 22. Furter, R. Protein Sci. 1998, 7, 419-426.
- 23. Anderson, J. C.; Schultz, P. G. *Biochemistry* **2003**, *42*, 9598-9608.

- 24. Santoro, S. W.; Wang, L.; Herberich, B.; King, D. S.; Schultz, P. G. *Nat. Biotechnol.*2002, 20, 1044-1048.
- 25. Wang, L.; Schultz, P. G. Chem. Biol. 2001, 8, 883-890.
- 26. Wang, L.; Schultz, P. G. Chem. Commun. 2002, 1-11.
- Wang, L.; Xie, J.; Schultz, P. G. Annu. Rev. Biophys. Biomol. Struct. 2006, 35, 225-249.
- 28. Xie, J. M.; Schultz, P. G. Curr. Opin. Chem. Biol. 2005, 9, 548-554.
- Wang, L.; Zhang, Z. W.; Brock, A.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A.
 2003, 100, 56-61.
- Zhang, Z. W.; Smith, B. A. C.; Wang, L.; Brock, A.; Cho, C.; Schultz, P. G. Biochemistry 2003, 42, 6735-6746.
- 31. Zeng, H. Q.; Me, J. M.; Schultz, P. G. Bioorg. Med. Chem. Lett. 2006, 16, 5356-5359.
- 32. Link, A. J.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 11164-11165.
- 33. Link, A. J.; Vink, M. K. S.; Tirrell, D. A. J. Am. Chem. Soc. 2004, 126, 10598-10602.
- Kodama, K.; Fukuzawa, S.; Nakayama, H.; Kigawa, T.; Sakamoto, K.; Yabuki, T.; Matsuda, N.; Shirouzu, M.; Takio, K.; Tachibana, K.; Yokoyama, S. *ChemBioChem* 2006, 7, 134-139.
- 35. Nguefack, J. F.; Bolitt, V.; Sinou, D. Tetrahedron Lett. 1996, 37, 5527-5530.
- Sakakibara, Y.; Ido, Y.; Sasaki, K.; Sakai, M.; Uchino, N. Bull. Chem. Soc. Jpn. 1993, 66, 2776-2778.
- 37. Schoenbe, A.; Heck, R. F. J. Org. Chem. 1974, 39, 3327-3331.
- 38. Yoneyama, M.; Kakimoto, M.; Imai, Y. Macromolecules 1988, 21, 1908-1911.
- Yoneyama, M.; Konishi, T.; Kakimoto, M. A.; Imai, Y. Makromol. Chem., Rapid Commun. 1990, 11, 381-386.
- Chin, J. W.; Santoro, S. W.; Martin, A. B.; King, D. S.; Wang, L.; Schultz, P. G. J. Am. Chem. Soc. 2002, 124, 9026-9027.
- 41. Wang, P., Ph.D. thesis, California Institute of Technology, Pasadena, 2003.
- Chin, J. W.; Martin, A. B.; King, D. S.; Wang, L.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 2002, 99, 11020-11024.
- 43. Chin, J. W.; Schultz, P. G. ChemBioChem 2002, 3, 1135-1137.
- 44. Deiters, A.; Groff, D.; Ryu, Y. H.; Xie, J. M.; Schultz, P. G. Angew. Chem. Int. Ed.
 2006, 45, 2728-2731.
- 45. Wu, N.; Deiters, A.; Cropp, T. A.; King, D.; Schultz, P. G. J. Am. Chem. Soc. 2004, 126, 14306-14307.
- 46. Xie, J. M.; Wang, L.; Wu, N.; Brock, A.; Spraggon, G.; Schultz, P. G. *Nat. Biotechnol.* **2004**, *22*, 1297-1301.
- 47. Deiters, A.; Geierstanger, B. H.; Schultz, P. G. ChemBioChem 2005, 6, 55-58.
- Schultz, K. C.; Supekova, L.; Ryu, Y. H.; Xie, J. M.; Perera, R.; Schultz, P. G. J. Am. Chem. Soc. 2006, 128, 13984-13985.
- Summerer, D.; Chen, S.; Wu, N.; Deiters, A.; Chin, J. W.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 2006, 103, 9785-9789.
- 50. Wang, J. Y.; Xie, J. M.; Schultz, P. G. J. Am. Chem. Soc. 2006, 128, 8738-8739.
- Liu, D. R.; Magliery, T. J.; Pasternak, M.; Schultz, P. G. *Proc. Natl. Acad. Sci. U. S. A.* 1997, 94, 10092-10097.
- 52. Doctor, B. P.; Mudd, J. A. J. Biol. Chem. 1963, 238, 3677-3681.

- 53. Kwok, Y.; Wong, J. T. F. Can. J. Biochem. 1980, 58, 213-218.
- Sampson, J. R.; Direnzo, A. B.; Behlen, L. S.; Uhlenbeck, O. C. Science 1989, 243, 1363-1366.
- 55. Roe, B.; Sirover, M.; Dudock, B. Biochemistry 1973, 12, 4146-4154.
- Tamura, K.; Himeno, H.; Asahara, H.; Hasegawa, T.; Shimizu, M. *Nucleic Acids Res.* 1992, 20, 2335-2339.
- Kowal, A. K.; Kohrer, C.; RajBhandary, U. L. Proc. Natl. Acad. Sci. U. S. A. 2001, 98, 2268-2273.
- Ohno, S.; Yokogawa, T.; Fujii, I.; Asahara, H.; Inokuchi, H.; Nishikawa, K. J. Biochem. (Tokyo). 1998, 124, 1065-1068.
- Fukunaga, J.; Yokogawa, T.; Ohno, S.; Nishikawa, K. J. Biochem. (Tokyo). 2006, 139, 689-696.
- 60. Liu, D. R.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 1999, 96, 4780-4785.
- 61. Pastrnak, M.; Magliery, T. J.; Schultz, P. G. Helv. Chim. Acta 2000, 83, 2277-2286.
- Wang, L.; Magliery, T. J.; Liu, D. R.; Schultz, P. G. J. Am. Chem. Soc. 2000, 122, 5010-5011.
- 63. Francklyn, C.; Perona, J. J.; Puetz, J.; Hou, Y. M. RNA 2002, 8, 1363-1372.
- 64. de Pouplana, L. R.; Schimmel, P. Trends Biochem. Sci. 2001, 26, 591-596.
- 65. Ibba, M.; Soll, D. Annu. Rev. Biochem. 2000, 69, 617-650.
- 66. Cusack, S. Curr. Opin. Struct. Biol. 1997, 7, 881-889.
- 67. Schimmel, P. Annu. Rev. Biochem. 1987, 56, 125-158.
- 68. Wang, L.; Brock, A.; Herberich, B.; Schultz, P. G. Science 2001, 292, 498-500.
- 69. Ibba, M.; Kast, P.; Hennecke, H. Biochemistry 1994, 33, 7107-7112.

- 24
- 70. Kwon, I.; Wang, P.; Tirrell, D. A. J. Am. Chem. Soc. 2006, 128, 11778-11783.
- 71. Ohno, S.; Yokogawa, T.; Nishikawa, K. J. Biochem. (Tokyo). 2001, 130, 417-423.
- Kiga, D.; Sakamoto, K.; Kodama, K.; Kigawa, T.; Matsuda, T.; Yabuki, T.; Shirouzu, M.; Harada, Y.; Nakayama, H.; Takio, K.; Hasegawa, Y.; Endo, Y.; Hirao, I.; Yokoyama, S. *Proc. Natl. Acad. Sci. U. S. A.* 2002, *99*, 9715-9720.
- Sakamoto, K.; Hayashi, A.; Sakamoto, A.; Kiga, D.; Nakayama, H.; Soma, A.; Kobayashi, T.; Kitabatake, M.; Takio, K.; Saito, K.; Shirouzu, M.; Hirao, I.; Yokoyama, S. *Nucleic Acids Res.* 2002, *30*, 4692-4699.
- Bose, M.; Groff, D.; Xie, J. M.; Brustad, E.; Schultz, P. G. J. Am. Chem. Soc. 2006, 128, 388-389.
- Bain, J. D.; Glabe, C. G.; Dix, T. A.; Chamberlin, A. R.; Diala, E. S. J. Am. Chem. Soc. 1989, 111, 8013-8014.
- Noren, C. J.; Anthony-Cahill, S. J.; Griffith, M. C.; Schultz, P. G. Science 1989, 244, 182-188.
- 77. Wang, L.; Brock, A.; Schultz, P. G. J. Am. Chem. Soc. 2002, 124, 1836-1837.
- Kohrer, C.; Xie, L.; Kellerer, S.; Varshney, U.; RajBhandary, U. L. Proc. Natl. Acad. Sci. U. S. A. 2001, 98, 14310-14315.
- Kohrer, C.; Yoo, J. H.; Bennett, M.; Schaack, J.; RajBhandary, U. L. Chem. Biol.
 2003, 10, 1095-1102.
- Alfonta, L.; Zhang, Z. W.; Uryu, S.; Loo, J. A.; Schultz, P. G. J. Am. Chem. Soc. 2003, 125, 14662-14663.
- Chin, J. W.; Cropp, T. A.; Anderson, J. C.; Mukherji, M.; Zhang, Z. W.; Schultz, P. G. Science 2003, 301, 964-967.

- Cornish, V. W.; Benson, D. R.; Altenbach, C. A.; Hideg, K.; Hubbell, W. L.; Schultz,
 P. G. *Proc. Natl. Acad. Sci. U. S. A.* **1994**, *91*, 2910-2914.
- Mehl, R. A.; Anderson, J. C.; Santoro, S. W.; Wang, L.; Martin, A. B.; King, D. S.;
 Horn, D. M.; Schultz, P. G. *J. Am. Chem. Soc.* 2003, *125*, 935-939.
- Thorbjarnardottir, S.; Uemura, H.; Dingermann, T.; Rafnar, T.; Thorsteinsdottir, S.;
 Soll, D.; Eggertsson, G. J. Bacteriol. 1985, 161, 207-211.
- 85. Yoshimura, M.; Inokuchi, H.; Ozeki, H. J. Mol. Biol. 1984, 177, 627-644.
- Kiick, K. L.; van Hest, J. C. M.; Tirrell, D. A. Angew. Chem. Int. Ed. 2000, 39, 2148-2152.
- 87. Kiick, K. L.; Weberskirch, R.; Tirrell, D. A. FEBS Lett. 2001, 505, 465-465.
- Chen, P.; Qian, Q.; Zhang, S. P.; Isaksson, L. A.; Bjork, G. R. J. Mol. Biol. 2002, 317, 481-492.
- Watanabe, Y.; Tsurui, H.; Ueda, T.; Furushima, R.; Takamiya, S.; Kita, K.; Nishikawa, K.; Watanabe, K. *Biochim. Biophys. Acta - Gene Struct. Express.* 1997, 1350, 119-122.
- 90. Lim, V. I.; Curran, J. F. RNA 2001, 7, 942-957.
- Inagaki, Y.; Kojima, A.; Bessho, Y.; Hori, H.; Ohama, T.; Osawa, S. J. Mol. Biol. 1995, 251, 486-492.
- 92. Curran, J. F. Nucleic Acids Res. 1995, 23, 683-688.
- Boren, T.; Elias, P.; Samuelsson, T.; Claesson, C.; Barciszewska, M.; Gehrke, C. W.;
 Kuo, K. C.; Lustig, F. *J. Mol. Biol.* **1993**, *230*, 739-749.
- 94. Osawa, S.; Jukes, T. H.; Watanabe, K.; Muto, A. Microbiol. Rev. 1992, 56, 229-264.
- 95. Munz, P.; Leupold, U.; Agris, P.; Kohli, J. Nature 1981, 294, 187-188.

- 96. Lim, V. I.; Aglyamova, G. V. Biol. Chem. 1998, 379, 773-781.
- 97. Lim, V. I. J. Mol. Biol. 1997, 266, 877-890.
- 98. Lim, V. I. J. Mol. Biol. 1995, 252, 277-282.
- 99. Lim, V. I.; Venclovas, C. FEBS Lett. 1992, 313, 133-137.
- 100. Murakami, H.; Hohsaka, T.; Ashizuka, Y.; Hashimoto, K.; Sisido, M. *Biomacromolecules* **2000**, *1*, 118-125.
- 101. Murakami, H.; Hohsaka, T.; Ashizuka, Y.; Sisido, M. J. Am. Chem. Soc. 1998, 120, 7520-7529.
- 102. Muranaka, N.; Hohsaka, T.; Sisido, M. FEBS Lett. 2002, 510, 10-12.
- 103. Taki, M.; Hohsaka, T.; Murakami, H.; Taira, K.; Sisido, M. J. Am. Chem. Soc. 2002, 124, 14586-14590.
- 104. Hohsaka, T.; Ashizuka, Y.; Taira, H.; Murakami, H.; Sisido, M. *Biochemistry* 2001, 40, 11060-11064.
- 105. Anderson, R. D.; Zhou, J.; Hecht, S. M. J. Am. Chem. Soc. 2002, 124, 9674-9675.
- 106. Sisido, M.; Ninomiya, K.; Ohtsuki, T.; Hohsaka, T. Methods 2005, 36, 270-278.
- 107. Taki, M.; Hohsaka, T.; Murakami, H.; Taira, K.; Sisido, M. *FEBS Lett.* **2001**, *507*, 35-38.
- 108. Taki, M.; Matsushita, J.; Sisido, M. ChemBioChem 2006, 7, 425-428.
- 109. Hohsaka, T.; Kajihara, D.; Ashizuka, Y.; Murakami, H.; Sisido, M. J. Am. Chem. Soc.
 1999, 121, 34-40.
- 110. Hohsaka, T.; Ashizuka, Y.; Murakami, H.; Sisido, M. *Nucleic Acids Res.* **2001**, *29*, 3646-3651.
- 111. Magliery, T. J.; Anderson, J. C.; Schultz, P. G. J. Mol. Biol. 2001, 307, 755-769.

- 112. Anderson, J. C.; Magliery, T. J.; Schultz, P. G. Chem. Biol. 2002, 9, 237-244.
- 113. Anderson, J. C.; Wu, N.; Santoro, S. W.; Lakshman, V.; King, D. S.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 2004, 101, 7566-7571.
- 114. Dudock, B. S.; Diperi, C.; Michael, M. S. J. Biol. Chem. 1970, 245, 2465-2468.



Figure 1: Nonnatural amino acids that have been genetically incorporated into proteins in bacteria, yeast or mammalian cells. This figure is reprinted, with permission, from the *Annual Review of Biophysics and Biomolecular Structure*, Volume **35**, Wang, L.; Xie, J.; Schultz, P. G., Expanding the genetic code, 225,²⁷ Copyright (2006) by Annual Reviews www.annualreviews.org.



Figure 2: Structures of yeast tRNA^{Phe} (left) and *E. coli* tRNA^{Phe} (right). This figure is reprinted from Dudock, B. S.; Diperi, C.; Michael, M. S. *J. Biol. Chem.* **1970**, *245*, 2465-2468,¹¹⁴ Copyright 1970 with permission from the American Society for Biochemistry and Molecular Biology.





Figure 3: (a) Anticodon-loop tRNA library (left) and all-loop tRNA library (right) derived from *M. jannaschii* tRNA^{Tyr}_{CUA} (N means all four nucleotides (A/U/G/C)). (b) A general selection scheme for suppressor tRNAs that are orthogonal to *the E. coli* endogenous systems and charged efficiently by a cognate synthetase. These figures are reprinted from from *Chem. Biol.*, **8**, Wang, L.; Schultz, P.G., A general strategy for orthogonal tRNA, 883,²⁵ Copyright (2001) with permission from Elsevier.



Figure 4: (a) Crystal structure of the *T. thermophilus* PheRS (*t*PheRS, pdb 1B70) in ribbon model. Only ($\alpha\beta$) monomeric portion of structure is shown. (b) Active sites of *t*PheRS. Only residues within 6 Å of substrate Phe are shown. Substrate is shown as space-filling model, while residues surrounding the substrate are shown in stick model. (c) Sequence alignment of PheRS from 21 different organisms. Only residues flanking equivalent residues of V261 and A314 in *T. thermophilus* are represented. The shaded residues are the conserved residues equivalent to V261 and A314 from *T. thermophilus*. Sequences adjacent to V261 and A314 are also highly conserved. The sequences are obtained from aminoacyl-tRNA synthetase database (http://rose.man.poznan.pl/aars/). These figures were adapted from Wang's thesis (2003).⁴¹



Starting E. coli library



Figure 5: (**A**) Phe substrate (red) and four residues (blue) within 7 Å of the *para*-position of the phenyl ring of the substrate inside the binding pocket of a homology model of yPheRS. (**B**) A screening scheme for yPheRS library. GFP6 in yPheRS expression library *E. coli* cells outfitted with ytRNA^{Phe}_{AAA} and yPheRS library was expressed in the presence of 2Nal (**a**). Weakly fluorescent cells that contain active yPheRS variants were enriched by FACS (**b**). GFP6 in the collected cells was expressed in the absence of 2Nal (**c**). Highly fluorescent cells that contain selective yPheRS variants were enriched by FACS (**d**). After two rounds of screening, ten colonies were isolated from the enriched cells and characterized (**e**).



Figure 6: A schematic diagram describing the concept of breaking the degeneracy of the genetic code.

Chapter 2

Site-Specific Incorporation of Phenylalanine Analogs into Proteins in Vivo by an Engineered Yeast Phenylalanyl-tRNA Synthetase

The work in this chapter was performed in collaboration with Pin Wang, Soojin Son, and Yi Tang.

Abstract

Aminoacyl-tRNA synthetases (aaRSs) catalyze the aminoacylation reaction to establish the rules of the genetic code. Precise manipulation of synthetase activity can alter aminoacylation specificity to attach non-canonical amino acids to the cognate transfer RNAs (tRNA). The codon-anticodon interaction between messenger RNA (mRNA) and tRNA determines which amino acid is delivered into a growing polypeptide chain. Thus introduction of nonnatural amino acids into proteins in vivo relies heavily on manipulation of amino acid specificity of the aaRS. In this report, we describe the generation and characterization of a mutant phenylalanyl-tRNA synthetase (PheRS) from Saccharomyces *cerevisiae* with a point mutation (T415G) in the α -subunit of the enzyme. The rationale for the mutation is to allow binding of bulkier amino acid substrates. The promiscuous substrate specificity of this mutant was extensively explored by ATP-PP_i exchange assays in vitro. A broad activation profile toward many nonnatural amino acids was observed. A phenylalanine auxotrophic E. coli strain transformed with this mutant synthetase and cognate suppressor tRNA enable the assignment of an amber nonsense codon to the amino acid tryptophan (Trp) or to the nonnatural amino acid 3-(2-naphthyl)alanine. Expression strains engineered as phenylalanine, tryptophan double auxotrophs or phenylalanine, tryptophan and lysine triple auxotrophs outfitted with this pair of mutant synthetase and tRNA makes possible the efficient incorporation of *p*-bromophenyl-, *p*-iodophenyl-, and *p*azidophenylalanine into recombinant proteins. Therefore, this synthetase and its cognate tRNA could serve as "21st" pair for site-specific incorporation of novel amino acid into proteins in vivo.

Introduction

Aminoacyl-tRNA synthetases (aaRSs) insure the high fidelity of transforming genetic code sequences into biologically functional proteins through a two-step aminoacylation reaction.¹⁻⁶ In the first step, the cognate amino acid is activated by aaRS in presence of ATP to form the amino acid adenylate; subsequently aaRS catalyzes the esterification reaction to join the amino acid to the 2'- or 3'-OH of the terminal ribonucleotide of its cognate tRNA. Through the aminoacylation reaction, the rules of the genetic code are settled; the strict correlation between triple nucleotide code and amino acid is established with the assistance of the aaRS.^{2,7,8} Manipulation of such a reaction could alter the genetic code to allow incorporation of novel amino acid into proteins in vivo.^{9,10}

With depletion of the intracellular pool of one natural amino acid and use of the corresponding bacterial auxotrophic cells, the rates of aminoacylation can be perturbed to enable charging of the nonnatural amino acid supplemented in the culture medium as a surrogate into cognate tRNA.⁹ Through subsequent translation by ribosomal machinery, the genetic code is re-assigned and the nonnatural amino acid is incorporated into proteins. We have successfully implemented such a method to incorporate alkenes,¹¹ alkynes,¹² alkyl azide¹³ and fluorinated side chains¹⁴ into proteins. After enhancing the cellular aminoacylation reactivity by over-expression of wild-type aaRS in the host, we found that some sluggish amino acid analogs could also be introduced into proteins.¹⁴⁻¹⁶ The method described above requires the cellular wild-type aaRSs to recognize these novel amino acids. Engineering new synthetase activity is almost inevitable when one wants to incorporate amino acids that are not recognized by the wild-type enzymes. To date, we and others have

shown that the re-design of the synthetic site of synthetase¹⁷⁻¹⁹ and the attenuation of the editing function of synthetase^{20,21} could be two complementary strategies to further expand our ability to introduce nonnatural amino acid into proteins.

We have shown that this multi-site incorporation method can utilize altered sets of 20 amino acids to design and engineer proteins and protein-like macromolecules.⁹ By manipulation of both synthetase and tRNA in the host, one can also accomplish sitespecific introduction of a single copy of a novel amino acid into proteins in vivo.²²⁻³¹ This method is derived from an in vitro approach to nonnatural amino acid incorporation through nonsense (stop codon) suppression, in which a stop codon (amber codon) was suppressed by a suppressor tRNA that had been chemically misacylated with the amino acid analog of interest.³²⁻³⁶ Such a chemical method to alter the aminoacylation in vitro suffers from the technical difficulty and the intrinsic low yield of protein production, which limits its application. In 1998, Furter modified the cellular aminoacylation reaction by imparting a "twenty-first pair" comprising a yeast suppressor tRNA and a yeast phenylalanyl-tRNA synthetase (vPheRS).²⁶ This approach allowed site-specific incorporation of L-p-fluorophenylalanine in vivo in response to an amber codon. More recently Schultz and colleagues have devised powerful selection methods to find useful mutant forms of tyrosyl-tRNA synthetase from archaebacterium Methanococcus jannaschii.^{10,37} By introduction of such a mutant and a cognate suppressor, many nonnatural amino acids have been incorporated into proteins in vivo.^{10,22-25,28-31,38-46}

In this report, we present a rationally designed variant of *y*PheRS with new amino acid specificity, allowing incorporation of a variety of phenylalanine analogs (2-4, 7, and 8) into proteins site-specifically. The variant of yPheRS contains a single T415G mutation at

the active site of the synthetase, which was shown to be very effective to open up the binding pocket. We performed a study on the amino acid activation kinetics of this mutant and found that this mutant can activate several interesting and useful aromatic amino acids (1-8). We also observed that this variant activates tryptophan (Trp, 6) 82-fold faster than cognate substrate phenylalanine (Phe, 1). A phenylalanine auxotrophic bacterial strain transformed with such a mutant and cognate suppressor tRNA can assign analog 7 to the amber codon with high fidelity (>90%). Such a transformed strain also allows the incorporation of 2-4 into proteins at the amber site with misincorporation of Trp. Further refinements of such a system by constructing a Phe/Trp/Lys triple auxotrophic strain enable us to incorporate analog 2-4 into the desired amber site with fidelity of 86%, 82%, and 90%, respectively.

Materials and Methods

Materials. Amino acids **1**, **5** and **6** were obtained from Sigma (St. Louis, MO). Amino acids **2-4**, and **7** were purchased from Chem-Impex (Wood Dale, IL). Amino acid **8** was purchased from Biosynth International (Naperville, IL). [³²P]-labeled sodium pyrophosphate was purchased from NEN Life Science (Boston, MA).

Plasmid Construction for Synthetase Expression. The yPheRS gene was amplified from template plasmid pUC-ASab2^{26,47} encoding the α - and β -subunits of PheRS gene with a 14-base pair (bp) intergenic sequence containing a translational reinitiation site upstream of the ATG start codon of the β -subunit gene. The following primers were used for PCR: 5'-CGA TTT TCA CAC AGG ATC CAG ACC ATG AAT CTA G-3' (primer 1 with restriction site BamHI) and 5'-GAC GGC CAG TGA ATT CGA GCT CGG TAC-3' (primer 2 with restriction site KpnI). The resulting DNA was introduced into the BamHI and KpnI sites of pQE32 to give pQE32-yFRS. The mutant yPheRS was generated by using the 4-primer mutagenesis method. Two complementary oligonucleotides, designated as primer 3 (5'-CTA CCT ACA ATC CTT ACG GCG AGC CAT CAA TGG AAA TC-3') for the forward primer and primer 4 (5'-GAT TTC CAT TGA TGG CTC GCC GTA AGG ATT GTA GGT AG-3') for the reverse primer, were synthesized to carry the specific mutation at position 415 of the α -subunit of yPheRS. In the initial two reactions, primer 1/primer 4 and primer 2/primer 3 were added into individual tubes and two DNA fragments were generated from these two PCRs. With the mixture of two reaction products and additional outside primers, a 3400 bp fragment of DNA was obtained and purified. This DNA was subjected to digestion by BamHI and KpnI and inserted into pQE32 to yield pQE32-T415G. The cloned enzymes contain the N-terminal sequence MRGSHHHHHHGIQTMNLE to facilitate protein purification. The entire yPheRS gene was sequenced for each of these constructs. Proofreading polymerase Pfx (Invitrogen) was used for all PCR amplication described in this work.

Synthetase Expression and Purification. The plasmids pQE32-yFRS and pQE32-T415G were individually transformed into *E. coli* strain BLR (Novagen) to form expression strains BLR(pQE32-yFRS) and BLR(pQE32-T415G). Synthetase expression was conducted in LB media. At an OD of 0.6, expression of the wild-type and mutant forms of yPheRS was induced with 1 mM isopropyl-1-thio- β -D-galactopyranoside (IPTG) for 4 hours. Cells were harvested and proteins were purified over a nickel-nitrilotriacetic

acid affinity column under native conditions according to the manufacturer's protocol (Qiagen). Imidazole in the elution buffer was removed by desalting column (Bio-rad) and proteins were eluted into a buffer containing 50 mM Tris-HCl (pH=7.5), 1 mM DTT. Aliquots of proteins were stored at -80°C with 50% of glycerol.

Amino Acid Activation Assay. The amino acid-dependent ATP-PP_i exchange reaction was used to evaluate the activation of amino acid analogs by *y*PheRS. This assay was performed in 200 μ L of reaction buffer containing 50 mM HEPES (pH=7.6), 20 mM MgCl₂, 1 mM DTT, 2 mM ATP and 2 mM [³²P]-PP_i with specific activity of 0.2-0.5 TBq/mol. Depending on the rate of activation of the analog by the synthetase, the amino acid concentration varied from 10 μ M to 5 mM and the enzyme concentration varied from 10 nM to 100 nM. Aliquots (20 μ L) were removed from the reaction solution at various time points and quenched into 500 μ L of buffer solution containing 200 mM NaPP_i, 7% w/v HClO₄ and 3% w/v activated charcoal. The charcoal was spun down and washed twice with 500 μ L of 10 mM NaPP_i and 0.5% HClO₄ solution. The [³²P]-labeled ATP absorbed into the charcoal was quantified via liquid scintillation methods. The specificity constants were calculated by nonlinear regression fit of the data to a Michaelis-Menten model.

Plasmid and Strain Construction for in Vivo Incorporation Assays. pQE16 (Qiagen), chosen as the expression plasmid, encodes the marker protein murine dihydrofolate reducase (mDHFR) with a C-terminal hexa-histidine tag under control of a bacteriophage T5 promoter and t₀ terminator. A Quick-change mutagenesis kit was used to place an

amber codon (UAG) at position 38 of mDHFR (mDHFR (38Am)) (with two complementary oligonucletides (5'-CCG CTC AGG AAC GAG TAG AAGTAC TTC CAA AGA ATG-3'; 5'-CAT TCT TTG GAA GTA CTT CTA CTC GTT CCT GAG CGG-3') to afford pQE16am. The mutant vPheRS (T415G) gene was amplified from pQE32-T415G and a constitutive tac promoter with an abolished lac repressor binding site was added upstream of the start codon of this gene.²⁶ The entire expression cassette was inserted into the PvuII site of pQE16 to yield pQE16am-T415G. The mutant yeast suppressor tRNA (*mu*tRNA^{Phe}(CUA)) was constitutively expressed under control of an *lpp* promoter. The expression cassette for mutRNA^{Phe}(CUA) was inserted into pREP4 to form pREP4-tRNA as described by Furter.²⁶ A Phe auxotrophic strain AFK10, Hfr(Cavalli) pheS13rel-1 tonA22 thi T2^R pheA18), constructed in this laboratory,²⁶ was used as the expression strain. A Phe/Trp double auxotrophic strain AFW (K10, Hfr(Cavalli) pheS13rel-1 tonA22 thi T2^R pheA18, trpB114) and a Phe/Trp/Lys triple auxotrophic strain AFWK (K10, Hfr(Cavalli) pheS13rel-1 tonA22 thi T2^R pheA18, trpB114, lysA) were prepared by P1 phage-mediated transduction with *trpB::Tn10* and *lvsA::Tn10* transposons.

Analog Incorporation Assay in Vivo. The auxotrophic strains AF, AFW, and AFWK were transformed with the plasmids pQE16am-T415G and pREP4-tRNA to afford expression strains AF [pQE16am-T415G/pREP4-tRNA] and AWF [pQE16am-T415G/pREP4-tRNA] respectively. Expression of mDHFR (38Am) was investigated in 20 mL cultures. The *E. coli* expression strains were grown in M9 minimal medium supplemented with glucose, thiamin, MgSO₄, CaCl₂, 20 amino acids (at 25 mg/L), and antibiotics (kanamycin and ampicillin). When cells reached an OD₆₀₀ of 1.0, they were

sedimented by centrifugation, washed twice with cold 0.9% NaCl, and shifted to supplemented M9 medium containing 17 amino acids (at 25 mg/L), 3 mM analog of interest, and the indicated concentrations of Phe, Trp and Lys. Protein expression was induced by addition of isopropyl-β-D-thiogalactopyranoside (1 mM). After 4 hours, cells were pelleted and the protein was purified by passage through a Ni-NTA spin column according to the supplier's instructions (Qiagen).

Compositional Analysis of Mutant mDHFR. Mutant mDHFR (38Am) was purified under denaturing conditions and isolated in elution buffer (8 M urea, 100 mM NaH₂PO₄, 10 mM Tris, pH = 4.5). For trypsin digestion, 10 μ L of the solution was diluted into 90 μ L of 75 mM (NH₄)₂CO₃ and the pH was adjusted to 8. 2 μ L of modified trypsin (Promega, 0.2 μ g/ μ L) was added. The sample was incubated at room temperature overnight. For digestion by endoproteinase Lys-C, 10 μ L of the solution was diluted into 90 μ L of 25 mM Tris-HCl, pH = 8 and 1 mM EDTA and pH was adjusted to optimal working pH 8 for Lys-C. 2 μ L of Lys-C (Calbiochem, 0.2 μ g/ μ L) was added and the reaction was incubated at 37 °C for 10 hours. The digestion reaction was stopped by addition of 2 μ L trifluoroacetic acid (TFA). The solution was subjected to ZipTip_{C18} (Millipore) purification and the digested peptides were eluted with 3 μ L of 50% CH₃CN, 0.1% TFA, of which 1 μ L was used for matrix-assisted laser desorption ionization mass spectrometry (MALDI-MS) analysis with μ -cyano-4-hydroxycinnamic acid and 2,5-dihydroxybenzoic acid as the matrix. The analysis was performed on a PerSeptive Biosystems (Framingham, Massachusetts) Voyager DE PRO MALDI-TOF mass spectrometer operating in linear and positive ion modes.

LC-MS/MS analysis of protease-digested peptides was conducted on Finnigan LCQ ion trap mass spectrometer with HPLC pump and ESI probe. 3 μ L of digested peptide solution eluted from ZipTip_{C18} was diluted with 20 μ L of distilled water and injected into the HPLC pump. Peptides were separated by Magic C18 column (Michrom, 5 μ L, 200Å, 0.3x150mm) and eluted at a flow rate of 30 μ L/min using a linear gradient of 0-60% of acetonitrile in 45 min. The column eluent was transferred to the electrospray source and tandem mass sequencing was carried out by fragmentation of the precursor ion with m/z corresponding to the protease-digested fragment (residues 26-39; NGDLPWPPLRNE**Z**K) including an amber site at position 38 of mutant mDHFR (38Am).

Results and Discussion

Rationales for Engineering New Synthetase Specificity. Engineering the new synthetase specificity toward amino acid analogs is one of the critical steps for the successful implementation of the general strategy to incorporate nonnatural amino acids into proteins in vivo.²⁹ We have been especially interested in exploring the possibilities of incorporating substituted phenylalanine (Figure 2) such as *p*-bromophenylalanine (2), *p*-iodophenylalanine (3), *p*-azidophenylalanine (4), and 3-(2-naphthyl)alanine (7). These non-proteinogenic amino acids either carry chemical functionality not present in proteins or are characterized by enhanced hydrophobic interactions. The crystal structure of *Thermus thermophilus* PheRS (tPheRS) complexed with phenylalanine is available⁴⁸ and the

sequence identity (*vs Saccharomyces cerevisiae*) in the active site region is about 40% (Figure 1). We used this crystal structure as a starting point for the design of a mutant yeast PheRS that charges the nonnatural amino acids of interest. Safro and colleagues identified two residues, V261 and A314, which constitute a back-wall structure to hold substrate phenylalanine and play an important role to hinder binding of amino acids larger than Phe into the active site of *T. thermophilus* PheRS.⁴⁹ In fact, an *E. coli* PheRS variant prepared by mutation of the corresponding residues (T251G and A314G, Figure 1), has been shown to exhibit relaxed substrate specificity.¹⁷ In *S. cerevisiae* PheRS, the corresponding residues are T415 and G460. We reasoned that mutation from T415 to G might enlarge the active site to accommodate larger phenylalanine analogs.

Synthetase Expression and Purification. The wild-type yPheRS gene was obtained by PCR amplification of vector pUC-ASab2 encoding the α - and β -subunits of yeast PheRS (our assignment of α - and β -subunits are opposed to what was assigned by Sanni; we followed the tradition that the catalytic subunit of PheRS is designated the α - subunit.^{47,48,50}). A 14-bp intergenic sequence obtained from upstream region of the *E. coli* pheST gene was inserted in the middle of the α - and β -subunits of yeast PheRS. Four-primer mutagenesis was employed to create the mutant T415G. The synthetases were over-expressed by *E. coli* BLR strains under control of the bacteriophage T5 promoter. An N-terminal hexa-histidine tag fused to the α -subunit of yPheRS facilitated purification of the enzymes under native conditions. Both *E. coli* and yeast synthetases are $\alpha_2\beta_2$ hetero-tetramers and their molecular weights for each subunit are rather different (α (ePheRS)=37

kDa; α (yPheRS)=57 kDa; β (ePheRS)=87 kDa; β (yPheRS)=67.5). SDS-PAGE analysis of purified yPheRS indicated no observable contamination by *E. coli* PheRS. The intensities of the bands for the α - and β -subunits are not stoichiometric (the α -subunit is stronger than the β -subunit); Sanni and colleagues reported a similar observation and suggested that this result stems from proteolysis of the β -subunit and from one β -fragment co-migrating with the α -subunit.⁴⁷ So a fraction of the purified yPheRS used in our experiments must be inactive. Our aim is to investigate the altered substrate specificity from the mutation (T415G for α -subunit) and compare the ability of the mutant enzyme to activate different amino acid analogs (**2-8**) in vitro; as long as we paralleled the measurement and used the same batch of the enzyme, it is still possible to deduce the specificity constants for the different analogs and make comparisons.

Amino Acid Specificity of Mutant Yeast PheRS. The activities of the yPheRS variant with Phe analogs (2-8) were examined via amino acid-dependent ATP-PP_i exchange assay. Kinetic parameters for amino acid 1-8 were shown in Table 1. Our measured value of K_m for 1 by the wild-type yeast PheRS matched previously reported values,⁵¹ although the obtained k_{cat} was lower than that reported,⁵¹ and this difference might come from the different buffer conditions and different methods to determine the enzyme concentration. The efficiency k_{cat}/K_m for 1 by the mutant T415G is reduced approximately 40-fold compared to 1 by the wild-type enzyme. The loss of catalytic activity is apparent in both the increasing value of K_m and decreasing value of k_{cat} , which suggests that the T415G mutation significantly impairs both the binding and catalytic events for 1. However, this mutant exhibited higher activity for analogs 2-4 and 6-8, with values of k_{cat}/K_m ranging

from 8- to 58-fold higher than that for Phe (1). It appears that the mutation effectively opens up the active site of the synthetase so that T415G favors binding and catalysis of bulkier substrates. Canonical amino acid Trp (6) showed especially high activity, as manifested by 82-fold higher specificity constant than that of 1.

Site-Specific Incorporation of Phe Analogs into mDHFR (38Am) in Vivo Using a Phenylalanine Auxotrophic Strain. To test the utility of the engineered yPheRS to incorporate aromatic amino acid analogs into proteins in vivo at programmed sites, a constitutive expression system for the mutant synthetase was constructed. An amber stop codon was placed at residue 38 of the marker protein mDHFR (38Am), which was equipped with a C-terminal hexa-histidine tag followed by an ochre stop codon (Figure 3). The levels of full length protein production were examined by SDS-PAGE analysis. Without over-expression of suppressor mutRNA^{Phe}(CUA), we did not obtain detectable levels of full length mDHFR (38Am).⁵² As a positive control, we co-expressed wild-type yPheRS with mutRNA^{Phe}(CUA), and efficient suppression by phenylalanine was observed.⁵² When mutant vPheRS was co-expressed with mutRNA^{Phe}(CUA) in the presence of analogs 2-8 (3 mM), full length mDHFR (38Am) was formed.⁵² Overall protein yields varied depending on the analogs used and were ca. 2-10 mg/L, estimated by SDS-PAGE. The same strain yield of mDHFR without amber codon in minimal medium was approximately 25 mg/L.

It should be noted that the observed pattern of protein expression by SDS-PAGE does not necessarily mean that the analogs of interest are inserted at the amber codon position. Proteolytic peptide analysis was performed by MALDI-MS and liquid chromatography tandem mass spectrometry for purified protein samples. Our protocol for digestion by endoproteinase Lys-C yielded two peptides in the mass range 1,600-1,800 Da; these fragments were assigned to residues 26-39 (Peptide A) and 40-53 (Peptide B) of mDHFR (38Am) (Figure 3). Peptide A includes an amber codon at the 38th position; Peptide B contains a single phenylalanine residue at position 41. In a control experiment, in which mDHFR (38Am) was expressed in medium supplemented with 1 (3 mM) (Figure 4a), the mass of Peptide A indicates that 1 was incorporated into the amber site. Our in vitro assay shows that T415G mutant synthetase activates 6 82-fold faster than 1; while our in vivo translational experiment reveals that 1 is favored over 6 to enter amber site, despite the fact that Phe auxotrophic strain can still synthesize 6. In the medium supplemented with 7 (3 mM) and 1 (0.03 mM), we unambiguously confirmed incorporation of 7 into the amber site as shown in Figure 4b; Peptide A was shifted up in mass by 50 Da, consistent with the increased mass of 7 relative to 1; and Peptide B was unchanged in mass, indicating that the Phe codon site is decoded as 1. Liquid chromatography tandem mass spectrometry was employed to confirm the sequence of Peptide A. The precursor ion at m/z 867.7 Da, which corresponds to the doubly charged ion of Peptide A, was separated and fragmented with an ion trap mass spectrometer. As shown in Figure 5, the sequence obtained from the spectrum clearly reveals site-specific incorporation of 7 into mDHFR at position 38; we could find no indication of any other amino acid at the amber site after carefully checking all possible peptides with masses that could be associated with Peptide A, suggesting that the fidelity of incorporation of 7 is better than 95%. The kinetics of activation of 7 and 1 by mutant yPheRS showed that the efficiency (k_{cat}/K_m) for 7 is 1,500±100 (M⁻¹s⁻¹) (Table 1), 8-fold larger than that for 1. Therefore, when the ratio of 7 to 1 in the culture medium is

high (100 to 1), tRNA^{Phe}(CUA) should be charged predominantly with 7, consistent with our observation of high percentage incorporation of 7 into proteins at amber sites.

Similar analyses show incorporation of **2-4** into mDHFR (38Am). As a representative, incorporation of **2** into mDHFR (38Am) was shown (Figure 4c). We observed competitive misincorporation of Trp (**6**) and lysine (Lys, **8**) at the amber site. Incorporation of Trp is not surprising in view of the in vitro data. The rates of activation of **2-4** by T415G are 10-33% of that of **6** (Table 1). Therefore, **6** is able to compete significantly with **2-4** for the attachement into mutRNA^{Phe}(CUA) and insertion into the amber site. Lys misincorporation at the amber codon is likely due to mischarging of Lys to mutRNA^{Phe}(CUA) by *E. coli* lysyl-tRNA synthetase (eLysRS), because mutRNA^{Phe}(CUA) has U36 and A73, which are the main identity elements for eLysRS .^{53,54} No observation of misincorporation of **6** or **8** with **7** (Figure 4b) could be due to the better recognition of **7** by mutant synthetase than that of **2-4** (Table 1) and the better charging of **7** into mutRNA^{Phe}(CUA) by mutant synthetase than that of eLysRS.

Site-Specific Incorporation of Phe Analogs into mDHFR in Vivo Using a Phe/Trp Double Auxotrophic Strain. To test our hypothesis of intracellular tryptophan inhibiting the incorporation of analogs (2-4), we decided to construct a Phe/Trp double auxotrophic strain so that the cellular concentration of Trp could be controlled. This strain, designated AFW, was transformed with mutant yPheRS, mDHFR (38Am) and mutRNA^{Phe}(CUA) genes to form a new expression system. In order to confirm our ability to control the cellular pool of Trp in this new strain, we performed a comparison experiment where we set one high concentration of Trp (3 mM) and one low concentration of Trp (0.03 mM) in

M9 minimal medium to examine the effect on site-specific incorporation of 2 (3 mM) in response to an amber stop codon. Proteolytic cleavage of mDHFR (38Am) by trypsin generated two peptides in the mass range 1550-1800 Da, which consistently appeared in the mass spectra and could be assigned to residues 26-39 (Peptide A) and 85-98 (Peptide C) of mDHFR (Figure 4, Figure 5). Peptide C contains a single phenylalanine residue at position 95. MALDI-MS analysis revealed that Trp concentration has a significant effect on incorporation of **2**. As shown in Figure 6a, **2** is incorporated into mDHFR along with Trp; the ratio of MALDI signal intensities suggests that 2 and 6 are incorporated at comparable rates at the amber site, although signal intensity and the peptide concentration are not strictly correlated. In contrast, as we lowered the Trp concentration to 0.03 mM, the signal for Trp incorporation greatly decreased; the amber position was mostly assigned by analog 2 (Figure 6b). Since the biosynthesis of Trp in the auxotrophic strain is impeded, cellular concentration of Trp is dependent on the amino acid transport from medium to cytoplasm. Our previous work showed that the cellular concentration of 2 is 14-17 times higher than that of Phe in the Phe auxotrophic strain cultured in media supplemented with 2 mM of 2 and 0.12 mM of Phe.¹⁹ Compared with Trp, we believe that the 10-fold lower amino acid activation of 2 by mutant yPheRS (Table 1) can be compensated by high concentration of 2 inside the cells; mutRNA^{Phe}(CUA) is charged mostly with 2. However, we observed that Lys was still misincorporated at amber codon (Figure 6b).

Site-Specific Incorporation of Phe Analogs into mDHFR in Vivo Using a Phe/Trp/Lys Triple Auxotrophic Strain. Based on our hypothesis that Lys was mischarged into mutRNA^{Phe}(CUA) by eLysRS, we decided to control the intracellular concentration of Lys

by constructing a Phe/Trp/Lys triple auxotrophic strain (AFWK). When a high concentration of Lys (0.2 mM) was supplemented into the medium, misincorporation of Lys was detected (Figure 6b). However, when low concentration of Lys (0.01 mM) was supplemented into the medium, the signal for the peptide containing Lys at amber codon is no longer detected (Figure 6c). In conclusion, under the condition that high concentrations of **2** and low concentrations of Phe, Trp and Lys were supplemented into the medium, incorporation of **2** is predominant at the programmed amber stop codon in mDHFR (38Am) with 86% fidelity. Further experiments on analog **3** (Figure 6d) and analog **4** manifest a similar pattern of the site-directed analog incorporation with 82% and 90% fidelity, respectively.⁵²

Conclusions

The amino acid binding pocket of yeast PheRS can be engineered very simply to accommodate a variety of aromatic side chain functionalities. Cells outfitted with mutant yeast synthetase (T415G) and suppressor tRNA can site-specifically incorporate several amino acid analogs (2-4, and 7) into proteins in response to the UAG codon. Coupled with recent reports from several laboratories,^{22-25,27,29-31,38-42,44,45,55-57} these results begin to demonstrate that the import of heterologous synthetase pairs should constitute a general strategy for site-specific incorporation of amino acids into proteins in vivo. Chemical and physical properties of these new side chains can enlarge our ability to manipulate proteins provide us tools to study and design new functions and additional of biomacromolecules.^{29,33,41,58-60} For instance, bromine (2) and iodine (3) once introduced into proteins can facilitate phasing of crystallographic data and therefore be useful in X-ray

diffraction studies of protein structure. Photoactivity of aryl azide (4) can be used for intramolecular crosslinking and protein immobilization.⁶¹ Hydrophobic nature of naphthalene (7) can be applied to design specific ligand-receptor recognition.⁶² Chemical utilities of these side chains equip us with new means to site-specifically modify proteins, most of which are orthogonal to natural amino acid side chains in the proteins. Proteins with aryl-halide side chains can be subjected to palladium-mediated coupling reactions to attach molecules containing the ethynyl group.⁶³ Azide chains in the proteins can be applied to staudinger ligation to tether molecules associated with the triarylphosphine group.¹³ Recently, there is a growing interesting in possible bioconjugation of proteins using copper(I)-catalyzed azide-alkyne cycloaddition reactions.⁶⁴⁻⁶⁶ Our group is currently doing these investigations using recombinant proteins biosynthesized in vivo in the engineered bacterial hosts described in this report.

Acknowledgments

We would like to thank Dr. Mona Shahgholi for her assistance with tryptic digest mass spectrometry. We are grateful to Dr. Aaron J. Link for helpful discussions on P1 phage-mediated transduction.

References

- 1. Cusack, S. Curr. Opin. Struct. Biol. 1997, 7, 881-889.
- 2. de Pouplana, L. R.; Schimmel, P. Cell. Mol. Life Sci. 2000, 57, 865-870.
- 3. Francklyn, C.; Perona, J. J.; Puetz, J.; Hou, Y. M. RNA 2002, 8, 1363-1372.
- 4. Ibba, M.; Soll, D. Annu. Rev. Biochem. 2000, 69, 617-650.

- 5. Schimmel, P. Annu. Rev. Biochem. 1987, 56, 125-158.
- Soll, D.; RajBhandary, U. L., Structure, Biosynthesis, and Function. In *tRNA*, ASM Press: Washington, D. C., 1995.
- Nangle, L. A.; Lagardt, V. D.; Dodring, V.; Schimmel, P. J. Biol. Chem. 2002, 277, 45729-45733.
- 8. Tamura, K.; Schimmel, P. Nat. Biotechnol. 2002, 20, 669-670.
- 9. van Hest, J. C. M.; Tirrell, D. A. Chem. Commun. 2001, 1897-1904.
- 10. Wang, L.; Schultz, P. G. Chem. Commun. 2002, 1-11.
- 11. van Hest, J. C. M.; Tirrell, D. A. FEBS Lett. 1998, 428, 68-70.
- 12. van Hest, J. C. M.; Kiick, K. L.; Tirrell, D. A. J. Am. Chem. Soc. 2000, 122, 1282-1288.
- Kiick, K. L.; Saxon, E.; Tirrell, D. A.; Bertozzi, C. R. Proc. Natl. Acad. Sci. U. S. A.
 2002, 99, 19-24.
- 14. Tang, Y.; Tirrell, D. A. J. Am. Chem. Soc. 2001, 123, 11089-11090.
- Kiick, K. L.; van Hest, J. C. M.; Tirrell, D. A. Angew. Chem. Int. Ed. 2000, 39, 2148-2152.
- 16. Kiick, K. L.; Weberskirch, R.; Tirrell, D. A. FEBS Lett. 2001, 505, 465-465.
- Datta, D.; Wang, P.; Carrico, I. S.; Mayo, S. L.; Tirrell, D. A. J. Am. Chem. Soc. 2002, 124, 5652-5653.
- 18. Kirshenbaum, K.; Carrico, I. S.; Tirrell, D. A. ChemBioChem 2002, 3, 235-237.
- 19. Sharma, N.; Furter, R.; Kast, P.; Tirrell, D. A. FEBS Lett. 2000, 467, 37-40.
- Doring, V.; Mootz, H. D.; Nangle, L. A.; Hendrickson, T. L.; de Crecy-Lagard, V.;
 Schimmel, P.; Marliere, P. *Science* 2001, *292*, 501-504.
- 21. Tang, Y.; Tirrell, D. A. Biochemistry 2002, 41, 10635-10645.

- Chin, J. W.; Martin, A. B.; King, D. S.; Wang, L.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 2002, 99, 11020-11024.
- 23. Chin, J. W.; Santoro, S. W.; Martin, A. B.; King, D. S.; Wang, L.; Schultz, P. G. J. Am. *Chem. Soc.* **2002**, *124*, 9026-9027.
- 24. Deiters, A.; Geierstanger, B. H.; Schultz, P. G. ChemBioChem 2005, 6, 55-58.
- Deiters, A.; Groff, D.; Ryu, Y. H.; Xie, J. M.; Schultz, P. G. Angew. Chem. Int. Ed.
 2006, 45, 2728-2731.
- 26. Furter, R. Protein Sci. 1998, 7, 419-426.
- 27. Kwon, I.; Wang, P.; Tirrell, D. A. J. Am. Chem. Soc. 2006, 128, 11778-11783.
- Mehl, R. A.; Anderson, J. C.; Santoro, S. W.; Wang, L.; Martin, A. B.; King, D. S.;
 Horn, D. M.; Schultz, P. G. J. Am. Chem. Soc. 2003, 125, 935-939.
- 29. Wang, L.; Brock, A.; Herberich, B.; Schultz, P. G. Science 2001, 292, 498-500.
- 30. Wang, L.; Brock, A.; Schultz, P. G. J. Am. Chem. Soc. 2002, 124, 1836-1837.
- 31. Wang, L.; Xie, J. M.; Deniz, A. A.; Schultz, P. G. J. Org. Chem. 2003, 68, 174-176.
- Bain, J. D.; Glabe, C. G.; Dix, T. A.; Chamberlin, A. R.; Diala, E. S. J. Am. Chem. Soc.
 1989, 111, 8013-8014.
- 33. Dougherty, D. A. Curr. Opin. Chem. Biol. 2000, 4, 645-652.
- 34. Gilmore, M. A.; Steward, L. E.; Chamberlin, A. R., Incorporation of noncoded amino acids by in vitro protein biosynthesis. In *Implementation and Redesign of Catalytic Function in Biopolymers*, 1999; Vol. 202, pp 77-99.
- Heckler, T. G.; Chang, L. H.; Zama, Y.; Naka, T.; Chorghade, M. S.; Hecht, S. M. Biochemistry 1984, 23, 1468-1473.

- Noren, C. J.; Anthony-Cahill, S. J.; Griffith, M. C.; Schultz, P. G. Science 1989, 244, 182-188.
- 37. Wang, L.; Schultz, P. G. Chem. Biol. 2001, 8, 883-890.
- 38. Anderson, J. C.; Schultz, P. G. Biochemistry 2003, 42, 9598-9608.
- 39. Chin, J. W.; Schultz, P. G. ChemBioChem 2002, 3, 1135-1137.
- Deiters, A.; Cropp, T. A.; Mukherji, M.; Chin, J. W.; Anderson, J. C.; Schultz, P. G. J. Am. Chem. Soc. 2003, 125, 11782-11783.
- Schultz, K. C.; Supekova, L.; Ryu, Y. H.; Xie, J. M.; Perera, R.; Schultz, P. G. J. Am. Chem. Soc. 2006, 128, 13984-13985.
- 42. Wang, J. Y.; Xie, J. M.; Schultz, P. G. J. Am. Chem. Soc. 2006, 128, 8738-8739.
- 43. Wang, L.; Xie, J.; Schultz, P. G. Annu. Rev. Biophys. Biomol. Struct. 2006, 35, 225-249.
- 44. Wang, L.; Zhang, Z. W.; Brock, A.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 2003, 100, 56-61.
- 45. Xie, J. M.; Schultz, P. G. Curr. Opin. Chem. Biol. 2005, 9, 548-554.
- Xie, J. M.; Wang, L.; Wu, N.; Brock, A.; Spraggon, G.; Schultz, P. G. *Nat. Biotechnol.* 2004, 22, 1297-1301.
- 47. Sanni, A.; Walter, P.; Ebel, J. P.; Fasiolo, F. Nucleic Acids Res. 1990, 18, 2087-2092.
- Reshetnikova, L.; Moor, N.; Lavrik, O.; Vassylyev, D. G. J. Mol. Biol. 1999, 287, 555-568.
- Fishman, R.; Ankilova, V.; Moor, N.; Safro, M. Acta Crystallogr. D. Biol. Crystallogr.
 2001, 57, 1534-44.

- Fayat, G.; Mayaux, J. F.; Sacerdot, C.; Fromant, M.; Springer, M.; Grunbergmanago, M.; Blanquet, S. J. Mol. Biol. 1983, 171, 239-261.
- 51. Gabius, H. J.; Vonderhaar, F.; Cramer, F. Biochemistry 1983, 22, 2331-2339.
- 52. Wang, P., Ph.D. thesis, California Institute of Technology, Pasadena, 2003.
- McClain, W. H.; Foss, K.; Jenkins, R. A.; Schneider, J. Proc. Natl. Acad. Sci. U. S. A. 1990, 87, 9260-9264.
- Tamura, K.; Himeno, H.; Asahara, H.; Hasegawa, T.; Shimizu, M. *Nucleic Acids Res.* 1992, 20, 2335-2339.
- 55. Chin, J. W.; Cropp, T. A.; Anderson, J. C.; Mukherji, M.; Zhang, Z. W.; Schultz, P. G. *Science* 2003, *301*, 964-967.
- Wu, N.; Deiters, A.; Cropp, T. A.; King, D.; Schultz, P. G. J. Am. Chem. Soc. 2004, 126, 14306-14307.
- 57. Zhang, Z. W.; Alfonta, L.; Tian, F.; Bursulaya, B.; Uryu, S.; King, D. S.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 2004, 101, 8882-8887.
- Cornish, V. W.; Benson, D. R.; Altenbach, C. A.; Hideg, K.; Hubbell, W. L.; Schultz,
 P. G. *Proc. Natl. Acad. Sci. U. S. A.* **1994**, *91*, 2910-2914.
- Liu, H. T.; Wang, L.; Brock, A.; Wong, C. H.; Schultz, P. G. J. Am. Chem. Soc. 2003, 125, 1702-1703.
- Summerer, D.; Chen, S.; Wu, N.; Deiters, A.; Chin, J. W.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 2006, 103, 9785-9789.
- 61. Fleming, S. A. Tetrahedron 1995, 51, 12479-12520.
- Chan, D. C.; Chutkowski, C. T.; Kim, P. S. Proc. Natl. Acad. Sci. U. S. A. 1998, 95, 15613-15617.
- 63. Genet, J. P.; Savignac, M. J. Organomet. Chem. 1999, 576, 305-317.
- 64. Kolb, H. C.; Finn, M. G.; Sharpless, K. B. Angew. Chem. Int. Ed. 2001, 40, 2004-2021.
- 65. Link, A. J.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 11164-11165.
- 66. Link, A. J.; Vink, M. K. S.; Tirrell, D. A. J. Am. Chem. Soc. 2004, 126, 10598-10602.



Figure 1: (a) Sequence alignment of PheRS variants from *Thermus thermophilus*, *Escherichia coli*, *Saccharomyces cerevisiae*. Only portions of sequences flanking the V261 and A314 positions in *T. thermophilus* are shown. The sequences adjacent to V261 and A314 are highly conserved. The equavilent residues in *E. coli* and *S. cerevisiae* that correspond to residues V261 and A314 in *T. thermophilus* are shown in red. (b) Stereoview of active site of PheRS from *T. thermophilus*. Substrate phenylalanine is shown in yellow.



Figure 2: Amino acids involved in this study. (1) phenylalanine; (2) *p*-bromophenylalanine; (3) *p*-iodophenylalanine; (4) *p*-azidophenylalanine; (5) tyrosine; (6) trytophan; (7) 3-(2-naphthyl)alanine; and (8) L-lysine.

Table 1: Kinetic parameters for ATP-PPi exchange of amino acids (1-7) by the
 mutant yeast PheRS

Amino Acid	Enzyme	$K_{\rm m}$ (μ M)	$k_{\text{cat}} (\text{s}^{-1})$	$k_{\rm cat}/K_{\rm m}({\rm M}^{-1}{\rm s}^{-1})$	$k_{\text{cat}}/K_{\text{m}}$ (rel)*
1	T415G	264±42	0.05 ± 0.002	180±30	1
2	T415G	22±3	0.03 ± 0.001	$1,500\pm200$	8
3	T415G	12±2	0.05 ± 0.001	4,400±800	24
4	T415G	11±3	0.05 ± 0.002	4,600±1,200	25
5	T415G	757±149	0.04 ± 0.003	48±10	1/4
6	T415G	20±5	0.30 ± 0.006	15,000±4,000	82
7	T415G	27±2	0.04 ± 0.001	1,600±100	8
1	Wild-type	68±20	0.52±0.093	7,600±2,700	41

frelative to $k_{\text{cat}}/K_{\text{m}}$ for 1 by 1415G



Figure 3: Amino acid sequence of marker protein mDHFR (38Am). Four proteolytic peptide fragments (Peptide A, Peptide B, and Peptide C) used for MALDI-MS and LC-MS/MS analyses are underscored. The positions with amber codon and ochre codon are labeled.



Figure 4: MALDI-MS of proteolytic peptide fragments derived from mDHFR expressed in media supplemented with (a) **1** (3 mM); (b) **7** (3 mM)/**1** (0.03 mM); (c) **2** (3 mM)/**1** (0.03). mDHFR (38Am) was expressed in phenylalanine auxotrophic strain (AF) transformed with the mutant yeast PheRS and cognate suppressor tRNA. No tryptophan is added, except that 1 mM of Trp is supplemented in (c). Protein was digested by endoproteinase Lys-C. Peptide B, containing one Phe codon, remains the same for all the experimental conditions, indicating normal Phe codons are assigned as Phe. Peptide A contains one amber codon; the residue responding to this codon is designated as **Z**. In case of Lys misincorporation into Peptide A (c), C-terminal Lys was further cleaved to produce a shorter Peptide A (NGDLPWPPLRNEK).



Figure 5: The tandem mass spectrum of Peptide A (NGDLPWPPLRNE**Z**K) derived from mDHFR (38Am) expressed in media supplemented with 7 (3 mM)/1 (0.03 mM). mDHFR (38Am) was expressed in phenylalanine auxotrophic strain (AF) transformed with the mutant yeast PheRS and cognate suppressor tRNA. Partial sequence of PWPPLRNE and residue Z (corresponds to 7) of Peptide A can be assigned from the annotated y and b ion series respectively.





Figure 6: MALDI-MS of proteolytic peptide fragments of mDHFR (38Am) expressed in media supplemented with (a) **2** (3 mM)/**6** (3 mM)/**1** (0.03 mM)/**8** (0.2 mM); (b) **2** (3 mM)/**6** (0.01 mM)/**1** (0.03 mM)/**8** (0.2 mM); (c) **2** (3 mM)/**6** (0.01 mM)/**1** (0.03 mM)/**8** (0.01 mM); (d) **3** (3 mM)/**6** (0.01 mM)/**1** (0.03 mM)/**8** (0.01 mM). mDHFR (38Am) was expressed in an phenylalanine, tryptophan and lysine triple auxotrophic strain (AFWK) transformed with the mutant yeast PheRS and cognate suppressor tRNA. The purified mDHFR (38Am) was digested by modified trypsin. Peptide C, containing one Phe codon, remains the same for all the experimental conditions, indicating normal Phe codons are assigned as Phe. Residue *Z*, the amino acid incorporated in response to the amber codon, is assigned based on the mass units of Peptide A.

Chapter 3

Design of a Bacterial Host for Site-Specific Incorporation of *p*-Bromophenylalanine into Recombinant Proteins

The text in this chapter is reprinted with permission from Kwon, I.; Wang, P.; Tirrell, D. A. *J. Am. Chem. Soc.* **2006**, *128*, 11778-11783, Copyright 2006, American Chemical Society.

Introduction of a yeast suppressor tRNA (ytRNA^{Phe}_{CUA}) and a mutant yeast phenylalanyl-tRNA synthetase (yPheRS (T415G)) into an *E. coli* expression host allows in vivo incorporation of phenylalanine analogs into recombinant proteins in response to amber stop codons. However, high fidelity incorporation of non-natural amino acids is precluded in this system by mischarging of ytRNA^{Phe}_{CUA} with tryptophan (Trp) and lysine (Lys). Here we show that ytRNA^{Phe}_{CUA} and yPheRS can be re-designed to achieve high fidelity amber codon suppression through delivery of p-bromophenylalanine (pBrF). Two strategies were used to reduce misincorporation of Trp and Lys. First, Lys misincorporation was eliminated by disruption of a Watson-Crick base pair between nucleotides 30 and 40 in ytRNA^{Phe}_{CUA}. Loss of this base pair reduces mischarging by the E. coli lysyl-tRNA synthetase. Second, the binding site of yPheRS was re-designed to enhance specificity for pBrF. Specifically, we used the T415A variant, which exhibits five-fold higher activity toward pBrF as compared to Trp in ATP-PPi exchange assays. Combining mutant ytRNA^{Phe}_{CUA} and yPheRS (T415A) allowed incorporation of pBrF into murine dihydrofolate reductase in response to an amber codon with at least 98% fidelity.

Introduction

Non-natural amino acids carrying a wide variety of novel functional groups have been incorporated into recombinant proteins in both prokaryotic and eukaryotic cells.¹⁻²⁴ Although global replacement of natural amino acids with non-natural analogs is useful for many purposes,^{1,16,25,26} there are situations in which single-site substitution by a nonnatural amino acid is required. In such circumstances, a codon must be assigned uniquely to the non-natural amino acid. Amber (UAG), ${}^{2,4,7-9,27-29}$ ochre (UAA), 30,31 opal (UGA) stop codons, 30,32 and four base codons 32,33 have been explored for this purpose. The amber codon has been used most widely, ${}^{2,8,9,27-29}$ because it is the least common stop codon in *E. coli* and because several naturally occurring suppressor tRNAs recognize it efficiently. 34,35 Use of the amber codon to encode non-natural amino acids requires outfitting the cell with an "orthogonal pair" comprising a suppressor tRNA and a cognate aminoacyl-tRNA synthetase (aaRS) that operate independently of the endogenous synthetase-tRNA pairs in *E. coli*. The site-specific incorporation of a non-natural amino acid into recombinant proteins via this strategy was reported by Furter in 1998, ⁹ and the Schultz laboratory has developed powerful selection methods to identify heterologous synthetases and tRNAs. 2,22,32,36,37 The tyrosyl pair derived from the archaebacterium *Methanococcus jannaschii* has been especially useful in this regard. ³⁸

In a complementary approach, rational modification and virtual screening methods have been used to design the amino acid binding pocket of the *E. coli* PheRS (ePheRS).^{5,39} On the basis of the crystal structure of the PheRS (tPheRS) from *Thermus thermophilus*, Safro and colleagues proposed that Val 261 and Ala 314 in the amino acid binding pocket are critical in the discrimination of Phe from its amino acid competitors.⁴⁰ Sequence alignment indicates that Ala 314 in tPheRS corresponds to Ala 294 in ePheRS, and the Hennecke group showed that the substrate specificity of ePheRS can be relaxed by a point mutation at Ala 294. The A294G mutant was shown to enable incorporation of *para*-chlorophenylalanine into recombinant proteins.⁴¹ A subsequent computational simulation, consistent with the Safro prediction, identified two cavity-forming mutations (T251G and A294G) in ePheRS binding pocket. These two mutations led to relaxed substrate specificity and efficient in vivo replacement of Phe by *para*-

acetylphenylalanine (pAcF).⁵ Sequence alignment shows that Thr 415 in yPheRS is equivalent to Thr 251 in ePheRS. We therefore anticipated that the yPheRS (T415G) variant would activate a variety of Phe analogs.⁴² However, when the T415G variant was co-transformed with ytRNA^{Phe}_{CUA} into an *E. coli* host, substantial misincorporation of Trp and Lys was observed. Use of a triple auxotrophic *E. coli* host (Phe, Trp, and Lys) led to good incorporation (85-95%) of a variety of novel aromatic amino acids in response to amber codons,³² but the limited specificity of this expression system prompted us to consider its refinement.

In this report, we describe the design of a bacterial host capable of high fidelity site-specific incorporation of *para*-bromophenylalanine (pBrF) in response to an amber codon. Introduction of aryl halides, such as pBrF^{19,22} or *para*-iodophenylalanine (pIF)^{13,22} into recombinant proteins allows site-specific modification via versatile palladium catalyzed cross-coupling reactions with terminal alkene or alkyne reaction partners.⁴³⁻⁴⁸ In order to achieve high fidelity incorporation of pBrF, two different strategies were applied to reduce misincorporation of Trp and Lys, while providing good yields of recombinant protein. First, Lys misincorporation was eliminated by modification of the sequence of $ytRNA^{Phe}_{CUA}$ to reduce mischarging of $ytRNA^{Phe}_{CUA}$ by eLysRS. Second, the binding site of yPheRS was re-designed to enhance specificity for pBrF. Finally, the combined use of these two strategies provided high fidelity amber codon suppression through delivery of pBrF.

Materials and Methods

Materials. Unless otherwise noted, reagents were obtained from commercial suppliers and were used without further purification. Natural amino acids were from Sigma (St. Louis, MO), and [³H]-amino acids from Amersham Pharmacia Biotech (Piscataway, NJ). pBrF was obtained from Chem-Impex (Wood Dale, IL). Purified total yeast tRNA was purchased from Roche Biochemical (Indianapolis, IN). The nickel-nitrilotriacetic acid affinity column and repressor plasmid pREP4 were from Qiagen (Valencia, CA). *E. coli* strain BLR (RecA⁻ derivative of BL21) was purchased from Novagen (Madison, WI).

Plasmid Construction for Synthetase Expression. Genes encoding the α - and β subunits of yPheRS were amplified from template plasmid pUC-ASab2^{9,49} and inserted between the BamHI and KpnI sites of pQE32 to give pQE32-yPheRS. pQE32-T415G and pQE32-T415A were constructed from pQE32-yPheRS by PCR mutagenesis at position 415 of the α -subunit of yPheRS by use of a QuikChange mutagenesis kit (Stratagene). Two complementary oligonucleotides, 5'-CT ACC TAC AAT CCT TAC GCC GAG CCA TCA ATG GAA ATC-3' for the forward primer and 5'-GAT TTC CAT TGA TGG CTC GGC GTA AGG ATT GTA GGT AG-3' for the reverse primer, were used to introduce the T-to-A mutation at position 415 of the α -subunit of yPheRS. The entire yPheRS gene was verified by DNA sequencing for each of these constructs. Proofreading polymerase Pfx was used in all PCR described in this work. The E. coli *lvsS* gene was amplified by PCR from template plasmid pXLysKS1,⁴⁸ by using two primers, 5'-GCA CTG ACC ATG GCT GAA CAA CAC GCA CAG-3' (which includes an NcoI restriction site) and 5'-GGA CTT CGG ATC CTT TCT GTG GGC GCA TCG C-3' (which carries a *BamH*I restriction site). The resulting DNA was inserted between the *NcoI* and *BamH*I sites of pQE60 to yield pQE60-eLysS. The cloned enzymes contained N-terminal or C-terminal hexa-histidine tags to facilitate protein purification.

Synthetase Expression and Purification. The plasmids pQE32-yPheRS, pQE32-T415G, pQE32-T415A, and pQE60-eLysS were individually co-transformed with a repressor plasmid pREP4 into E. coli strain BLR to form expression strains BLR (pQE32-yFRS), BLR (pQE32-T415G), BLR (pQE32-T415A), and BLR (pQE60-eLysS), respectively. Synthetase expression was conducted in 2xYT media with 100 µg/mL of ampicillin and 35 µg/mL of kanamycin. At an OD of 0.6, expression of each of the synthetase variants was induced with 1 mM IPTG. After four hours, cells were harvested and proteins were purified over a nickel-nitrilotriacetic acid affinity column under native conditions according to the manufacturer's protocol (Qiagen). The imidazole in the elution buffer was removed by a desalting column (Amersham Lifescience) and proteins were eluted into a buffer containing 50 mM Tris-HCl (pH=7.5), 1 mM DTT. Aliquots of proteins were stored in -80 °C in 50% glycerol. Concentrations of the yPheRS variants and eLvsRS, the lysS gene product, were determined by UV absorbance at 280 nm using calculated extinction coefficients⁵⁰ of 99,060 and 29,420 cm⁻¹ M⁻¹ for vPheRS and eLysRS, respectively.

Amino Acid Activation Assay. The amino acid-dependent ATP-PP_i exchange reaction was used to determine the kinetics of activation of amino acid analogs by yPheRS. The reaction buffer consisted of 50 mM N-(2-hydroxethyl) piperazine-N'-(2-ethanesulfonic acid) (potassium-HEPES) (pH=7.6), 20 mM MgCl₂, 1 mM dithiothreitol (DTT), 2 mM adenosine triphosphate (ATP) and 2 mM [³²P]-pyrophosphate (PPi) (NEN Life Science) with specific activity of 10-50 Ci/mol. The amino acid concentration varied from 10 μ M to 2.5 mM and the enzyme concentration varied from 10 nM to 100 nM. Aliquots (20

 μ L) were removed from the reaction solution at various time points and quenched into 500 μ L of buffer solution containing 200 mM NaPP_i, 7% w/v HClO₄ and 3% w/v activated charcoal. The charcoal was sedimented by centrifugation and washed twice with 500 μ L of 10 mM NaPP_i and 0.5% HClO₄ solution. The [³²P]-labeled ATP absorbed on the charcoal was quantified via liquid scintillation methods. The specificity constants were calculated by a nonlinear regression fit of the data to a Michaelis-Menten model.

Plasmid Construction for ytRNA^{Phe} Expression. The mutant yeast amber suppressor tRNA (ytRNA^{Phe}_{CUA}) was constitutively expressed under control of an lpp promoter. The expression cassette for ytRNA^{Phe}_{CUA} was inserted into the repressor plasmid pREP4 to form pREP4 ytRNA^{Phe} as described by Furter.⁹ A mutant yeast suppressor ytRNA^{Phe}_{CUA}30U40G (ytRNA^{Phe}_{CUA} _{UG}) was constructed from ytRNA^{Phe}_{CUA} by use of a QuikChange mutagenesis kit. Two complementary oligonucleotides, designated as primer UG_f (5'- GAA CAC AGG ACC TCC ACA TTT AGA GTA TGG CGC TCT CCC -3') for the forward primer and primer UG r (5'- GGG AGA GCG CCA TAC TCT AAA TGT GGA GGT CCT GTG TTC -3') for the reverse primer, were used to introduce the desired mutations at positions 30 and 40 of the yeast suppressor tRNA. The resulting plasmid carrying the gene encoding ytRNA^{Phe}_{CUA UG} is designated pREP4 ytRNA^{Phe} UG. In order to construct plasmids for in vitro transcription of ytRNA^{Phe}, the ytRNA^{Phe}_{CUA} and ytRNA^{Phe}CUA UG amplified from pREP4 ytRNA^{Phe} genes were and pREP4_ytRNA^{Phe}_UG, respectively. At the end of the tRNA sequence, a *BstN*I site was inserted to allow linearization prior to transcription. A T7 promoter sequence was added for in vitro transcription of ytRNA^{Phe} by T7 RNA polymerase. The following primers were used for PCR: 5'-CTG GGT AAG CTT CGC TAA GGA TCT GCC CTG GTG

CGA ACT CTG-3' (which includes restriction sites for *Hind*III and *BstN*I) and 5'-GAT TAC GGA TTC CTA ATA CGA CTC ACT ATA GCG GAC TTA GCT C-3' (which carries an *EcoR*I restriction site and a T7 promoter sequence). The resulting DNA fragments were introduced between the *Hind*III and *EcoR*I sites of pUC18 to yield pUC18_ytRNA^{Phe}_CUA and pUC18_ytRNA^{Phe}_UG, respectively. In order to facilitate DNA manipulation, a *BstN*I site close to the T7 promoter sequence of pUC18_ytRNA^{Phe}_CUA was removed to increase the size of the DNA fragment containing the ytRNA^{Phe}_{CUA} gene from 180 bp to 500 bp after *BstN*I digestion. Two complementary oligonucleotides, 5'-CGG AAG CAG AAA GTG TAA AGA GCG GGG TGC CTA ATG AGT G-3' for the forward primer and 5'-CAC TCA TTA GGC ACC CCG CTC TTT ACA CTT TAT GCT TCC G-3' for the reverse primer, were used to introduce this mutation.

In Vitro Transcription. Linearized DNAs were prepared by BstNI digestion of pUC18_ytRNA^{Phe}_CUA and pUC18_ytRNA^{Phe}_UG as described previously.⁵¹ In vitro transcription of linearized DNA templates, and purification of transcripts were performed as described previously with minor alterations.⁵² The in vitro transcription of linearized DNA to produce 76-mer tRNA transcripts was performed with the Ambion T7-MEGAshortscript kit. Transcripts were isolated by extraction with 25:24:1 phenol:CHCl₃:isoamyl alcohol. The organic layer was re-extracted with water and a 24:1 CHCl₃:isoamyl alcohol extraction was performed on the aqueous layers. The water layer was then mixed with an equal volume of isopropanol, precipitated overnight at -20°C, pelleted, dried, and re-dissolved in water. Unreacted nucleotides in the tRNA solution

were eliminated using CHROMA SPIN-30 DEPC-H₂O (BD Bioscience) spin columns. Concentrations of the transcripts were determined by UV absorbance at 260 nm.

Aminoacylation Assay. Aminoacylation of wild-type ytRNA^{Phe}_{GAA} with Phe and Trp by vPheRS variants was performed as described earlier.⁵¹ Aminoacylation reactions were carried out in buffer (100 µL) containing 30 mM HEPES (pH=7.45), 15 mM MgCl₂, 4 mM DTT, 25 mM KCl, and 2 mM ATP at 30 °C. Purified yeast total tRNA was used in the assay at a final concentration of 4 mg/mL (ytRNA^{Phe}_{GAA} concentration approximately 2.24 μ M). For aminoacylation with Phe, 13.3 μ M [³H]-Phe (5.3 Ci/mmol) and 80 nM yPheRS variants were used; for aminoacylation with Trp, 3.3 μ M [³H]-Trp (30.0 Ci/mmol) and 160 nM yPheRS variants were used. Aminoacylation of vtRNAPhe transcripts reactions was carried out in buffer (100 µL) containing 100 mM potassium-HEPES (pH=7.4), 10 mM MgCl₂, 1 mM DTT, 0.2 mM EDTA, 2 mM ATP, and 4 units/mL yeast inorganic pyrophosphatase (Sigma) at 37°C for eLysRS. For aminoacylation with Lys, 4 μ M of ytRNA^{Phe} transcript, 1.1 μ M [³H]-Lys (91 Ci/mmol) and 80 nM eLvsRS were used. The tRNAs were annealed before use by heating to 85 °C for 4 min in annealing buffer (60 mM Tris, pH=7.8, 2 mM MgCl₂) followed by slow cooling to room temperature. Reactions were initiated by adding the enzyme and 10 μ L aliquots were quenched by spotting on Whatman filter disks soaked with 5% trichloroacetic acid (TCA). The filters were washed for three 10 min periods in ice-cold 5% TCA, washed in ice-cold 95% ethanol, and counted via liquid scintillation methods.

Strain and Plasmid Construction for in Vivo Incorporation of pBrF. A Phe/Trp double auxotrophic strain (AFW) and a Phe/Trp/Lys triple auxotrophic strain (AFWK) were constructed from the Phe auxotrophic strain AF (K10, Hfr(Cavalli) pheS13rel-1 tonA22 thi T2^R pheA18, trpB114)⁹ by P1 phage-mediated transposon transduction as described previously.⁴² pQE16 (Qiagen) was chosen as the expression plasmid; pQE16 bears a gene encoding the marker protein murine dihydrofolate reducase (mDHFR) outfitted with a C-terminal hexa-histidine tag under control of a bacteriophage T5 promoter and t₀ terminator. A QuikChange mutagenesis kit with two complementary oligonucleotides (5'-CCG CTC AGG AAC GAG TAG AAGTAC TTC CAA AGA ATG-3'; 5'-CAT TCT TTG GAA GTA CTT CTA CTC GTT CCT GAG CGG-3') was used to place an amber codon (TAG) at the 38th position of mDHFR; the resulting plasmid was designated pQE16am. The mutant yPheRS genes T415G and T415A were amplified from pQE32-T415G and pQE32-T415A, respectively, and a constitutive tac promoter with an abolished lac repressor binding site was added upstream of the start codon of this gene.⁹ The resulting expression cassettes were inserted into the PvuII site of pQE16 to yield pQE16am-T415G and pQE16am-T415A.

In Vivo Incorporation of pBrF. The auxotrophic strains AF, AFW, and AFWK were transformed with pQE16am and pREP4_ytRNA^{Phe} vectors to enable pBrF incorporation. mDHFR expression was investigated in 20 mL cultures. The *E. coli* expression strains were grown in M9 minimal medium supplemented with glucose, thiamin, MgSO₄, CaCl₂, 20 amino acids (at 25 μ g/mL), and antibiotics (35 μ g/mL of kanamycin and 100 μ g/mL of ampicillin). When the cultures reached an OD₆₀₀ of 0.8-1.0, cells were sedimented by centrifugation, washed twice with cold 0.9% NaCl, and shifted to expression media

supplemented with 17 amino acids (at 20 μ g/mL), 6 mM of pBrF, and the indicated concentrations of Phe, Trp and Lys. Protein expression was induced by addition of 1 mM IPTG. After four hours, cells were pelleted by centrifugation, and the protein was purified on a Ni-NTA spin column under denaturing conditions according to the supplier's instructions (Qiagen). After purification, expression levels of mDHFR were determined by UV absorbance at 280 nm using a calculated extinction coefficient⁵⁰ of 24,750 cm⁻¹ M⁻¹.

Quantitative Analysis of pBrF Incorporation by Liquid Chromatography Mass Spectrometry (LC-MS). LC-MS and LC/MS/MS analyses of tryptic digests of mDHFR were conducted on a Finnigan LCQ ion trap mass spectrometer equipped with an HPLC pump and ESI probe. Mutant mDHFR was prepared in elution buffer (8 M urea, 100 mM NaH_2PO_4 , 10 mM Tris, pH=4.5). After concentration of the protein by ultracentrifugation (Millipore), 10 µL of the concentrate was diluted into 90 µL of 75 mM NH₄HCO₃ for trypsin digestion. Modified trypsin (1 μ L, Promega, 0.2 μ g/ μ L) was added. The sample was incubated at 37 °C for 2 to 6 h and the reaction was stopped by addition of 12 μ L of 5% trifluoroacetic acid (TFA) solution. Digested peptide solution was subjected to desalting on a C18 Vydac Microspin column (The Nest group) and eluted with 50 µL of 80% acetonitrile/20% 0.1% w/v formic acid. The peptide solution eluted from the Microspin column was dried, redissolved in 10% acetonitrile/90% 0.1% TFA solution, and injected into the HPLC. Peptides were separated on a Magic C₁₈ column (Michrom, 300 Å, 0.3x150 mm) and eluted at a flow rate of 30 µL/min using a gradient of 10-95% of solvent A (90% acetonitrile/10% 0.1 M aqueous acetic acid solution) and solvent B

(2% acetonitrile/98% 0.1 M aqueous acetic acid solution) for 30 min. The column eluent was introduced to the electrospray source and sequencing was carried out by fragmentation of the precursor ion corresponding to the fragment bearing the residue at position 38 of mutant mDHFR.

Results and Discussion

Aminoacylation of ytRNA^{Phe}CUA and ytRNA^{Phe}CUA UG with Lys. When the yeast phenylalanine amber suppressor (ytRNA^{Phe}_{CUA_WT}) was co-expressed with wild-type yPheRS in Furter's E. coli expression strain, 60% of the amber codon sites were decoded as Lys.⁹ Use of the G37A mutant of ytRNA^{Phe}_{CUA WT} (ytRNA^{Phe}_{CUA}) reduced the extent of Lys misincorporation to 5%.⁹ Because insertion of Lys in response to the amber codon is likely a consequence of charging of ytRNA^{Phe}_{CUA} by the *E. coli* lysyl-tRNA synthetase (eLysRS), we modified ytRNA^{Phe}_{CUA} to eliminate cross-reactivity with eLysRS. We focused on the base pair between nucleotides 30 and 40, which is thought to enhance recognition of yeast isoleucine amber suppressor (ytRNA^{Ile}_{CUA}) by eLysRS.⁵³ Buttcher and coworkers showed that introduction of a wobble base pair between nucleotides 30 and 40 reduced charging of ytRNA^{Ile}_{CUA} by eLysRS.⁵³ Similarly, we generated a mutant yeast phenylalanine amber suppressor (ytRNA^{Phe}_{CUA UG}) containing the wobble base pair 30U-40G, and compared the rates of charging of ytRNA^{Phe}_{CUA} and ytRNA^{Phe}_{CUA} ug with Lys by eLysRS (Figure 1).⁵⁴ As expected, the rate of aminoacylation of ytRNA^{Phe}_{CUA UG} was about three-fold lower than that of ytRNA^{Phe}_{CUA} (Figure 1); however, there was only about a 40% reduction in the rate of aminoacylation of ytRNAPhe_{CUA_UG} with Phe as compared to that of ytRNA^{Phe}_{CUA}. Since ytRNA^{Phe}_{CUA} variants are competitively

aminoacylated by either eLysRS or yPheRS, the three-fold reduction in the rate of aminoacylation by eLysRS was expected to reduce the level of Lys-charged ytRNA^{Phe}.

Occupancy of the Amber Site. Occupancy of the amber site in mDHFR-38Am was analyzed by LC-MS/MS (Figure 2). The results are summarized in Table 1. LC-MS/MS has been used successfully for quantitative analysis of protein modification.⁴³ Mutant mDHFR containing the amber codon at the 38th position (mDHFR 38Am) was expressed in E. coli strains (AF, AFW, and AFWK) co-transformed with pQE16_yPheRS (T415G) and either pREP4_ytRNA^{Phe} CUA or pREP4_ytRNA^{Phe}_UG. We focused on Peptide 38 (residues 26-39; NGDLPWPPLRNEAmK) which contains the amber site. Peptide 38 variants containing Trp and pBrF are designated Peptides W38 and Z38, respectively. When Lys is incorporated in response to the amber codon, there are two consecutive lysines at the C-terminus of Peptide 38, and the C-terminal Lys can be further cleaved by trypsin. fully cleaved (NGDLPWPPLRNEK) partially The and cleaved (NGDLPWPPLRNEKK) variants of Peptide K38 were designated K38S (short) and K38L (long), respectively. Peptides K38S, K38L, W38 and Z38 were readily separated by liquid chromatography (Figure 2a), and the relative yields of the four variants were determined by comparing the integrated areas of the corresponding chromatographic signals. Furthermore, the fragment ion masses could be unambiguously assigned (data not shown), confirming the identity of the amino acid inserted in response to the amber codon. The amplitude of the most intense signal in the chromatograms in Figure 2 is ca. 1×10^8 (instrument-dependent arbitrary units) compared to a noise level of ca. 5×10^5 , suggesting that misincorporation of natural amino acids can be analyzed at the 0.5% level.

Reduced Lys Misincorporation by ytRNA^{Phe}_{CUA_UG}. When mDHFR(38Am) was expressed in a triple auxotrophic *E. coli* host (AFWK) outfitted with ytRNA^{Phe}_{CUA} and yPheRS (T415G) in medium supplemented with 0.03 mM Phe, 0.01 mM Trp, 1 mM Lys and 6 mM pBrF, Lys misincorporation was observed at a level of 7.6% (Figure 2a). The level of Lys misincorporation could be reduced to 2.7% by restricting the Lys concentration to 0.01 mM (Table 1). However, when ytRNA^{Phe}_{CUA_UG} was co-transformed into *E. coli* strain AFWK with yPheRS (T415G), Lys misincorporation was not detected, even in media supplemented with 1.0 mM Lys (Figure 2b). Considering the detection limit of the ESI MS detector, this establishes an upper limit of 0.5% Lys misincorporation. Similar results were obtained with Phe auxotrophic strain AF outfitted with ytRNA^{Phe}_{CUA_UG} (Figure 2e); the Lys auxotrophic strain is not required to eliminate Lys misincorporation.

Redesign of Phenylalanyl-tRNA Synthetase. Although use of ytRNA^{Phe}_{CUA_UG} eliminated Lys at the amber site, ca. 10% Trp misincorporation was still detected (Figure 2b). Trp misincorporation could result from mischarging of ytRNA^{Phe}_{CUA} with Trp either by yPheRS or by *E. coli* tryptophanyl-tRNA synthetase (eTrpRS). Because misincorporation of Trp was not detected in the absence of yPheRS (T415G), mischarging of ytRNA^{Phe}_{CUA} by eTrpRS is probably negligible. However, the Trp-dependent ATP-PPi exchange rate of yPheRS (T415G) was comparable to that observed for pBrF (Table 2). Therefore, mischarging of ytRNA^{Phe}_{CUA} with Trp by yPheRS (T415G) is the more likely explanation for Trp misincorporation at the amber codon. We speculated that yPheRS (T415A) might exclude Trp.

ATP-PPi Exchange and Aminoacylation of ytRNA^{Phe} **Catalyzed by yPheRS Variants.** The kinetics of amino acid-dependent ATP-PPi exchange catalyzed by three variants of yPheRS (wild-type, T415G, and T415A) were measured; the kinetic parameters are reported in Table 2. Wild-type yPheRS activates neither Trp nor pBrF, whereas the T415G variant shows ten-fold higher activity toward pBrF and Trp as compared to Phe. As expected, the Trp-dependent ATP-PPi exchange rate of yPheRS (T415A) is ca. thirty-fold lower than that of yPheRS (T415G). In contrast, the effects of the G415A mutation on the rates of activation of Phe and pBrF are much smaller (Table 2). Aminoacylation of ytRNA^{Phe}_{CUA} with Phe was comparable for yPheRS (T415G) and for yPheRS (T415A) (Figure 3a); however, aminoacylation of ytRNA^{Phe}_{CUA} with Trp by yPheRS (T415A) is much slower than that by yPheRS (T415G) (Figure 3b).

The improved selectivity of yPheRS (T415A) with respect to exclusion of Trp may be understood at least in part on the basis of the side-chain volumes of the amino acid substrates and the active-site residues of the synthetase. Replacement of Gly415 by Ala reduces the volume of the substrate-binding site by ca. 16 Å^{3,55} Occlusion of the binding site reduces the binding affinity of both Trp and pBrF, but because Trp is the larger of the two amino acids (by ca. 5 Å³), it suffers the larger reduction in affinity. The G415A mutation increases K_m by a factor of ca. 17 for Trp as compared to ca. 7 for pBrF (Table 2).

High Fidelity Site-Specific Incorporation of pBrF. Use of expression hosts outfitted with yPheRS (T415A) reduced misincorporation of Trp at the amber site to levels below the limit of detection (Figure 2c-e). Our initial studies used the triple auxotrophic

expression host AFWK. However, construction of the triple auxotroph was timeconsuming and the limited cellular pools of several of the natural amino acids reduced protein yield. In order to overcome these problems, a single (Phe) auxotrophic expression host (AF) outfitted with the yPheRS (T415A) and ytRNA^{Phe}_{CUA_UG} was tested for pBrF incorporation. High fidelity incorporation of pBrF in response to the amber codon was observed (Figure 2e and Table 1), indicating that the triple auxotrophic expression host is not necessary. Furthermore, the AF expression host outfitted with yPheRS (T415A) and ytRNA^{Phe}_{CUA_UG} yielded twice as much protein as the triple auxotroph (4.3 mg/L vs. 2.0 mg/L) (Table 1).

Conclusions

We describe the engineering of a bacterial expression host that allows high fidelity incorporation of pBrF into recombinant proteins in response to amber stop codons. Rational modification of the yeast amber suppressor tRNA eliminated cross-reactivity with the *E. coli* LysRS, while the substrate specificity of yPheRS was enhanced by a point mutation in the amino acid binding pocket. When the modified yeast amber suppressor tRNA and the mutant yPheRS were co-expressed in a Phe auxotrophic *E. coli* expression host, at least 98% of the amber sites in full-length recombinant mDHFR were occupied by pBrF. Aryl bromides at programmed positions should enable chemoselective ligation of proteins with terminal alkene or alkyne reaction partners.⁴³

Acknowledgments

We thank Dr. P. Plateau for providing the plasmid pXLysKS1 and Dr. M. Shahgholi for assistance with mass spectrometry. We are grateful to Dr. K. Shin, Dr. J.

Petersson, and Dr. D. Beene for help with in vitro transcription of tRNA. We thank I. C. Tanrikulu for help with analysis of the PheRS crystal structure and for review of this manuscript. We also thank Dr. A. J. Link and K. E. Beatty for discussion and valuable comments on this project. This work was supported by National Institutes of Health grant GM62523.

References

- Beatty, K. E.; Xie, F.; Wang, Q.; Tirrell, D. A. J. Am. Chem. Soc. 2005, 127, 14150-14151.
- Bose, M.; Groff, D.; Xie, J. M.; Brustad, E.; Schultz, P. G. J. Am. Chem. Soc. 2006, 128, 388-389.
- 3. Budisa, N.; Pal, P. P. Biol. Chem. 2004, 385, 893-904.
- Chin, J. W.; Cropp, T. A.; Anderson, J. C.; Mukherji, M.; Zhang, Z. W.; Schultz, P. G. Science 2003, 301, 964-967.
- Datta, D.; Wang, P.; Carrico, I. S.; Mayo, S. L.; Tirrell, D. A. J. Am. Chem. Soc. 2002, 124, 5652-5653.
- Dedkova, L. M.; Fahmi, N. E.; Golovine, S. Y.; Hecht, S. M. J. Am. Chem. Soc. 2003, 125, 6616-6617.
- Deiters, A.; Cropp, T. A.; Mukherji, M.; Chin, J. W.; Anderson, J. C.; Schultz, P. G. J. Am. Chem. Soc. 2003, 125, 11782-11783.
- 8. Deiters, A.; Geierstanger, B. H.; Schultz, P. G. ChemBioChem 2005, 6, 55-58.
- 9. Furter, R. Protein Sci. 1998, 7, 419-426.

- 10. Hendrickson, T. L.; de Crecy-Lagard, V.; Schimmel, P. Annu. Rev. Biochem. 2004, 73, 147-176.
- 11. Ibba, M.; Stathopoulos, C.; Soll, D. Curr. Biol. 2001, 11, R563-R565.
- 12. Kiick, K. L.; Weberskirch, R.; Tirrell, D. A. FEBS Lett. 2001, 505, 465-465.
- 13. Kirshenbaum, K.; Carrico, I. S.; Tirrell, D. A. ChemBioChem 2002, 3, 235-237.
- 14. Kwon, I.; Kirshenbaum, K.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 7512-7513.
- 15. Link, A. J.; Mock, M. L.; Tirrell, D. A. Curr. Opin. Biotechnol. 2003, 14, 603-609.
- 16. Link, A. J.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 11164-11165.
- 17. Link, A. J.; Vink, M. K. S.; Tirrell, D. A. J. Am. Chem. Soc. 2004, 126, 10598-10602.
- Sakamoto, K.; Hayashi, A.; Sakamoto, A.; Kiga, D.; Nakayama, H.; Soma, A.; Kobayashi, T.; Kitabatake, M.; Takio, K.; Saito, K.; Shirouzu, M.; Hirao, I.; Yokoyama, S. *Nucleic Acids Res.* 2002, *30*, 4692-4699.
- 19. Sharma, N.; Furter, R.; Kast, P.; Tirrell, D. A. FEBS Lett. 2000, 467, 37-40.
- 20. Tang, Y.; Tirrell, D. A. J. Am. Chem. Soc. 2001, 123, 11089-11090.
- 21. Wang, L.; Brock, A.; Herberich, B.; Schultz, P. G. Science 2001, 292, 498-500.
- 22. Wang, L.; Xie, J. M.; Deniz, A. A.; Schultz, P. G. J. Org. Chem. 2003, 68, 174-176.
- 23. Xie, J. M.; Schultz, P. G. Curr. Opin. Chem. Biol. 2005, 9, 548-554.
- 24. Zhang, Z. W.; Gildersleeve, J.; Yang, Y. Y.; Xu, R.; Loo, J. A.; Uryu, S.; Wong, C. H.; Schultz, P. G. *Science* 2004, *303*, 371-373.
- 25. Lim, V. I. J. Mol. Biol. 1995, 252, 277-282.
- 26. Tang, Y.; Ghirlanda, G.; Petka, W. A.; Nakajima, T.; DeGrado, W. F.; Tirrell, D. A. *Angew. Chem.Int. Ed.* **2001**, *40*, 1494-1496.
- Cload, S. T.; Liu, D. R.; Froland, W. A.; Schultz, P. G. Chem. Biol. 1996, 3, 1033-1038.

- 28. Cornish, V. W.; Benson, D. R.; Altenbach, C. A.; Hideg, K.; Hubbell, W. L.; Schultz,
 P. G. *Proc. Natl. Acad. Sci. U. S. A.* **1994**, *91*, 2910-2914.
- Mendel, D.; Cornish, V. W.; Schultz, P. G. Annu. Rev. Biophys. Biomol. Struct. 1995, 24, 435-462.
- Kohrer, C.; Sullivan, E. L.; RajBhandary, U. L. Nucleic Acids Res. 2004, 32, 6200-6211.
- Kohrer, C.; Yoo, J. H.; Bennett, M.; Schaack, J.; RajBhandary, U. L. Chem. Biol.
 2003, 10, 1095-1102.
- 32. Anderson, J. C.; Schultz, P. G. Biochemistry 2003, 42, 9598-9608.
- Hohsaka, T.; Muranaka, N.; Komiyama, C.; Matsui, K.; Takaura, S.; Abe, R.;
 Murakami, H.; Sisido, M. *FEBS Lett.* 2004, *560*, 173-177.
- Thorbjarnardottir, S.; Uemura, H.; Dingermann, T.; Rafnar, T.; Thorsteinsdottir, S.;
 Soll, D.; Eggertsson, G. J. Bacteriol. 1985, 161, 207-211.
- 35. Yoshimura, M.; Inokuchi, H.; Ozeki, H. J. Mol. Biol. 1984, 177, 627-644.
- Liu, D. R.; Magliery, T. J.; Pasternak, M.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 1997, 94, 10092-10097.
- 37. Santoro, S. W.; Wang, L.; Herberich, B.; King, D. S.; Schultz, P. G. *Nat. Biotechnol.*2002, 20, 1044-1048.
- 38. Cropp, T. A.; Schultz, P. G. Trends Genet. 2004, 20, 625-630.
- Wang, P.; Vaidehi, N.; Tirrell, D. A.; Goddard, W. A. J. Am. Chem. Soc. 2002, 124, 14442-14449.
- 40. Fishman, R.; Ankilova, V.; Moor, N.; Safro, M. Acta Crystallogr. D. Biol. Crystallogr. 2001, 57, 1534-44.
- 41. Ibba, M.; Kast, P.; Hennecke, H. Biochemistry 1994, 33, 7107-7112.

- 42. Wang, P., Ph.D. thesis, California Institute of Technology, Pasadena, 2003.
- Kodama, K.; Fukuzawa, S.; Nakayama, H.; Kigawa, T.; Sakamoto, K.; Yabuki, T.; Matsuda, N.; Shirouzu, M.; Takio, K.; Tachibana, K.; Yokoyama, S. *ChemBioChem* 2006, 7, 134-139.
- 44. Nguefack, J. F.; Bolitt, V.; Sinou, D. Tetrahedron Lett. 1996, 37, 5527-5530.
- Sakakibara, Y.; Ido, Y.; Sasaki, K.; Sakai, M.; Uchino, N. Bull. Chem. Soc. Jpn. 1993, 66, 2776-2778.
- 46. Schoenbe.A; Heck, R. F. J. Org. Chem. 1974, 39, 3327-3331.
- 47. Yoneyama, M.; Kakimoto, M.; Imai, Y. Macromolecules 1988, 21, 1908-1911.
- 48. Yoneyama, M.; Konishi, T.; Kakimoto, M. A.; Imai, Y. Makromol. Chem., Rapid Commun. 1990, 11, 381-386.
- 49. Sanni, A.; Walter, P.; Ebel, J. P.; Fasiolo, F. Nucleic Acids Res. 1990, 18, 2087-2092.
- 50. Gill, S. C.; Vonhippel, P. H. Anal. Biochem. 1989, 182, 319-326.
- 51. Sampson, J. R.; Uhlenbeck, O. C. Proc. Natl. Acad. Sci. U. S. A. 1988, 85, 1033-1037.
- 52. Nowak, M. W.; Gallivan, J. P.; Silverman, S. K.; Labarca, C. G.; Dougherty, D. A.; Lester, H. A., In vivo incorporation of unnatural amino acids into ion channels in *Xenopus oocyte* expression system. In *Ion Channels, Pt B*, **1998**, Vol. 293, pp 504-529.
- Buttcher, V.; Senger, B.; Schumacher, S.; Reinbolt, J.; Fasiolo, F. Biochem. Biophys. Res. Commun. 1994, 200, 370-377.
- 54. Nishikawa and coworkers used an alternative strategy to reduce misacylation of yeast amber suppressor tRNATyr by eLysRS. Fukunaga, J.; Yokogawa; T., Ohno; Nishikawa, K. J. Biochem. 2006, 139, 689-696. In.

55. Volumes were estimated as solvent-excluded volumes (1.4 Å probe) by using Chem3D Pro 5.0 (CambridgeSoft: Cambridge, MA)



Figure 1: Aminoacylation of ytRNA^{Phe}_{CUA} (\blacksquare) and ytRNA^{Phe}_{CUA_UG} (\bigcirc) with Lys by eLysRS. The reaction was carried out at 37°C in mixtures containing 2 mM ATP, 1.1 μ M [³H]-Lys, 80 nM eLysRS, and 4 μ M of ytRNA^{Phe}_{CUA} or ytRNA^{Phe}_{CUA_UG}.



Figure 2: LC-MS chromatograms of tryptic digests of mDHFR. Peptide 38 (residues 26-39; NGDLPWPPLRNEAmK; Am indicates an amber codon) contains an amber codon at the 38th position. Peptide 38 variants containing Lys (Peptides K38S and K38L), Trp (Peptide W38), and pBrF (Peptide Z38) were se*para*ted and detected by MS. mDHFRs were synthesized in a triple auxotrophic expression host outfitted with: (a) ytRNA^{Phe}_{CUA} and yPheRS (T415G); (b) ytRNA^{Phe}_{CUA_UG} and yPheRS (T415G); (c) ytRNA^{Phe}_{CUA} and yPheRS (T415A); (d) ytRNA^{Phe}_{CUA_UG} and yPheRS (T415A); or in a single auxotrophic strain with ytRNA^{Phe}_{CUA_UG} and yPheRS (T415A) (e). The expression media were supplemented with 6.0 mM pBrF, 0.01 mM Trp, 1.0 mM Lys, 0.03 mM Phe (a, b and c) or 0.01 mM Phe (d, e and f), and 25 µg/mL of 17 amino acids.



(**A**) and yPheRS (T415A) (**•**). Reactions were carried out at 30°C in mixtures containing 2.24 μ M ytRNA^{Phe}_{GAA}, 2 mM ATP, and either (a) 13.3 μ m [³H]-Phe and 80 nM yPheRS variants or (b) 3.3 μ m [³H]-Trp and 160 nM yPheRS variants in aminoacylation buffer.

Figure 3: Charging of Phe (a) and Trp (b) by wild-type yPheRS (■), yPheRS (T415G)
Host strain	yPheRS	ytRNA ^{Phe}	Occupan	Occupancy of amber sites (%)		
			Lys	Trp	pBrF	$(mg/L)^a$
AFWK ^b	T415G	CUA ^c	7.6	9.5	83	1.4 ± 0.4
AFWK	T415G	CUA	2.7 ^d	10.2	87	1.5 <u>±0.2</u>
AFWK	T415G	CUA_UG ^e	\mathbf{ND}^{f}	10.4	90	1.4 ± 0.1
AFWK	T415A	CUA	2.7	ND	97	3.1 ± 0.3
AFWK	T415A	CUA_UG	ND	ND	> 98	2.0 ± 0.2
AF^{g}	T415A	CUA_UG	ND	ND	> 98	4.3 ± 0.4

Table 1: Occupancy of amber sites and expression yields.

^a Volumetric yields are given as mg of purified mDHFR_38Am per liter of culture volume.

^b Triple (Phe, Lys, and Trp) auxotrophic *E. coli* strain.

^{c,e} The ytRNA^{Phe}_{CUA} and ytRNA^{Phe}_{CUA_UG} are designated as CUA and CUA_UG, respectively.

^d 0.01 mM of Lys was supplemented into the media rather than 1.0 mM.

^fNot detected in either LC/MS or MALDI analyses.

^g Single (Phe) auxotrophic *E. coli* strain.

Table 2: Kinetic parameters for ATP-PPi exchange of amino acids by the wild-typeyPheRS, yPheRS (T415G), and yPheRS (T415A) variant.

Amino Acid	Enzyme	$K_{m}\left(\mu M ight)$	k_{cat} (s ⁻¹)	$\begin{array}{c} k_{cat}/K_{m} \\ (M^{-1}s^{-1}) \end{array}$	$\frac{k_{cat}}{(rel)^a}$
Phe	T415A	185±56	0.207±0.02	1,163±240	2.1
Trp	T415A	913±310	0.129±0.01	152.4±49	0.28
pBrF	T415A	255±73	0.217±0.03	892±220	1.6
Phe	T415G	499±51	0.278±0.06	553.0±79	1.0
Trp	T415G	55.0±24	0.261±0.08	5,001±1,000	9.0
pBrF	T415G	36.3±5.0	0.211±0.09	6,116±3,400	11
Phe	Wild-type	29.2±9.8	0.302±0.13	10,669±3,700	19
pBrF	Wild-type	ND^b	ND	ND	ND

^a Relative to k_{cat}/K_m for Phe by T415G.

^b Not detected.

Chapter 4

Site-Specific Incorporation of Tryptophan Analogs into Recombinant Proteins in Vivo

Abstract

An expansion of the number of genetically encoded tryptophan (Trp) analogs in Escherichia coli has been restricted by two limitations. First, Trp analogs inactive in the native E. coli translation system could not be incorporated into proteins. Second, most of the previous attempts to incorporate Trp analogs have been limited to global replacement of all Trp residues throughout a target protein. Here we show that a rationally designed yeast phenylalanyl-tRNA synthetase (yPheRS (T415G)) efficiently activated four Trp analogs: 6-chlorotrytophan (6ClW), 6-bromotryptophan (6BrW), 5-bromotrytophan (5BrW), and benzothienylalanine (BT), which were not utilized by the endogenous E. coli translational system. Introduction of the yPheRS (T415G) and a mutant yeast phenylalanine amber suppressor (ytRNA^{Phe}_{CUA UG}) into an *E. coli* expression host allowed the site-specific incorporation of three Trp analogs: 6ClW, 6BrW, and BT, into murine dihydrofolate reductase in response to amber stop codons with at least 98% fidelity. Subsequently, all three Trp analogs were introduced at the Trp66 position in the chromophore of a cyan fluorescent protein (CFP) to investigate changes in spectral properties. CFP variants containing Trp analogs showed a blue-shift in the fluorescence emission peak as well as in the absorption maximum. In particular, a CFP variant with BT featured an unusually large Stokes' shift (56 nm). As shown in this report, an expanded set of the genetically encoded Trp analogs would enable the design of new proteins with novel properties.

Introduction

Tryptophan (Trp) is an attractive target for protein engineering for several reasons. First, it is the main source of UV absorbance and fluorescence of proteins. Second, Trp is involved in various interactions in proteins, such as ligand binding¹⁻³ and DNA-protein interaction.^{4,5} Third, numerous Trp analogs are available through diverse indole chemistry. Fourth, it is the rarest amino acid,⁶ and so its substitution by other amino acids minimally perturbs structure.⁷ Ever since 7-azatryptophan and 2-azatryptophan were first incorporated into proteins in *E. coli* in the 1950s.^{8,9} many researchers have tried to replace Trp with Trp analogs. In the last few decades, a residue-specific incorporation method utilizing a Trp auxotrophic strain realized the incorporation of various Trp analogs containing fluoro, hydroxyl, methyl, amino, selenophene, and thienyl functional groups.¹⁰⁻¹² Fluorinated Trp analogs, in particular, have been used in ¹⁹F-NMR studies.^{13,14} Trp analogs containing selenophene and thienyl functional groups were used in the X-ray crystallography of proteins.^{15,16} Aminotryptophan was used to design protein-based pH sensors¹⁷ and greatly changed the spectral properties of fluorescent proteins.¹⁸

Although diverse Trp analogs have been introduced into proteins, further expansion of the genetic code for Trp has been restricted by two limitations. First, Trp analogs incorporated into proteins in *E. coli* have so far been limited to the substrates for the native *E. coli* translation system. Therefore, a large number of Trp analogs, such as benzothienylalanine (BT), 5-methyltryptophan, 6-methyltryptophan, 6-aminotryptophan, and 5-cyanotryptophan, were found not to be incorporated into proteins.¹⁰ Recently, several research groups have devised various methods to introduce translationally

inactive nonnatural amino acids into recombinant proteins in *E. coli*.^{10,12,19} Overexpression of aminoacyl-tRNA synthetase (aaRS),^{20,21} utilization of attenuated editing mutant aaRS,^{22,23} rational design of the binding pocket of aaRS,^{24,25} and screening of an aaRS library^{26,27} have been effective in expanding the genetic code for leucine, isoleucine, valine, methionine, phenylalanine, and tyrosine. Recently, Schultz and colleagues modified *Bacillus stearothermophilus* trytophanyl-tRNA synthetase (*Bs*TrpRS) to incorporate 5-hydroxytryptophan into a recombinant protein in mammalian cells.²⁸ However, until now there have been no reports about modification of substrate specificity of aaRS to expand the number of genetically encoded Trp analogs in *E. coli*. In this report, we have shown that the rational design of the binding pocket of yPheRS can expand the genetic code for Trp in *E. coli*.

Second, most of the previous attempts to incorporate Trp analogs have been limited to the global replacement of Trp throughout a protein. Although Trp is the rarest amino acid, the replacement of Trp with Trp analogs at non-permissive sites could result in the loss of function. For example, the replacement of three Trp residues in barstar with 4-aminotryptophans greatly reduced its stability due to increased polarity of 4-aminotryptophan at position 53 inside the protein core.¹¹ In order to overcome such a limitation, we have explored the feasibility of site-specific incorporation of Trp analogs into recombinant proteins in *E. coli*, which would allow incorporation of Trp analogs only at permissive sites. In 1998, Furter showed site-specific incorporation of *p*-fluorophenylalanine into murine dihydrofolate reductase (mDHFR) in *E. coli* through the use of an orthogonal yPheRS/ytRNA^{Phe} pair.²⁹ More recently, Schultz and colleagues have developed powerful screening methods to change the substrate specificity of

tyrosyl-tRNA synthetase (*mj*TyrRS) derived from *Methanococcus jannaschii* toward nonnatural amino acids.^{19,26,30} *E. coli* strains equipped with the screened orthogonal *mj*TyrRS/*mj*tRNA^{Tyr} pairs have allowed incorporation of diverse Phe and Tyr analogs into proteins in a site-specific manner. Tirrell and colleagues refined the orthogonality of the yPheRS/ytRNA^{Phe} pair and the specificity of yPheRS to achieve high fidelity incorporation of *p*-bromophenylalanine into proteins *in vivo*.²⁵

In this report, we describe the design of a bacterial host to achieve site-specific incorporation of Trp analogs into a recombinant protein in response to an amber stop codon in *E. coli*. Since no orthogonal tryptophanyl-tRNA synthetase/tRNA^{Trp} pairs have been reported for *E. coli* translation systems, we needed an alternate orthogonal pair. Our previous studies showed that a rationally designed vPheRS (T415G) variant efficiently activates Trp as well as various Phe analogs.^{25,31} We hypothesized that the relaxed substrate specificity of yPheRS (T415G) may allow it to recognize a variety of Trp analogs. Therefore, a mutant vPheRS (T415G) and a mutant yeast amber suppressor $(ytRNA^{Phe}_{CUA UG})^{32}$ were introduced into an *E. coli* expression strain to insert Trp analogs at a specific site in a protein. Four Trp analogs, BT, 6ClW, 6BrW, and 5BrrW, were examined for incorporation in vivo. In our preliminary studies, none of the four Trp analogs tested were incorporated at UGG codons, even in a Trp auxotrophic E. coli strain under a Trp-depleted condition (data not shown). BT is isosteric to Trp and so might result in minimal perturbation of a protein structure upon the replacement of Trp. Introduction of aryl halides into recombinant proteins can allow for site-specific modification through a palladium-catalyzed coupling reactions with alkenes and alkynes.^{33,34} Bromine and iodine, once introduced into proteins, can facilitate the phasing of crystallographic data and therefore can be useful in X-ray diffraction studies of protein structures.³⁵ As expected, introduction of an orthogonal yeast PheRS (T415G) $/ytRNA^{Phe}_{CUA_UG}$ pair into *E. coli* expression strains enabled high-fidelity (ca. 98%) incorporation of BT, 6BrW, and 6ClW into mDHFR in a site-specific manner.

The *E. coli* strain, equipped with an engineered orthogonal pair, has been used to change the spectral properties of fluorescent proteins. Considering that Trp is a main contributor to UV absorbance and the fluorescence of proteins, it is not surprising that the replacement of Trp with other amino acids resulted in changes in the spectral properties of proteins in many different ways. Schultz and his colleagues have characterized the spectral properties of green fluorescent protein (GFP) variants containing various Phe analogs in the chromophore.³⁶ Fluorescence emission peaks of all GFP variants were blue-shifted to some extent, depending on the electron-donating ability of the Phe analogs. Similarly, Budisa and his colleagues replaced Trp residues in either GFP or the cyan fluorescent protein (CFP) with translationally active Trp analogs.¹¹ In particular, the fluorescence emission peak of 4-aminotryptophan in the chromophore of a fluorescent protein was red-shifted significantly.¹⁸ In this report, we have explored whether we could design novel spectral classes of fluorescent proteins containing Trp analogs that would be translationally inactive for the endogenous E. coli translation system. Specifically, we replaced a Trp residue in the chromophore of a CFP variant (CFP6) with 6ClW, 6BrW, and BT to generate CFP6 6ClW, CFP6 6BrW, and CFP6 BT, respectively. Similar to halogenated Phe analogs, fluorescence emission peaks of CFP6 6ClW and CFP6 6BrW were blue-shifted. However, quantum yields of halogenated Trp analogs were 3- to 4-fold higher than their Phe counterparts. CFP6 BT exhibited a larger Stokes' shift (56 nm) than

that of CFP6 (ca. 40 nm).

Materials and Methods

Materials. Amino acids **1-2** were obtained from Sigma (St. Louis, MO). **3** and **4** were purchased from Biosynth International (Naperville, IL). **5** and **6** were from Aldrich (St. Louis, MO) and Chem-Impex (Wood Dale, IL), respectively. [³²P]-labeled sodium pyrophosphate was purchased from NEN Life Science (Boston, MA).

Amino Acid Activation Assay. The kinetics of activation of amino acids by the yPheRS (T415G) variant were determined by the amino acid dependent adenosine triphosphate (ATP)-[³²P]-pyrophosphate (PP_i) exchange reaction. The plasmid construction, expression, and purification of the yPheRS (T415G) variant with C-terminal hexa-histidine tag were described previously.²⁵ The assay buffer contained 50 mM *N*-(2-hydroxethyl) piperazine-*N*-(2-ethanesulfonic acid) (potassium-HEPES) (pH=7.6), 20 mM MgCl₂, 1 mM dithiothreitol (DTT), 2 mM ATP, and 2 mM [³²P]- PP_i. The amino acid concentrations varied from 100 nM to 2.5 mM and the enzyme concentration varied from 100 nM to 400 nM. Aliquots (20 μ L) of solution were quenched at various time points in 500 μ L of buffer solution containing 200 mM NaPP_i, 7% w/v HClO₄ and 3% w/v activated charcoal. The charcoal was centrifuged and washed three times with 500 μ L of 10 mM NaPP_i and 0.5% HClO₄ solution. The [³²P]-labeled ATP absorbed on the charcoal was counted via liquid scintillation methods. The kinetic parameters were calculated by a nonlinear regression fit of the data to a Michaelis-Menten model.

Strain and Plasmid Construction for in Vivo Incorporation Assays. pQE16am-T415G²⁵ encodes a murine dihydrofolate reductase (mDHFR) outfitted with a C-terminal hexa-histidine tag and an amber codon at position 38. The GFP_{UV} gene was cloned from pGFPuv (Clontech) and inserted into the PstI site of pQE9 (Qiagen) to yield a pQE9 GFP_{UV} plasmid. A series of QuikChange mutations (Stratagene) were performed on GFP_{UV} gene to generate a GFP6 gene containing four sequence changes (F64L, S65T, S99F, and T153M) (see Chapter 6). Either an amber codon (UAG) or a tryptophan codon (UGG) was inserted at position 66 of GFP6 by QuikChange mutagenesis using two complementary pairs of oligonucleotides, Am66 f (5'-CTT GTC ACT ACT CTG ACC TAG GGT GTT CAA TGC TTT TCC-3')/Am66 r (5'-GGA AAA GCA TTG AAC ACC CTA GGT CAG AGT AGT GAC AAG-3') and W66 f (CTT GTC ACT ACT CTG ACC TGG GGT GTT CAA TGC TTT TCC-3')/W66 r (5'-GGA AAA GCA TTG AAC ACC CCA GGT CAG AGT AGT GAC AAG-3'), to yield pQE9 CFP6 66Am and pQE9 CFP6, respectively. The entire expression cassette from either CFP6 66Am or CFP6 was inserted between the AatII and NheI sites of pQE16am-T415G to generate pQE9_CFP6_66Am-T415G or pQE9_CFP6-T415G, respectively. pREP4 ytRNA^{Phe} UG was used to constitutively express a mutant yeast amber suppressor tRNA (ytRNA^{Phe}_{CUA UG}) shows minimal cross-reactivity with the *E. coli* lysyl-tRNA synthetase.²⁵ A Phe/Trp double auxotrophic strain (AFW) was derived from the Phe auxotrophic strain AF (K10, Hfr(Cavalli) pheS13rel-1 tonA22 thi T2^R pheA18, trpB114) by P1 phage-mediated transposon transduction as described previously.^{25,31} The double auxotrophic strain, AFW, was co-transformed with either pQE16am-T415G or pQE9 CFP6 66Am-T415G, and with pREP4 ytRNA^{Phe} UG to yield expression strains

AFW[pQE16am-T415G/pREP4_ytRNA^{Phe}_UG] or AFW[pQE9_CFP6_66Am-T415G/pREP4_ytRNA^{Phe}_UG], respectively. DH10B (Invitrogen) *E. coli* strain was co-transformed with pQE9_CFP6-T415G and pREP4_ytRNA^{Phe}_UG to generate DH10B[pQE9_CFP6-T415G/pREP4_ytRNA^{Phe}_UG].

In Vivo Incorporation Assays. *E. coli* expression strains were grown in M9 minimal medium supplemented with 0.4% glucose, 35 mg/L thiamin, 1mM MgSO₄, 1mM CaCl₂, 20 amino acids (at 25 mg/L), 35 mg/L kanamycin, and 200 mg/L ampicillin. The overnight culture was diluted 20-fold in fresh M9 minimal medium. The cells were grown to an OD₆₀₀ between 0.9 and 1.0, sedimented by centrifugation, and washed twice with cold 0.9% NaCl. The cell pellet was resuspended in fresh M9 minimal medium containing 18 amino acids (at 25 mg/L), 3 mM analog of interest, and the indicated concentrations of Phe and Trp. After 10 min, 1 mM of isopropyl- β -D-thiogalactopyranoside (IPTG) was added to induce protein expression. After 4 hours, cells were pelleted and kept at -80 °C.

A different procedure was used for expression strain DH10B[pQE9_CFP6-T415G/pREP4_ytRNA^{Phe}_UG]. A single colony was inoculated into 5 mL of 2xYT medium and grown overnight. 5 mL of the overnight culture was transferred to 100 mL of fresh 2xYT medium. When the cultures reached an OD₆₀₀ of 0.6, 1 mM IPTG was added to induce protein expression. After 3 hours, cells were sedimented and stored at -80 °C. Whole cell lysates were analyzed by SDS-PAGE.

Quantitative Analysis of Analog Incorporation by Liquid Chromatography Mass **Spectrometry (LC-MS).** Quantitative analysis of amber codon occupancy by various amino acids was performed by LC-MS as described previously.²⁵ In a minor alteration. mutant mDHFR was purified on a Ni-NTA spin column under denaturing conditions according to the supplier's instructions (Qiagen). After purification, mutant mDHFR concentrations were determined by UV absorbance at 280 nm using a calculated extinction coefficient³⁷ of 24,750 cm⁻¹ M⁻¹. After concentration of the protein by ultrafiltration (Millipore), 10 µL of the solution was diluted into 90 µL of 75 mM $(NH_4)_2CO_3$ for trypsin digestion. 1 µL of modified trypsin (Promega, 0.2 µg/µL) was added. The samples were incubated for 2 hrs at 37 °C. The reaction was quenched by addition of 13 µL of 5% trifluoroacetic acid (TFA) solution. The solution was then directly subjected to LC-MS/MS analysis. LC-MS/MS analysis of protease-digested peptides was conducted on a QSTAR XL LC-MS/MS system (Applied Biosystems/MDS SCIEX) with HPLC system (NanoLC-2DTM, Eksigent Technologies) and ESI probe. Peptides were separated on a C18 reversed column (3 µm, 300Å, 0.1 x 70mm) and eluted at a flow rate of 0.2 µL/min using a gradient of 0-50% of solvent B (89%) acetonitrile/10% H₂O/0.02% TFA/1% formic acid) and solvent A (2% acetonitrile/97% H₂O/0.02% TFA/1% formic acid) in 55 min. The column eluent was transferred to the electrospray source and sequencing was carried out by fragmentation of the precursor ion corresponding to the peptide containing the residue at position 38 of mutant mDHFR.

Fluorescent Protein Isolation. Fluorescent proteins were purified by Ni-NTA affinity chromatography according to the manufacturer's protocol (Qiagen) under native

conditions. Purified proteins were desalted over a Sephadex PD-10 column (Amersham Pharmacia, Piscataway, NJ) using 20 mM Tris-HCl, 1 mM EDTA, pH 8.0. Protein samples other than CFP6 were further purified by ammonium sulfate precipitation.³⁸ Dry ammonium sulfate was added to the bright fluorescent solution to 40% saturation (about 1.6 M). Then solutions were incubated on ice for 25 min and the precipitated proteins were removed by centrifugation at 14,000 rpm for 15 min at 4 °C. Dry ammonium sulfate was added to the fluorescent supernatants to 70% saturation (about 2.8 M). The solutions were incubated on ice for 30 min and the fluorescent proteins were collected by centrifugation at 14,000 rpm for 30 min at 4 °C. The fluorescent proteins were resuspended in 20 mM Tris-HCl, 1 mM EDTA, pH 8.0 and desalted over polyacrylamide P-6 columns (Bio-rad) using 20 mM Tris-HCl, 1 mM EDTA, pH 8.0 buffer. Protein concentrations were determined by the BCA protein assay kit (Pierce Biotechnology, Rockford, IL) using bovine serum albumin as a reference.

Characterization of Spectral Properties. Absorption spectra of fluorescent proteins were measured on a Cary 50 UV-visible spectrophotometer (VARIAN, Palo Alto, CA). Protein samples were diluted to four different concentrations ($0.1 < OD_{\lambda max} < 0.5$). Extinction coefficients were calculated by applying Beer's law to the absorption spectra of diluted protein samples. A molar mass of 30,287 Da was used for all fluorescent proteins. Fluorescence spectra were measured on a PTI QuantaMaster fluorescence spectrofluorometer (Photon Technology International, Birmingham, NJ) at a scan rate of 1 nm/s. For the quantum yield calculation, protein samples in assay buffer and fluorescenin (Invitrogen) in borate buffer (pH 9.1) were adjusted to similar OD (OD_{$\lambda ex} < 0.1$). The</sub>

fluorescence emission spectra were recorded with excitation at the maxima in the corresponding excitation spectra. The emission spectra were corrected by the correction functions supplied with the instrument. The quantum yields of fluorescent proteins were calculated by comparing the integrated intensities of the corrected emission spectra with that of fluorescein of known quantum yield (0.91).³⁹

Results and Discussion

Rational Design of yPheRS and Choice of Trp Analogs. In order to increase the number of genetically encoded Trp analogs, we engineered the substrate specificity of yPheRS. Studies of the crystal structure of PheRS (tPheRS) derived from Thermus thermophilus revealed that Val 261 in the amino acid binding pocket is critical for distinguishing Phe from other non-aromatic amino acids.⁴⁰ Sequence alignments showed that Val 261 in tPheRS corresponds to Thr 415 in yPheRS. Introduction of a single mutation (Thr to Gly) at position 415 in the binding pocket of yPheRS (yPheRS (T415G)) created extra room to accommodate various Phe analogs.^{25,31} There was an interesting finding that yPheRS (T415G) activated Trp and 2-naphthylalanine as well as substituted Phe analogs. Considering the spectrum of nonnatural amino acids recognized by yPheRS (T415G), we hypothesized that yPheRS (T415G) may activate Trp analogs. In order to achieve site-specific incorporation of Trp analogs, we needed to choose Trp analogs that can be recognized by yPheRS (T415G) but by the native E. coli translation system. We have chosen four Trp analogs (6ClW, 6BrW, 5BrW, and BT) that can not be utilized by the endogenous E. coli translational system. Based on previous data, 10,11 we proposed a model for the adaptability of Trp analogs to the binding pocket of E. coli

TrpRS (eTrpRS) (Figure 2). Recognition of Trp analogs by eTrpRS depends on the size and position of substituents on the indole ring. Methyl (ca. 16.8 Å³),⁴¹ amino (ca. 12.3 Å³), and fluorine groups (ca. 3.2 Å³) were the largest functional groups tolerated at the C4, C5 and C6 positions of the indole ring, respectively.¹⁰ Therefore, introduction of bulkier groups at each position would result in translationally inactive Trp analogs. Based on our hypothetical model, we chose three translationally inactive Trp analogs, **3**, **4**, and **5** containing chlorine (ca. 14.3 Å³) or bromine (ca. 20 Å³) at the C6 position, and bromine at the C5 position of the indole ring. Additionally, we chose **6**, known as an translationally inactive Trp analog.⁴² Replacement of the nitrogen in the indole ring of Trp with other hetero-atoms, such as selenium, sulfur, or oxygen, results in translationally inactive amino acids.^{15,42} As expected, none of the four Trp analogs (**3-6**) were incorporated into proteins, even in Trp auxotrophic *E. coli* strain under Trp depleted conditions (data not shown).

Activation of Trp Analogs by yPheRS (T415G). The relative rates of activation of Phe, Trp and Trp analogs (1-6) by yPheRS (T415G) were examined by ATP-PP_i exchange assays. The kinetic parameters are shown in Table 1. Derivatives of Trp (3-6) revealed different activities with respect to substitution at different positions of the indole ring. Substitution at position 6 (3 and 4) was highly favorable; 3 displayed the highest activity among all the analogs we tested. The increase in side-chain volume upon replacement of 2 with 3 (ca. 39 Å³) matched very well the cavity (ca. 39 Å³) generated by the T415G mutation in the binding pocket of yPheRS. 4 exhibited a k_{cat}/K_m value comparable to that of Trp. Notably, 4 showed 7-fold larger K_m value than that of 3, indicating less favorable

binding of **4** to the binding pocket of yPheRS. This result is consistent with the fact that the calculated increase in side-chain volume (45 Å³) upon replacement of **2** with **4** is larger than the cavity generated by the T415G mutation. Substitution at position 5 of the indole ring was less favorable than that at position 6. The activation rate of **5** by yPheRS (T415G) was one half of that of **4**. Replacement of an imine in the indole ring with a sulfur atom reduced binding to yPheRS (T415G). The K_m of **6** was 3-fold greater than that of **1**. However, the activation rate of **6** by yPheRS (T415G) was 6.4-fold faster than that of **2**. The spectrum of activity of the yPheRS (T415G) supported our proposal that the yPheRS (T415G) may recognize Trp analogs.

Site-Specific Incorporation of Trp Analogs into mDHFR in Vivo. To investigate the utility of yPheRS (T415G) to incorporate Trp analogs into proteins in vivo, mDHFR was chosen as a test protein. An amber stop codon was placed at the 38^{th} position of mDHFR to generate mDHFR_38Am. The expression of full length mDHFR_38Am with various Trp analogs (**3-6**) was examined by SDS-PAGE analysis (Figure 3). All of the Trp analogs supported expression of full length mDHFR_38Am. The band intensities of mDHFR_38Am variants were in the order of **4** > **3** > **5** > **6**. Although expression of full length mDHFR_38Am strongly indicated that the amber stop codon was supported of Trp analogs were incorporated at the amber site. In order to verify incorporation of Trp analogs at the amber site, the occupancy of the site in mDHFR_38Am by various amino acids was determined by LC-MS/MS analysis. Trypsin digestion of purified mDHFR_38Am variants generated diverse peptide fragments. Among them, a peptide

containing the amber site (residues 26-39, NGDLPWPPLRNEAmK) was designated as peptide 38. Peptide 38 variants containing Phe, Trp or Trp analogs were separated on a C₁₈ reversed phase column and relative yields were determined by comparing the integrated areas of the corresponding signals in the chromatogram. The results are summarized in Table 2. Furthermore, the identity of each peptide was confirmed by tandem-mass spectrometry. As a representative, the identity of peptide 38 containing 6BrW at the amber site was determined from the spectrum shown in Figure 4. We observed that 3, 4, and 6 occupied at least 98% of the amber sites in full length mDHFR 38Am (Table 2). However, 5 occupied only 2% of the amber site, though it was efficiently activated by vPheRS (T415G). Instead, 2 occupied 97% of the amber site. The discrepancy between in vitro activation and in vivo incorporation for 5 could be rationalized by the model proposed by Sisido and colleagues for the adaptability of nonnatural amino acids to the E. coli ribosome.⁴³ The model suggests that the E. coli ribosome has the ability to define the allowed and excluded regions of the aromatic groups of aryl-L-alanines to be bound for further translation. The addition of bromine increases the volume of the indole ring by 20 Å³. We reason that the addition of bromine at the C5 position of the indole ring may result in exclusion of 5 by the ribosome. Another interesting finding from the experiment for analog 5 is that at similarly low concentrations of Phe and Trp in the culture medium, Phe was incorporated into the amber site instead of Trp. According to our in vitro activation assays, Trp was activated 18-fold faster than Phe by yPheRS (T415G). This discrepancy between the results of in vitro activation assays and in vivo translation might be explained by elevated Trp degradation in the presence of Trp analogs. Expression of tryptophanase, a major enzyme

involved in Trp degradation, is regulated by tryptophan-induced transcription antitermination in *E. coli*.^{44,45} Even 1-methyltryptophan, a Trp analog, acted as an effective inducer in vitro.⁴⁵ Therefore, we reasoned that supplementation of **5** into the culture medium accelerated Trp degradation via formation of tryptophanase and reduced the intracellular concentration of Trp levels lower than that of Phe. Since the ytRNA^{Phe}_{CUA_UG} charged with **5** was rejected in ribosome, the ytRNA^{Phe}_{CUA_UG} charged with **5** was rejected in ribosome, the ytRNA^{Phe}_{CUA_UG} charged with **2** was dominantly utilized for translation.

Expression yields of mDHFR_38Am with various Trp analogs were determined by measuring the concentration of purified mDHFR_38Am. The results are summarized in Table 2. One interesting finding was that 4 provided 25% higher expression level of mDHFR_38Am as compared to 3 despite its 3.5-fold lower k_{cat}/K_m value. This discrepancy can be rationalized by higher k_{cat} of 4 than that of 3. The concentration (3 mM) of Trp analogs supplemented into culture medium was at least 900-fold higher than K_m values of both 3 and 4. In such a case, the k_{cat} value became more important than k_{cat}/K_m in determining relative activation rates. Therefore, 4 with 60% higher k_{cat} value than that of 3 could be more efficiently used for protein translation.

Spectral Analysis of Fluorescent Proteins. The spectral properties of CFP6 and CFP6_66Am proteins containing Trp analogs are summarized in Table 3. Absorption maxima, emission maxima, and extinction coefficient of CFP6 were consistent with literature values.⁴⁶ Compared to CFP6, CFP6_6CIW and CFP6_6BrW showed blue-shifts of 14 nm in the fluorescence emission maxima (Figure 6). Similar trends were found in previous work in which Tyr at position 66 of GFP was replaced by *p*-

bromophenylalanine.³⁶ In addition, solution studies indicated that the absorption and fluorescence spectra of fluorinated tryptophans were blue-shifted;¹⁶ the electronwithdrawing property of halogen functional groups can account for the blue-shift in the fluorescence emission peak. Compared with CFP6, the quantum yields of the CFP6 6CIW and CFP6 6BrW decreased 2.5 to 3-fold (Table 3), probably due to collisional perturbation of spin-orbital coupling in the π -electron orbitals of the fluorophore by the halogen atom.⁴⁷ CFP6_BT showed blue-shifts of 16 and 34 nm in fluorescence emission and absorption maxima, respectively. Similar blue-shift in absorption maxima in CFP variants was observed by replacing a tryptophan with sulfur containing tryptophan surrogates, such as β -(thieno[2,3-b]pyrrolyl)-L-alanine ([2.3]Tpa) and β -(thieno[3,2-b]pyrrolyl)-L-alanine ([3.2]Tpa).¹⁶ Extinction coefficient of CFP6 BT (8,100 M⁻¹cm⁻¹) was similar to those of CFP variants containing either [2,3]Tpa (6840 M⁻¹ 1 cm⁻¹) or [3,2]Tpa (6350 M⁻¹ cm⁻¹). Introduction of either [2,3]Tpa or [3,2]Tpa in the chromophore of CFP resulted in the disappearance of fluorescence;¹⁶ however, CFP6 BT exhibited substantial fluorescence, with quantum yield only 3-fold smaller than that of CFP6. CFP6 BT is the only fluorescent protein containing a sulfur atom in its chromophore. CFP6 BT has several advantages as a fluorescence resonance electron transfer (FRET) partner. It can be efficiently excited by a common violet diode laser (405-415 nm), which can simplify cell screening on the basis of FRET signals. In addition, the Stokes' shift of CFP6 BT (56 nm) is larger than that of CFP6 (37 nm). This large Stokes' shift is advantageous for fluorescence measurements, since it allows emission photons to be detected against a low background and isolated from excitation photons. CFP and yellow fluorescent protein (YFP) have been commonly used as a FRET

pair. Because the fluorescence emission spectrum of CFP_BT (Figure 6 b) overlaps the excitation spectrum of GFP, it can be used as a FRET partner with GFP.

Conclusions

In this paper, we have shown that properly engineered *E. coli* strains can allow high-fidelity incorporation of Trp analogs into recombinant proteins in response to an amber stop codon. Rational modification of the binding pocket in yPheRS enabled activation of Trp analogs (**3-6**). When the engineered yPheRS and the yeast amber suppressor tRNA were introduced into a Phe/Trp auxotrophic *E. coli* expression host, at least 98% of the amber sites in recombinant mDHFR were occupied by Trp analogs (**3**, **4** and **6**). In a subsequent experiment, three Trp analogs (**3**, **4**, and **6**) were introduced at the Trp66 position of CFP. CFP variants containing Trp analogs (**4**, **5**, and **6**) in the chromophore showed blue-shifts in fluorescence emission and absorption maxima. In particular, CFP6BT provided unusually large Stokes' shift (56 nm). Introduction of aryl bromides into recombinant proteins should allow site-specific modification via bio-orthogonal palladium-catalyzed coupling reactions. An expanded set of genetically encoded Trp analogs should enable design of new proteins with novel properties.

Acknowledgments

We thank I. C. Tanrikulu, R. Connor, and T. H. Yoo for discussion and valuable comments. We are grateful to Dr. J. Zhou for help with LC-MS analysis. This work was supported by National Institutes of Health grant.

References

- 1. Fulton, K. F.; Jackson, S. E.; Buckle, A. M. Biochemistry 2003, 42, 2364-2372.
- Uchiyama, T.; Katouno, F.; Nikaidou, N.; Nonaka, T.; Sugiyama, J.; Watanabe, T. J. Biol. Chem. 2001, 276, 41343-41349.
- Bray, M. R.; Johnson, P. E.; Gilkes, N. R.; McIntosh, L. P.; Kilburn, D. G.; Warren, R. A. J. *Protein Sci.* 1996, *5*, 2311-2318.
- Marie, G.; Serani, L.; Laprevote, O.; Cahuzac, B.; Guittet, E.; Felenbok, B. *Protein Sci.* 2001, *10*, 99-107.
- Eason, D. E.; Shepherd, A. T.; Blanck, G. Biochim. Biophys. Acta Gene Struct. Express. 1999, 1446, 140-144.
- Dayoff, M. O., *Atlas of protein sequence and structure* National Biomedical Research Foundation: Washington DC, 1972.
- 7. Wong, C. Y.; Eftink, M. R. Protein Sci. 1997, 6, 689-697.
- 8. Brawerman, G.; Ycas, M. Arch. Biochem. Biophys. 1957, 68, 112-117.
- 9. Pardee, A. B.; Shore, V. G.; Prestidge, L. S. Biochim. Biophys. Acta 1956, 21, 406-407.
- 10. Budisa, N. Angew. Chem. Int. Ed. 2004, 43, 6426-6463.
- 11. Budisa, N.; Pal, P. P. Biol. Chem. 2004, 385, 893-904.
- 12. Link, A. J.; Mock, M. L.; Tirrell, D. A. Curr. Opin. Biotechnol. 2003, 14, 603-609.
- 13. Pratt, E. A.; Ho, C. Fed. Proc. 1974, 33, 1463-1463.
- Seifert, M. H.; Ksiazek, D.; Azim, M. K.; Smialowski, P.; Budisa, N.; Holak, T. A. J. Am. Chem. Soc. 2002, 124, 7932-7942.
- Bae, J. H.; Alefelder, S.; Kaiser, J. T.; Friedrich, R.; Moroder, L.; Huber, R.; Budisa, N. J. Mol. Biol. 2001, 309, 925-936.

- Budisa, N.; Pal, P. P.; Alefelder, S.; Birle, P.; Krywcun, T.; Rubini, M.; Wenger, W.;
 Bae, J. H.; Steiner, T. *Biol. Chem.* 2004, *385*, 191-202.
- Budisa, N.; Rubini, M.; Bae, J. H.; Weyher, E.; Wenger, W.; Golbik, R.; Huber, R.;
 Moroder, L. Angew. Chem. Int. Ed. 2002, 41, 4066-4069.
- Bae, J. H.; Rubini, M.; Jung, G.; Wiegand, G.; Seifert, M. H. J.; Azim, M. K.; Kim, J. S.; Zumbusch, A.; Holak, T. A.; Moroder, L.; Huber, R.; Budisa, N. *J. Mol. Biol.* 2003, 328, 1071-1081.
- 19. Wang, L.; Schultz, P. G. Chem. Commun. 2002, 1-11.
- 20. Kiick, K. L.; Weberskirch, R.; Tirrell, D. A. FEBS Lett. 2001, 505, 465-465.
- 21. Kirshenbaum, K.; Carrico, I. S.; Tirrell, D. A. ChemBioChem 2002, 3, 235-237.
- 22. Tang, Y.; Tirrell, D. A. Biochemistry 2002, 41, 10635-10645.
- Doring, V.; Mootz, H. D.; Nangle, L. A.; Hendrickson, T. L.; de Crecy-Lagard, V.; Schimmel, P.; Marliere, P. *Science* 2001, 292, 501-504.
- 24. Datta, D.; Wang, P.; Carrico, I. S.; Mayo, S. L.; Tirrell, D. A. J. Am. Chem. Soc. 2002, 124, 5652-5653.
- 25. Kwon, I.; Wang, P.; Tirrell, D. A. J. Am. Chem. Soc. 2006, 128, 11778-11783.
- Santoro, S. W.; Wang, L.; Herberich, B.; King, D. S.; Schultz, P. G. Nat. Biotechnol.
 2002, 20, 1044-1048.
- 27. Link, A. J.; Vink, M. K. S.; Agard, N. J.; Prescher, J. A.; Bertozzi, C. R.; Tirrell, D. A. Proc. Natl. Acad. Sci. U. S. A. 2006, 103, 10180-10185.
- 28. Zhang, Z. W.; Alfonta, L.; Tian, F.; Bursulaya, B.; Uryu, S.; King, D. S.; Schultz, P. G. Proc. Natl. Acad. Sci. U. S. A. 2004, 101, 8882-8887.
- 29. Furter, R. Protein Sci. 1998, 7, 419-426.

- 30. Wang, L.; Brock, A.; Schultz, P. G. J. Am. Chem. Soc. 2002, 124, 1836-1837.
- 31. Wang, P., Ph.D. thesis, California Institute of Technology, Pasadena, 2003.
- 32. The mutant yeast amber suppressor was used to reduce lysine misincorporation at the amber sites, because it showed minimal cross-reactivity with *E. coli* lysyl-tRNA synthetase.
- Kodama, K.; Fukuzawa, S.; Nakayama, H.; Kigawa, T.; Sakamoto, K.; Yabuki, T.; Matsuda, N.; Shirouzu, M.; Takio, K.; Tachibana, K.; Yokoyama, S. *ChemBioChem* 2006, 7, 134-139.
- Ojida, A.; Tsutsumi, H.; Kasagi, N.; Hamachi, I. *Tetrahedron Lett.* 2005, 46, 3301-3305.
- 35. Xie, J. M.; Wang, L.; Wu, N.; Brock, A.; Spraggon, G.; Schultz, P. G. Nat. Biotechnol.
 2004, 22, 1297-1301.
- 36. Wang, L.; Xie, J. M.; Deniz, A. A.; Schultz, P. G. J. Org. Chem. 2003, 68, 174-176.
- 37. Gill, S. C.; Vonhippel, P. H. Anal. Biochem. 1989, 182, 319-326.
- Yakhnin, A. V.; Vinokurov, L. M.; Surin, A. K.; Alakhov, Y. B. Protein Expr. Purif. 1998, 14, 382-386.
- 39. Velapoldi, R. A.; Tonnesen, H. H. J. Fluoresc. 2004, 14, 465-472.
- 40. Fishman, R.; Ankilova, V.; Moor, N.; Safro, M. Acta Crystallogr. D. Biol. Crystallogr.
 2001, 57, 1534-44.
- 41. Volumes were estimated as solvent-excluded volumes (1.4 Å probe) by using Chem3D Pro 5.0 (Cambridge, MA)
- 42. Budisa, N.; Alefelder, S.; Bae, J. H.; Golbik, R.; Minks, C.; Huber, R.; Moroder, L. *Protein Sci.* 2001, 10, 1281-1292.

- 43. Hohsaka, T.; Kajihara, D.; Ashizuka, Y.; Murakami, H.; Sisido, M. J. Am. Chem. Soc. 1999, 121, 34-40.
- 44. Gong, F.; Ito, K.; Nakamura, Y.; Yanofsky, C. Proc. Natl. Acad. Sci. U. S. A. 2001, 98, 8997-9001.
- 45. Gong, F.; Yanofsky, C. J. Biol. Chem. 2001, 276, 1974-1983.
- 46. Tsien, R. Y. Annu. Rev. Biochem. 1998, 67, 509-544.
- 47. Kasha, M. J. Chem. Phys. 1952, 20, 71-74.



Figure 1: Amino acids involved in this study. (1) tryptophan, (2) phenylalanine, (3) 6chlorotryptophan, (4) 6-bromotryptophan, (5) 5-bromotryptophan, (6) benzothienylalanine.



Figure 2: A hypothetical model for the adaptability of Trp analogs to the binding pocket of eTrpRS. Position and size of substituents on the indole ring that can be tolerated in the binding pocket of eTrpRS are represented as circles. Indole rings containing larger substituents can not fit into the binding pocket of eTrpRS.



Figure 3: SDS-PAGE analysis of mDHFR_38Am expression. Cells were equipped with yPheRS (T415G) and ytRNA^{Phe}_{CUA_UG}. Cultures were supplemented with 18 amino acids (at 25 mg/L), one of Trp analogs (at 3 mM) and the indicated concentrations of Phe and Trp. Lane 1: molecular weight standards, lane 2: before induction, lane 3: 6ClW, 0.03 mM Phe and 0.01 mM Trp, Lane 4: 6BrW, 0.015 mM Phe and 0.05 mM Trp, lane 5: 5BrW, 0.03 mM Phe and 0.01 mM Trp, lane 6: BT, 0.01 mM Phe and 0.0025 mM Trp.



Figure 4: Tandem mass spectrum of the peptide (NGDLPWPPLRNEZK). The partial sequence DLPWPPLRNEZ containing 6BrW (\mathbb{Z}) can be confirmed by assigning masses of series of fragmented ions (*b* and *y*).



Figure 5: Crystal structure of enhanced cyan fluorescent protein (PDB code 10XD)¹⁸ (a). The chromophore is shown in black. Chromophores of CFP6 variants containing 1, 3, 4, and 6 (b).



Figure 6: Absorption (**a**) and fluorescence emission spectra (**b**) for CFP6 variants containing different Trp analogs at the Trp66 position.

Amino Acid	$K_{m}\left(\mu M ight)$	k_{cat} (s ⁻¹)	$k_{cat}/K_m (M^{\text{-1}}s^{\text{-1}})$	$k_{cat}/K_{m}\left(rel\right)^{a}$
Phe (2)	55±14	0.20±0.11	3,500±1,100	1
Trp (1)	2.8±1.6	0.15±0.003	63,200±34,600	18
6ClW (3)	0.51±0.21	0.11±0.04	224,100±15,300	64
6BrW (4)	3.4±1.3	0.18±0.11	61,800±53,900	18
5BrW (5)	0.62±0.02	0.021±0.002	33,900±3,600	9.7
BT (6)	8.1±3.7	0.15±0.06	22,500±18,000	6.4

Table 1: Kinetic parameters for ATP-PPi exchange by the yPheRS (T415G)

^a relative to k_{cat}/K_m for Phe by yPheRS (T415G)

Trp analogs	Occupancy of amber sites (%)				Yield
	Phe	Trp	Lys	Naa	(mg/L)*
Trp (1)	0.4 ± 0.5	ND^{b}	1.2 ± 1.3	98.4 ± 1.7	5.4 ± 0.3
6ClW (3)	0.4 ± 0.7	ND	ND	99.6 ± 0.7	6.8 ± 1.3
6BrW (4)	97.0 ± 1.4	0.3 ± 0.6	0.7 ± 0.5	1.8 ± 0.7	3.6 ± 0.6
BT (6)	0.6 ± 0.6	0.7 ± 0.6	0.9 ± 1.5	97.9 ± 2.4	2.8 ± 0.6

 Table 2: Occupancy of amber sites and expression yields

^a Volumetric yields are given as mg of purified mDHFR_38Am per liter of culture. ^b Not detected in LC-MS analysis.

amino acid at position 66	absorption maximum (nm)	extinction coefficient $(M^{-1} \text{ cm}^{-1})$	emission maximum (nm)	quantum yield
Trp (1)	449	25900 ± 1000	488	0.14 ± 0.008
6ClW (3)	430	19800 ± 6400	474	0.050 ± 0.001
6BrW (4)	430	20600 ± 3800	474	0.058 ± 0.001
BT (6)	415	8100 ± 1100	472	0.048 ± 0.008

Table 3: Spectral properties of CFP6 variants

Chapter 5

Breaking the Degeneracy of the Genetic Code

The text in this chapter is reprinted with permission from Kwon, I.; Kirshenbaum, K.; Tirrell, D. A. *J. Am. Chem. Soc.* **2003**, *125*, 7512-7513, Copyright 2003 Am. Chem. Soc.

Abstract

A mutant yeast phenylalanine transfer RNA (ytRNA^{Phe}_{AAA}) containing a modified (AAA) anticodon was generated to explore the feasibility of breaking the degeneracy of the genetic code in *Escherichia coli*. By using an *E.coli* strain co-transformed with ytRNA^{Phe}_{AAA} and a mutant yeast phenylalanyl-tRNA synthetase, we demonstrate efficient codon-biased replacement of phenylalanine (Phe) by L-3-(2-naphthyl)alanine (2Nal), a non-proteinogenic analog. Site-specific incorporation of 2Nal in response to UUU codons was confirmed by mass spectrometric analysis of recombinant murine dihydrofolate reductase. These results illustrate a general method for increasing the number of distinct, genetically-encoded amino acids available for protein engineering and for exploration of the chemistry and physics of protein-like macromolecules.

Introduction

Organisms use a canonical set of 20 amino acids to generate the proteins that sustain the life of the cell. In recent years, several laboratories have pursued an expansion in the number of genetically-encoded amino acids, by using either a nonsense suppressor or a frameshift suppressor tRNA to incorporate non-canonical amino acids into proteins in response to amber or four-base codons, respectively.¹⁻¹⁰ Such methods have worked well for single-site insertion of novel amino acids; however, their utility in multi-site incorporation is limited by modest (20-60%) suppression efficiencies.^{1, 5, 11}

Efficient multi-site incorporation has been accomplished by replacement of natural amino acids in auxotrophic *E. coli* strains,¹²⁻¹⁵ and by using aminoacyl-tRNA synthetases with relaxed substrate specificity or attenuated editing activity.^{14, 16} Although this method

provides efficient incorporation of analogs at multiple sites, it suffers from the limitation that the novel amino acid must "share" codons with one of the natural amino acids. We present here a potential solution to this coding problem.

Material and Methods

Materials. All chemical reagents were of analytical grade, obtained from commercial suppliers, and used without further purification unless otherwise noted. L-3-(2-naphthyl)alanine and other natural amino acids were obtained from Sigma (St. Louis, MO).

Strains and Plasmids. An *E. coli* strain XL1-Blue (Stratagene) was used for plasmid propagation and isolation. A Phe auxotrophic strain K10-F6 Δ (K10, Hfr(Cavalli) *pheS13rel-1 tonA22* thi *T2*^R *pheA18*)⁴ was a gift from Rolf Furter (University of Massachusetts). Plasmids carrying modified yeast tRNA^{Phe} variants were derived from pRO117⁴ in which yeast tRNA^{Phe} expression cassette was inserted at the *Sna*I site of the pREP4 (Qiagen). Plasmids for expression of mDHFR and overexpression of mutant yeast phenylalanyl-tRNA synthetases (yPheRS) were derived from pRO148⁴ in which yPheRS genes under constitutive *tac* promoter control was inserted at the *PvuII* site of pQE16 (Qiagen).

Construction of Plasmid Carrying Mutant Yeast tRNA^{Phe}. pREP4_ytRNA^{Phe}_AAA containing a mutant yeast tRNA^{Phe} with a modified anticodon (AAA) (ytRNA^{Phe}_{AAA}) was generated by PCR mutagenesis kit (Stratagene) using pRO117⁴ as a template.
Construction of Plasmid Carrying mDHFR and Mutant Yeast Phenylalanyl-tRNA Synthetase. An intact mDHFR expression cassette was obtained by endonuclease restriction of pQE16 at the *Aat*II and *Nhe*I sites. This expression cassette was inserted between the *Aat*II and *Nhe*I sites of pRO148 to generate pQE16_mDHFR_yPheRS. A mutation of threonine to glycine at the 415th position of α -subunit of yPheRS was performed by PCR mutagenesis kit (Stratagene) using pQE16_mDHFR_yPheRS as a template to generate pQE16_mDHFR_yPheRS (T415G).

Protein Expression and Purification. Cultures of K10-F6Δ outfitted with pREP4_ytRNA_AAA and pQE16_mDHFR_yPheRS (T415G) were grown in M9 minimal medium supplemented with 0.4% (w/v) of glucose, 0.1 mM of CaCl₂, 1.0 mM of MgSO₄, 35 µg/mL of thiamine, 20 amino acids (25 mg/L), 100 µg/mL of ampicillin, and 35 µg/mL of kanamycin. When an optical density of the culture reached 0.8 to 1.0 at 600nm (OD₆₀₀), the cultures were centrifuged for 7 min (6,000 rpm) at 4 °C. The cell pellets were washed twice with 0.9% (w/v) of NaCl solution. The cells were resuspended in M9 minimal medium supplemented with 18 amino acids (25 mg/L) and indicated amounts of 2Nal and Phe. After 10 min incubation, 1 mM of isopropyl-1-thio-β-D-galactopyranoside (IPTG) was added to induce mDHFR protein expression. The OD₆₀₀ of the culture was measured 4 hrs after induction. Then the cells were harvested by centrifugation for 10 min (6,000 rpm) at 4°C and stored at -70°C. After thawing, mDHFR proteins were purified under denaturing conditions according to manufacturer's

protocol (Qiagen). Protein expression was evaluated by SDS-PAGE with coomassie blue staining. Loading of the gel was normalized for cell densities as determined by OD₆₀₀.

Amino Acid Analysis. The purified mDHFR solutions were concentrated 10-fold by ultrafiltration (Millipore) followed by a buffer exchange against 0.1% trifluoroacetic acid (TFA) solution. Samples were submitted to the Molecular Structure Facility at the University of California, Davis, for amino acid analysis on a Beckman 6300 instrument (Fullerton, CA).

Matrix-Assisted Laser Desorption Ionization Mass Spectrometry (MALDI–MS) Analysis. The purified mDHFR solutions were desalted by ZipTip_{C18} (Millipore) and eluted with 3 μ L of 50% CH₃CN/0.1% TFA. 1 μ L was used for matrix-assisted laser desorption ionization mass spectrometry (MALDI-MS) analysis with sinapinic acid (30% (v/v) of acetonitrile and 70% (v/v) of 0.1% TFA solution) as the matrix. The analysis was performed on a PerSeptive Biosystems (Framingham, Massachusetts) Voyager DE PRO MALDI-TOF mass spectrometer operating in linear and positive ion modes. For MALDI-MASS analysis of tryptic digests of purified mDHFR, 10 μ L of the concentrated protein solution was added to 90 μ L of 75 mM ammonium bicarbonate solution. 2 μ L of 0.2 g/L of modified trypsin (Promega) was added, and the solution was incubated at 37°C for 2.0 hrs. 12 μ l of 5% TFA solution was added to quench the reaction. Chromatography on Ziptip_{C18} columns (Millipore) provided purified peptide samples (1 μ L), which were added to 2,5-dihydroxybenzoic acid (DHB) MALDI matrix solution for MALDI-MASS analysis.

Results and Discussion

The genetic code is degenerate, in that the protein biosynthetic machinery utilizes 61 mRNA sense codons to direct the templated polymerization of a set of 20 amino acid monomers.¹⁷ Just two amino acids, methionine and tryptophan, are encoded by unique mRNA triplets. Re-assignment of degenerate sense codons therefore offers the prospect of a substantially expanded genetic code and a correspondingly enriched set of building blocks for natural and artificial proteins.

As a test case for establishing the feasibility of breaking the degeneracy of the code, we chose the biosynthetic machinery responsible for incorporation of phenylalanine (Phe) into the proteins of *E. coli*. Phe is encoded by two codons, UUC and UUU. Both codons are read by a single ytRNA, which is equipped with the anticodon sequence GAA. The UUC codon is therefore recognized through standard Watson-Crick basepairing between codon and anticodon; UUU is read through a G-U wobble base pair at the first position of the anticodon.¹⁸ Thermal denaturation of RNA duplexes has yielded estimates of the Gibbs free energies of melting of G-U, G-C, A-U, and A-C base pairs as 4.1, 6.5, 6.3, and 2.6 kcal/mol, respectively, at 37°C.¹⁹ Thus the wobble base pair, G-U, is less stable than the Watson-Crick base pair, A-U. On this basis, we proposed that a mutant ytRNA^{Phe} outfitted with the AAA anticodon (ytRNA^{Phe}_{AAA}) might be engineered to read UUU codons faster than wild-type ytRNA^{Phe}. If ytRNA^{Phe}_{AAA} can then be charged selectively with an amino acid analog, one should be able to accomplish codonbiased incorporation of the analog at multiple sites in recombinant proteins. With respect to reading of UUC, an unmodified A in the first position of the anticodon is known to read codons ending with C, as well as U, in the absence of ytRNAs containing G in the

first anticodon position.²⁰⁻²² However, the binding of *E. coli* tRNA^{Phe}_{GAA} should dominate the binding of ytRNA^{Phe}_{AAA} owing to differences in the stability of A-C and G-C base pairs (see above).

The approach used here is a modification of the method introduced by Furter for site-specific insertion of amino acid analogs *in vivo*.⁴ The method involves introduction into *E. coli* of a heterologous aminoacyl-tRNA synthetase and its cognate tRNA. If cross-charging between the heterologous pair and the translational apparatus of the host is slow or absent, and if the analog is charged only by the heterologous synthetase, insertion of the analog can be restricted (or at least biased) to sites characterized by productive base-pairing between the heterologous tRNA and the messenger RNA of interest.

In order to test these ideas, we prepared a yeast tRNA^{Phe} (ytRNA^{Phe}_{AAA}) with an altered anticodon loop. The first base (G³⁴) of the ytRNA^{Phe} anticodon (GAA) was replaced with A to provide specific Watson-Crick base-pairing to the UUU codon. Furthermore, G³⁷ in the extended anticodon site was replaced with A in order to increase translational efficiency.^{10, 23} We believe that charging of yeast tRNA^{Phe}_{AAA} by *E. coli* PheRS can be ignored, because the aminoacylation rate of yeast tRNA^{Phe}_{AAA} by *E. coli* PheRS is known to be 0.1% of that of *E. coli* tRNA^{Phe}_{GAA}.²⁴

Since wild-type yeast PheRS does not activate amino acids significantly larger than phenylalanine, a mutant form of the synthetase with relaxed substrate specificity was prepared in order to accommodate L-3-(2-naphthyl)alanine (2Nal).¹⁰ On the basis of prior work from this laboratory,¹⁶ the mutant yeast PheRS (yPheRS (T415G)) was prepared by introduction of a Thr415Gly mutation in the α -subunit of the synthetase. The kinetics of activation of 2Nal and Phe by yPheRS (T415G) was analyzed *in vitro* via the

pyrophosphate exchange assay. The specificity constant (k_{cat}/K_M) for activation of 2Nal by yPheRS (T415G) was found to be 1.55×10^{-3} (S⁻¹ μ M⁻¹), 8-fold larger than that for Phe.¹⁰ Therefore, when the ratio of 2Nal to Phe in the culture medium is high, ytRNA^{Phe}_{AAA} should be charged predominantly with 2Nal. Recently, we have shown that 2Nal can be incorporated with better than 95% efficiency via amber suppression in an *E. coli* strain co-transformed with yPheRS (T415G) and ytRNA^{Phe}_{CUA}.

Murine dihydrofolate reductase (mDHFR), which contains nine Phe residues, was chosen as the test protein. The expression plasmid pQE16 encodes mDHFR under control of a bacteriophage T5 promoter; the protein is outfitted with a C-terminal hexahistidine tag to facilitate purification via immobilized metal affinity chromatography. In this construct, four of the Phe residues of mDHFR are encoded by UUC codons, five by UUU. A full-length copy of the yPheRS (T415G) gene, under control of a constitutive tac promoter, was inserted into pQE16. The gene encoding ytRNA^{Phe}_{AAA} was inserted into the repressor plasmid pREP4 (Qiagen) under control of the constitutive promoter lpp. E. coli transformants harboring these two plasmids were incubated in Phe-depleted minimal medium supplemented with 3 mM 2Nal, and then treated with 1 mM IPTG to induce expression of mDHFR. Although the *E. coli* strain (K10-F6 Δ) used in this study is a Phe auxotroph,⁴ a detectable level of mDHFR was expressed even under conditions of nominal depletion of Phe (Figure 2), probably due to release of Phe through turnover of cellular proteins. In negative control experiments, mDHFR was expressed in the absence of either ytRNA^{Phe}_{AAA} or yPheRS (T415G). mDFHR expression levels in these experiments were similar, indicating that neither ytRNA^{Phe}_{AAA} nor yPheRS (T415G) significantly reduces the protein synthesis rate (Figure 2). However, MALDI-MS spectra

and amino acid analyses of purified mDHFRs showed differences among samples prepared under these conditions (Table 1). The molar mass of mDHFR prepared in the absence of 2Nal, ytRNA^{Phe}_{AAA}, or yPheRS (T415G) was 23,287 daltons, precisely that calculated for His-tagged mDHFR. However, when ytRNA^{Phe}_{AAA} and yPheRS (T415G) were introduced into the expression strain and 2Nal was added to the culture medium, the observed mass of mDHFR was 23,537 daltons. Because each substitution of 2Nal for Phe leads to a mass increment of 50 daltons, this result is consistent with replacement of five Phe residues by 2Nal. No detectable mass shift was found in the absence of either ytRNA^{Phe}_{AAA} or yPheRS (T415G), confirming that the intact heterologous pair is required for incorporation of 2Nal. For mDHFR isolated from the strain harboring the heterologous pair, amino acid analysis indicated replacement of 4.4 of the 9 Phe residues by 2Nal (Table 1). Without ytRNA^{Phe}_{AAA} or yPheRS (T415G), no incorporation of 2Nal into mDHFR was detected by amino acid analysis.

Because neither MALDI nor amino acid analysis of intact mDHFR shows which Phe residues have been replaced by 2Nal, tryptic digests were analyzed to determine the occupancy of individual Phe sites. Tryptic digestion of mDHFR yields peptide fragments that are readily analyzed by MALDI mass spectrometry as shown in Figure 3. Peptide 1_{UUU} (residues 184-191, YKFEVYEK) contains a Phe residue encoded as UUU, whereas Peptide 2_{UUC} (residues 62-70, KTWFSIPEK) and Peptide 3_{UUC} (residues 26-39, NGDLPWPPLRNEFK) each contain a Phe residue encoded by UUC. In the absence of 2Nal, Peptide 1_{UUU} was detected with a monoisotopic mass of 1105.55 daltons, in accord with its theoretical mass (Figure 3A). However, when 2Nal was added, a strong signal at a mass of 1155.61 daltons was detected, and the 1105.55 signal was greatly reduced in intensity (Figure 3B). As described earlier, each substitution of 2Nal for Phe leads to a mass increase of 50.06 daltons; the observed shift in the experimental mass is thus consistent with replacement of Phe by 2Nal in response to the UUU codon. Liquid chromatography-tandem mass spectrometry of Peptide 1_{UUU} (Nal) was also carried out in order to determine more directly the origin of the observed increase in mass. The fragment ion masses could be unambiguously assigned as shown in Figure 4, confirming replacement of Phe by 2Nal. The ratio of MALDI signal intensities, though not rigorously related to relative peptide concentrations, suggests that 2Nal incorporation is dominant at the UUU codon.

Similar analyses were conducted for Peptide 2_{UUC} and Peptide 3_{UUC} . In the absence of added 2Nal, the observed masses of Peptides 2_{UUC} and 3_{UUC} are 1135.61 (Figure 3A) and 1682.89 daltons (Figure 3D), respectively. These observed masses match well the corresponding theoretical masses (1135.61 daltons for Peptide 2_{UUC} , and 1682.86 daltons for Peptide 3_{UUC}). Upon addition of 2Nal to the expression medium, the signals at these masses (Figure 3B and 3E) were not substantially reduced, and only very weak signals were observed at masses of 1185.60 and at 1733.03, which would be expected for peptides containing 2Nal in place of Phe. 2Nal incorporation thus appears to be rare at UUC codons under the conditions used here for protein expression. Other peptides containing encoded Phe show similar codon-biased selective incorporation of 2Nal and Phe.

There is at least a formal possibility that the observed codon-biased incorporation of 2Nal might be dependent on codon context rather than - or in addition to - codon identity. In order to test this possibility, mutant mDHFR genes were prepared by mutating the UUU codon in Peptide 1_{UUU} to UUC, and the UUC codon in Peptide 3_{UUC} to UUU. The resulting peptides were designated Peptide 1_{UUC} and Peptide 3_{UUU} , respectively. In Peptide 1_{UUC} , 2Nal incorporation was greatly reduced (Figure 3C), whereas for Peptide 3_{UUU} , 2Nal is readily detected (Figure 3F). 2Nal incorporation is unambiguously codon-biased to UUU.

Conclusions

The results described here show conclusively that a heterologous pair comprising a genetically engineered tRNA and cognate aminoacyl-tRNA synthetase can be used to break the degeneracy of the genetic code in *E. coli*. This method should provide a general strategy for multi-site incorporation of non-canonical amino acids, without the requirement that one of the natural amino acids be excluded. Introduction of more than one set of orthogonal pairs should allow several types of non-canonical amino acids to be incorporated in site-specific fashion. Ongoing experiments address the quantitative selectivity and the generality of the approach demonstrated here.

Acknowledgments

This work was supported by NIH GM62523 and by the NSF MRSEC program. We thank Dr. Mona Shahgholi, Dr. Gary M. Hathaway, and Dr. Jie Zhou for mass spectrometry studies. We thank Dr. Pin Wang and Dr. Soojin Son for helpful discussions.

References

1. Anderson, R. D.; Zhou, J.; Hecht, S. M. J. Am. Chem. Soc. 2002, 124, 9674-9675.

- 2. Bain, J. D.; Switzer, C.; Chamberlin, A. R.; Benner, S. A. Nature 1992, 356, 537-539.
- Chin, J. W.; Santoro, S. W.; Martin, A. B.; King, D. S.; Wang, L.; Schultz, P. G. J. Am. Chem. Soc. 2002, 124, 9026-9027.
- 4. Furter, R. Protein Sci. 1998, 7, 419-426.
- Hohsaka, T.; Ashizuka, Y.; Taira, H.; Murakami, H.; Sisido, M. *Biochemistry* 2001, 40, 11060-11064.
- 6. Magliery, T. J.; Anderson, J. C.; Schultz, P. G. J. Mol. Biol. 2001, 307, 755-769.
- Noren, C. J.; Anthonycahill, S. J.; Griffith, M. C.; Schultz, P. G. Science 1989, 244, 182-188.
- 8. Wang, L.; Brock, A.; Herberich, B.; Schultz, P. G. Science 2001, 292, 498-500.
- 9. Wang, L.; Brock, A.; Schultz, P. G. J. Am. Chem. Soc. 2002, 124, 1836-1837.
- 10. Wang, P., Ph.D. thesis, California Institute of Technology, Pasadena, 2003.
- Bain, J. D.; Glabe, C. G.; Dix, T. A.; Chamberlin, A. R.; Diala, E. S. J. Am. Chem. Soc.
 1989, 111, 8013-8014.
- 12. Kast, P.; Hennecke, H. J. Mol. Biol. 1991, 222, 99-124.
- 13. Sharma, N.; Furter, R.; Kast, P.; Tirrell, D. A. FEBS Lett. 2000, 467, 37-40.
- 14. Tang, Y.; Tirrell, D. A. Biochemistry 2002, 41, 10635-10645.
- 15. Wilson, M. J.; Hatfield, D. L. Biochim. Biophys. Acta 1984, 781, 205-215.
- Datta, D.; Wang, P.; Carrico, I. S.; Mayo, S. L.; Tirrell, D. A. J. Am. Chem. Soc. 2002, 124, 5652-5653.
- 17. Crick, F. H.; Brenner, S.; Watstobi, R. J.; Barnett, L. Nature 1961, 192, 1227-1232.
- 18. Crick, F. H. C. J. Mol. Biol. 1966, 19, 548-555.
- 19. Meroueh, M.; Chow, C. S. Nucleic Acids Res. 1999, 27, 1118-1125.

- Boren, T.; Elias, P.; Samuelsson, T.; Claesson, C.; Barciszewska, M.; Gehrke, C. W.;
 Kuo, K. C.; Lustig, F. J. Mol. Biol. 1993, 230, 739-749.
- Chen, P.; Qian, Q.; Zhang, S. P.; Isaksson, L. A.; Bjork, G. R. J. Mol. Biol. 2002, 317, 481-492.
- Inagaki, Y.; Kojima, A.; Bessho, Y.; Hori, H.; Ohama, T.; Osawa, S. J. Mol. Biol.
 1995, 251, 486-492.
- 23. Yarus, M. Science 1982, 218, 646-652.
- 24. Peterson, E. T.; Uhlenbeck, O. C. Biochemistry 1992, 31, 10380-10389.

ytRNA ^{Phe} AAA	+	+	-	+
yPheRS (T415G)	+	-	+	+
2Nal (3mM)	-	+	+	+
Phe (3mM)	+	-	-	-
Mass of intact mDHFR	23287	23287	23287	23537
Number of 2Nal residues	ND ^a	ND	ND	4.4

 Table 1: Molar masses and numbers of 2Nal residues observed for mDHFR samples

 prepared under various conditions

a. Not detected



Phenylalanine auxotrophic E. coli strain

Figure 1: A strategy for multi-site incorporation of 2Nal into recombinant proteins by breaking the degeneracy of the phenylalanine codons. ytRNA images were obtained from the Protein Data Bank (www.rcsb.org/pdb/).



Figure 2: SDS-PAGE analysis of mDHFR prepared in minimal media supplemented with 3 mM 2Nal and free of exogenous Phe. Conditions are noted at the top of each lane. Lane 1 shows molecular weight standards.



m/z

Figure 3: Replacement of Phe by 2Nal can be detected in MALDI-MS spectra of tryptic fragments of mDHFR. Peptide 1_{UUU} (residues 184-191, YKFEVYEK) contains a Phe (F) residue encoded by UUU, whereas in Peptide 1_{UUC} this codon has been mutated to UUC. Peptides 2 (residues 62-70, KTWFSIPEK) and 3 (residues 26-39, NGDLPWPPLRNEFK) are designated similarly. Peptide 1_{UUU} (Nal) refers to the form of the peptide containing 2Nal in place of Phe. In the absence of 2Nal, Peptide 1_{UUU} was detected at a mass of 1105.55 (A). Upon addition of 2Nal, a strong signal corresponding to Peptide 1_{UUU} (Nal) was detected at m/z = 1155.61, and the signal for Peptide 1_{UUU} was greatly reduced (B). For Peptide 1_{UUC} , 2Nal incorporation was much less efficient (C). Signals corresponding to Peptides 2_{UUC} (B) and 3_{UUC} (E) were not substantially reduced upon addition of 2Nal, and only very weak signals for Peptides 2_{UUC} (Nal) and 3_{UUC} (Nal) were detected (B, E). When the UUC codon in Peptide 3_{UUC} is mutated to UUU, a strong signal for Peptide 3_{UUU} (Nal) is detected (F). These data confirm that incorporation of 2Nal is strongly biased to UUU codons.



Figure 4: Tandem mass spectrum of Peptide 1_{UUU} (Nal) YKF*EVYEK. The doublycharged ion at 578.6 daltons was selected and fragmented. The sequence of the peptide containing 2Nal (F*) can be read from the annotated b (black) or y (red) ion series.

Chapter 6

Breaking the Degeneracy of the Leucine Codons in Escherichia coli

Abstract

Previously we have shown that combination of yPheRS (T415G) and the mutant ytRNA^{Phe}_{AAA} enables incorporation of 2Nal in response to UUU (Phe) codons. However, this method has two limitations. First, the yPheRS (T415G) activates Trp, which leads to Trp misincorporation. Second, 2Nal was incorporated at UUC codons as well as UUU codons, due to the relaxed codon recognition of the AAA anticodon in ytRNA^{Phe}_{AAA}. In order to enhance the selectivity of yPheRS, we performed high-throughput screening of a yPheRS library. The screen revealed a yPheRS variant (yPheRS_naph) with 6-fold higher relative activity toward 2Nal (vs. Trp) in ATP-PPi exchange assays. To enhance codon selectivity, we explored degenerate leucine codons instead of phenylalanine degenerate codons. Combined use of ytRNA^{Phe} containing the CAA anticodon, which recognizes only the UUG (Leu) codon, and yPheRS_naph allowed incorporation of 2Nal into murine dihydrofolate reductase in response to six UUG codons with 50% fidelity, but not to other Leu codon sites.

Introduction

An expanded set of genetically encoded amino acids has allowed design of proteins with novel chemical, physical, or biological properties. Several methods have been developed to introduce nonnatural amino acids into recombinant proteins at programmed sites *in vivo*. First, residue-specific incorporation involves the global replacement of a particular natural amino acid with a nonnatural analog in target proteins.¹⁻¹² The strength of this technique lies in efficient protein translation and multi-site incorporation of a nonnatural amino acid, because sense codons are re-assigned for a

nonnatural amino acid. However, its application could be restricted due to the exclusion of one natural amino acid for protein translation and lack of site-specificity. Second, single-site-specific incorporation involves utility of a heterologous orthogonal aminoacyltRNA synthetase/tRNA pair and suppression of either a stop codon or a frameshift codon.¹³⁻²⁰ Access to all 20 natural amino acids and site-specificity of the technique allows the incorporation of a nonnatural amino acid at a single site in a target protein with minimal perturbation of the native protein structure and function. However, application of the method is limited to the insertion of a nonnatural amino acid at only one or two positions in a target protein, due to moderate suppression efficiency.

Although residue-specific incorporation and single-site-specific incorporation have been used for many different applications,²¹⁻²⁷ one can envision other circumstances in which a nonnatural amino acid needs to be inserted at multiple positions in the presence of all twenty natural amino acids. For example, by introducing a nonnatural amino acid only at permissive sites in a target protein, the protein may be equipped with multiple reaction sites or probes with minimal loss of its native properties. Multiple-site-specific incorporation *in vivo* was realized in our lab by breaking the degeneracy of the Phe codons.²⁸ The combined use of a mutant yeast phenylalanyl-tRNA synthetase (yPheRS (T415G)) and mutant yeast phenylalanyl-tRNA containing modified AAA anticodon (ytRNA^{Phe}_{AAA}) that favors UUU Phe codons over UUC Phe codons, achieved efficient incorporation of 2Nal into mDHFR at five Phe sites encoded as UUU. Although the method successfully demonstrated the concept of breaking the degeneracy of the genetic code, its application has been restricted by two limitations.

First, the rationally designed yPheRS (T415G) efficiently activated a natural

148

amino acid, Trp, due to the enlarged binding pocket generated by the T415G mutation.¹³ The relaxed substrate specificity of yPheRS (T415G) showed several drawbacks. Cell growth rate decreased, likely due to misincorporation of Trp at Phe sites in essential proteins. Leaky expression of target protein even under uninduced conditions was prominent, perhaps due to the impaired repressor proteins. Furthermore, misincorporation of Trp as well as 2Nal was observed at Phe residues encoded as UUU codons, which prevented high fidelity incorporation of 2Nal at programmed sites. These drawbacks prompted us to pursue more selective yPheRS variants. Previously we reported that an other rationally designed yPheRS (T415A) showed 10-fold higher activity toward pBrF than Trp. However, vPheRS (T415A) did not exhibit enhanced selectivity toward pBrF against Phe.¹³ Therefore, we need to explore different approaches to obtain highly selective yPheRS variants. Aminoacyl-tRNA synthetases are known to be readily evolvable. Schultz and co-workers have developed powerful screening methods to change the substrate specificity of tyrosyl-tRNA synthetase (mjTyrRS) derived from Methanococcus jannaschii toward nonnatural amino acids.^{14,18,22,29} Recently our lab reported a novel screening method to adapt E. coli methionyl-tRNA synthetase to a reactive methionine analog, azidonorleucine.⁷ However, until now there have been no reports about evolving eukaryotic aminoacyl-tRNA synthetases to change their substrate specificity toward a nonnatural amino acid. In this report, we describe the highthroughput screening of a yPheRS library to obtain yPheRS variants of which substrate specificity is changed to a nonnatural amino acid, 2Nal.

Second, 2Nal was misincorporated at the unwanted sites (UUC codon) as well as programmed sites (UUU codon) due to relaxed codon recognition of the AAA anticodon

of ytRNA^{Phe}_{AAA}. Misincorporation of a nonnatural amino acid at unwanted sites might cause perturbation or loss of native properties of the target protein.^{23,30} According to Crick's wobble rules proposed in 1966,³¹ the base A in the first position of the anticodon recognizes only the base U in the third position of the codon. Therefore, the AAA anticodon was expected to recognize only UUU, not UUC codons. The discrepancy between the experimental result and Crick's wobble rule may be explained by the expanded wobble rule proposed by Lim and Curran in 2001.³² The expanded wobble rule is based on new experimental finding³²⁻³⁹ and stereochemical modeling⁴⁰⁻⁴³ of codon-anticodon interactions. The expanded wobble rule indicates that A in the first position of the anticodon of the anticodon can recognize all four bases in the third position of the codon, though its affinity to C is lower than that to U.

Due to the poor discrimination of UUU codon froms UUC by the AAA anticodon of ytRNA^{Phe}, we explored breaking the degeneracy of leucine (Leu) codons. Several considerations recommend degenerate Leu codons. First, Leu is encoded as six codons: UUA, UUG, CUA, CUG, CUU, and CUC. Discrimination of UUG from CUN (N = A/U/G/C) codons should be highly efficient due to discrimination at the first position in codon. Second, our existing yeast orthogonal pair should be readily adapted to the incorporation of Phe analogs in response to UUG codons. In practical terms, generalization of the concept of breaking the degeneracy of the genetic code is limited by the availability of orthogonal pairs. Third, the modified CAA anticodon would more efficiently discriminate UUG from UUA. According to the expanded wobble rules, C in the first position of the anticodon recognizes only G in the third position of the codon. In this report, we show that an engineered ytRNA^{Phe} containing a modified CAA anticodon (CAA) can completely discriminate UUG from the remaining five Leu codons, and achieve incorporation of 2Nal at multiple programmed sites in recombinant proteins.

Materials and Method

Materials. Unless otherwise noted, reagents were obtained from commercial suppliers and were used without further purification. 2Nal was purchased from Chem-Impex (Wood Dale, IL). Restriction enzymes were obtained from New England Biolabs (Beverly, MA). Quikchange mutagenesis kits were purchased from Stratagene (La Jolla, CA). Nickel-nitrilotriacetic acid affinity columns and plasmid pREP4 were obtained from Qiagen (Valencia, CA). DNA primers were obtained from Integrated DNA Technologies (Coralville, IA) and Operon Technologies (Huntsville, AL). Sequences of the DNA primers used in this research are listed in Appendix C.

Preparation of *E. coli* **hosts**. Preparation of the Phe/Trp double auxotrophic strain (AFW) and the Phe/Trp/Lys triple auxotrophic strain (AFWK) was described previously.¹³ A Phe/Leu double auxotrophic strain, MPC390 (*leuB6(Am)*, *PheA18::Tn10*), was obtained from the *E. coli* Genetic Stock Center (CGSC) at Yale University. A Phe auxotrophic derivative of DH10B (Stratagene) *E. coli* strain was prepared by chemical mutagenesis⁴⁴ and designated DHF.

Plasmid Construction for Reporter Gene Expression. The GFP_{UV} gene was amplified from pGFPuv (Clontech, Palo Alto, CA) using two primers (GFP1 and GFP2) containing a *Pst*I restriction site. The amplified GFPuv gene was inserted into pQE9

(Qiagen) at the PstI site to generate pQE9_GFP_{UV}. Since GFP_{UV} has excitation maxima at 475 and 395 nm,⁴⁵ it was mutated into an EGFP variant (GFP3)⁴⁵ with excitation maximum at 488 nm suitable for FACS analysis. A series of PCR mutagenesis reactions were performed at four residues (F64L, S65T, S99F and T153M) using three pairs of complementary primers (F64LS65T f/F64LS65T r; S99F f/S99F r; T153M_f/T153M_r). The GFP3 gene expression cassette was inserted into pQE16_mDHFR_yPheRS (T415G)²⁸ between the AatII and NheI restriction sites to generate pQE9_GFP3_yPheRS (T415G). GFP3 has twelve Phe residues, of which seven are encoded as UUC. A GFP variant (GFP6) was prepared by replacing all UUC Phe codons and one CUG Leu codon (at position 64) with UUU codons via gene fragment reassembly using sixteen primers (Primer 1-16). The GFP6 expression cassette was removed by restriction digestion with the AatII and NheI, and inserted into pQE9_GFP3_yPheRS (T415G) between the AatII and NheI sites to yield pQE9 GFP6 yPheRS (T415G). In order to introduce an extra copy of the *lacI* gene into pQE9_GFP6_yPheRS (T415G), the lacI gene expression cassette was amplified from pREP4 (Qiagen) with two primers (lacI_AflII_f and lacI_SacII_r). AflII and SacII restriction sites were introduced into pQE9_GFP6_yPheRS (T415G) by PCR reactions with two primers (pQE_AflII_f and pQE_SacII_r). The amplified lacI gene expression cassette was inserted into pQE9_GFP6_yPheRS (T415G) between the AflII and SacII sites to generate pQE9_GFP6_lacI_yPheRS (T415G). A GFP variant (GFP3_WC) containing 12 Phe residues encoded as only UUC codons were generated by the gene fragment reassembly method⁴⁶ using twelve primers (Primer 1, 16, 17-26). The GFP3_WC gene expression cassette was ligated into pQE9_GFP6_lacI_yPheRS (T415)

between the *Aat*II and *Nhe*I sites to yield pQE9_GFP3_WC_lacI_yPheRS (T415G). Information of the marker proteins used in this work is summarized in Appendix B.

Construction of vPheRS Library. Four residues (N412, T415, S418 and S437) in the active site of yPheRS were saturated by two step PCR mutagenesis. First, NNK (N = A/U/G/C; K = G/U) degenerate codons were introduced into the 437th position in the α subunit of yPheRS by PCR mutagenesis with two complementary primers (437 f and 437_r) using pQE9_GFP6_lacI_yPheRS (T415G) as a template. The PCR product was digested by *Dpn*I for 1 hr at 37 °C and transformed into XL-1 blue (Stratagene) competent cells. The plasmids isolated from the culture of transformants were used as a template for the second PCR mutagenesis with one pair of complementary primers (412 415 418 f and 412 415 418 r) to saturate the 412th, 415th and 418th positions in the α-subunit of yPheRS. The PCR product was digested by DpnI for 1 hr at 37 °C and desalted on a spin column. The eluted plasmids were transformed into ElectroTen-Blue electro-competent cells (Stratagene) according to the manufacturer's protocol, and plasmids were isolated from the culture of pooled transformants. The plasmid fragments encoding yPheRS were removed by digestion with NsiI and BglII restriction enzymes and ligated to the large fragments of pQE9_GFP6_lacI_yPheRS (T415G) obtained by digestion with NsiI and BglII. The ligated plasmids were desalted and transformed into ElectroTen-Blue electro-competent cells to generate six million transformants. The yPheRS library plasmids (P_yFS_20) were isolated from the culture of pooled transformants.

Construction of Expression Library. pREP4_ytRNA_AAA, prepared previously,²⁸ was transformed into DHF electro-competent cells to generate DHF_AAA cells. The P_yFS_20 library was transformed into DHF_AAA electro-competent cells to generate the yPheRS expression library (L_01) consisting of ten million transformants. The transformants were transferred to 0.5 L 2xYT media with 200 μ g/mL ampicillin and 35 μ g/mL kanamycin. When OD₆₀₀ reached 1.0, glycerol stocks were prepared and kept in a -80 °C freezer.

Screening of yPheRS Library. A half mL of glycerol stock of the expression library was inoculated into 100 mL of M9 minimal medium supplemented with glucose, thiamin, MgSO₄, CaCl₂, 20 amino acids (at 25 mg/L), antibiotics (35 µg/mL of kanamycin and 200 μ g/mL of ampicillin). When the culture reached an OD₆₀₀ of 0.6–0.8, cells were spun down, washed twice with ice-cold 0.9% NaCl, and shifted to expression medium supplemented with 18 amino acids (at 25 mg/L), and the indicated concentrations of Phe, Trp and 2Nal. Expression of GFP was induced by addition of 1 mM IPTG. After 3 hrs, 1 mL of the culture (based on OD of 1.0) was collected and washed twice with 0.5 mL of PBS (pH 7.4). 300 μ L of cells were diluted with 3 mL of distilled water, and then subjected to cell sorting using a MoFlo cell sorter (DakoCytomation, Ft. Collins, CO). The excitation and emission wavelengths were 488 and 525 nm, respectively. The library was subjected to both positive and negative screening (Figure 3). FACS gates were set based upon FSC/SSC and FL. In order to enrich active yPheRS variants, 5x10⁴ weakly fluorescent cells were collected by positive screening in the first round. The screened cells were incubated in 2xYT medium containing 200 µg/mL of ampicillin and 35 µg/mL of kanamycin. When OD_{600} reached 1.0, glycerol stocks of the cells were prepared. 0.1 mL of the glycerol stocks was inoculated into 20 mL of M9 minimal medium containing 20 amino acids. In order to enrich selective yPheRS variants, the top 1% of highly-fluorescent cells were collected. One more negative screening was applied to enrich the population in bright cells. The positive and negative screening steps were repeated. Finally, one more positive screening was performed, and the collected cells were spread onto a 2xYT agar plate containing 35 µg/mL of kanamycin and 200 µg/mL of ampicillin. After overnight incubation at 37 °C, ten single colonies were isolated and subjected to characterization.

Characterization of the Isolated Clones. The ten single colonies were transferred to 0.5 L 2xYT media with 200 µg/mL ampicillin and 35 µg/mL kanamycin. When OD₆₀₀ reached 1.0, glycerol stocks were prepared and kept in a -80 °C freezer. 200 µL of each glycerol stock was inoculated into minimal medium and incubated until OD₆₀₀ reached 0.6. The cultures were washed twice with 0.9% NaCl solution, and resuspended with 20 mL of minimal medium supplemented with 18 amino acids, 50 µM Trp, and 5 µM Phe. The resuspended cells were divided into two fractions, and transferred into two flasks. 3 mM 2Nal was supplemented into one of them. GFP6 was expressed by addition of 1 mM IPTG. After 3 hrs, 1 mL of the culture was collected, and washed twice with 0.5 mL of PBS (pH 7.4). 100 µL of cells were diluted with 3 mL of distilled water. Fluorescence intensities of the cells were analyzed by a MoFlo cell sorter. At least 20,000 events were collected in each measurement. Data were analyzed with Summit software (DakoCytomation). One positive clone showing weakly fluorescence with 2Nal and

highly fluorescence without 2Nal was subjected to further characterization. The plasmid isolated from the positive clone was subjected to DNA sequencing, which revealed that a yPheRS variant contained four mutations (N412G, T415G, S418C, and S437F). The yPheRS variant and the plasmid encoding it were designated yPheRS_naph and pQE9_GFP6_lacI_yPheRS_naph, respectively.

Amino Acid Activation Assay. pQE32-yPheRS (T415G) was prepared previously.¹³ The plasmid fragment encoding yPheRS removed from was pQE9_GFP6_lacI_yPheRS_naph by digestion with NsiI and BglII, and then ligated with a large fragment of pQE32-yPheRS (T415G) generated by digestion with NsiI and BglII to yield pQE32-yPheRS_naph. Expression and purification of yPheRS variants were described previously.¹³ The kinetics of activation of amino acids by yPheRS (T415G) and yPheRS_naph were determined by the amino acid-dependent adenosine triphosphate $(ATP)-[^{32}P]$ -pyrophosphate (PP_i) exchange assay. The assay buffer included 50 mM N-(2hydroxethyl) piperazine-N-(2-ethanesulfonic acid) (potassium-HEPES) (pH=7.6), 20 mM MgCl₂, 1 mM dithiothreitol (DTT), 2 mM ATP, and 2 mM [³²P]- PP_i. The amino acid concentration varied from 100 nM to 10 mM and the enzyme concentration varied from 100 nM to 800 nM. Detail procedures of the assay were described previously.¹³

Construction of Plasmids and Expression Hosts for Incorporation of 2Nal at Phe Codons. The gene fragment containing four mutations in the binding pocket of yPheRS_naph was excised by *Nsi*I and *Bgl*II digestion of pQE9_GFP6_lacI_yPheRS_naph and inserted into pQE16_mDHFR_yPheRS (T415G)

and pQE9_GFP3_WC_lacI_yPheRS (T415G) between the NsiI and Bg/II sites to generate pQE16_mDHFR_yPheRS_naph and pQE9_GFP3_WC_lacI_yPheRS_naph, respectively. Both pQE16_mDHFR_yPheRS (T415G) and pQE16_mDHFR_yPheRS naph were co-transformed with pREP4_ytRNA^{Phe}_AAA into AFW competent cells to generate AFW [pQE16 mDHFR yPheRS (T415G)/pREP4 ytRNA^{Phe} AAA] and AFW [pQE16_mDHFR_yPheRS_naph/pREP4_ytRNA^{Phe}_AAA], respectively. In order to express intact mDHFR, pQE16 (Qiagen) and pREP4 plasmids were co-transformed into AFW competent cells to generate AFW [pQE16/pREP4]. Both pQE9_GFP6_lacI_yPheRS_naph and pQE9_GFP3_WC_lacI_yPheRS_naph were transformed DHF AAA electrocompetent into cells to construct DHF [pQE9_GFP6_lacI_yPheRS_naph/pREP4_ytRNA^{Phe}_AAA] DHF and [pQE9_GFP3_WC_lacI_yPheRS_naph/pREP4_ytRNA^{Phe}_AAA], respectively. In order to determine Phe codon occupancy by various amino acids, the AGA (Arg) codon in the second position of GFP6 was mutated to either a UUU or a UUC (Phe) codon by PCR mutagenesis. The PCR reactions were conducted with two pairs of complementary $(2F_UUU_f/2F_UUU_r;$ $2F_UUC_f/2F_UUC_r$) primers using pQE9_GFP6_lacI_yPheRS_naph as a template to generate pQE9_GFP6 (2FUUU)_lacI_yPheRS_naph (2FUUC)_lacI_yPheRS_naph, pQE9_GFP6 and respectively. Removal of a *lac* promoter and a *lac* operator, and replacement of a weak ribosome binding site with a stronger one in the upstream of α -subunit of yPheRS gene were performed by two QuickChange mutagenesis reactions using two pairs of complementary primers (R lacP f/R lacP r and Ins_SD_A_Del_lacO_f/Ins_SD_A_Del_lacO_r) using (2FUUU)_lacI_yPheRS_naph and

pQE9_GFP6 (2FUUC)_lacI_yPheRS_naph as templates to generate pQE9_GFP6 (2FUUU)_lacI_SD_yPheRS_naph and pQE9_GFP6 (2FUUC)_lacI_SD_yPheRS_naph and pQE9_GFP6 (2FUUC)_lacI_SD_yPheRS_naph were transformed into DHF_AAA electrocompetent cells to construct DHF [pQE9_GFP6 (2FUUU)_lacI_SD_yPheRS_naph/pREP4_ytRNA^{Phe}_AAA] and DHF [pQE9_GFP6 (2FUUC)_lacI_SD_yPheRS_naph/pREP4_ytRNA^{Phe}_AAA].

Residue- and Site-Specific Incorporation of 2Nal into Recombinant Proteins. The AAA anticodon of ytRNA^{Phe}AAA was mutated into GAA by PCR mutagenesis with one pair of primers (ytRNA-GAA_f/ytRNA-GAA_r) using pREP4_ytRNA_AAA as a template. The resulting pREP4_ytRNA_GAA plasmid contained the ytRNA^{Phe}_{GAA} gene. Both pQE9_GFP6 (2FUUU)_lacI_SD_yPheRS_naph and pREP4_ytRNA^{Phe}_GAA were co-transformed into DHF competent cells to generate DHF [pQE9 GFP6 (2FUUU)_lacI_SD_yPheRS_naph/pREP4_ytRNA^{Phe}_GAA]. pREP4_ytRNA_UG was prepared previously.¹³ The yPheRS_naph gene fragment excised by NsiI and Bg/II digestion of pQE9_GFP6_lacI_SD_yPheRS_naph was inserted into pQE16_mDHFR (38Am)_yPheRS (T415G) between the NsiI and BglII sites to generate pQE16_mDHFR (38Am)_SD_yPheRS_naph. An AAG (Lys) codon was changed to a UAG codon at position 158 of GFP6 sequence in pQE9_GFP6_lacI_yPheRS_naph by PCR mutagenesis with one pair of complementary primers (K158_UAG_f/K158_UAG_r) to yield pQE9_GFP6 (158Am)_lacI_SD_yPheRS_naph. Both pQE16_mDHFR (38Am)_yPheRS_naph and pQE9_GFP6 (158Am)_lacI_SD_yPheRS_naph were cotransformed with pREP4_ytRNA^{Phe}_UG into AFWK and DHF competent cells to construct AFKW [pQE16_mDHFR (38Am)_yPheRS_naph/pREP4_ytRNA^{Phe}_UG] and DHF [pQE9_GFP6 (158Am)_lacI_SD_yPheRS_naph/pREP4_ytRNA^{Phe}_UG], respectively.

Construction of Plasmids and Expression Hosts for Incorporation of 2Nal at Leu Codons. The AAA anticodon of ytRNA^{Phe}AAA was mutated to CAA by PCR mutagenesis with primers ytRNA-CAA_f and ytRNA-CAA_r using pREP4_ytRNA^{Phe}_AAA as a template to yield pREP4_ytRNA^{Phe}_CAA. The expression cassette of mDHFR was excised from pQE16 (Qiagen) by digestion with AatII and NheI and inserted into pQE9_GFP6_lacI_yPheRS_naph between the AatII and NheI sites to generate pQE16_mDHFR_lacI_yPheRS_naph. In order to increase the number of Leu residues encoded as UUG, UUC and UUU (Phe) codons in position 38 and 95 of mDHFR were changed to UUG by PCR mutagenesis reactions with two complementary pairs of primers using pQE16_mDHFR_lacI_yPheRS_naph as a template to generate pQE16_mDHFR2_lacI_yPheRS_naph. PCR mutagenesis reaction was performed to mutate UUG to UUA at position 100 of mDHFR2 with two complementary primers (L100_UUA_f/L100_UUA_r) to yield pQE16_mDHFR2 (100UUA)_lacI_yPheRS_naph. Either pQE16_mDHFR2_lacI_yPheRS_naph or pQE16 mDHFR2 (100UUA) lacI yPheRS naph was co-transformed with ytRNA^{Phe}_{CAA} into MPC390 competent cells MP to yield [pQE16_mDHFR2_lacI_yPheRS_naph/pREP4_ytRNA^{Phe}_CAA] or [pQE16_mDHFR2 (100UUA) lacI yPheRS naph/pREP4 ytRNA^{Phe} CAA], respectively. In order to

express intact mDHFR, pQE16 (Qiagen) and pREP4 were co-transformed into MPC390 competent cells to generate MP [pQE16/pREP4]. In order to generate GFP with Cterminal hexa-histidine tag (GFP3C), a PstI restriction site was added to pQE70 (Qiagen) using two complementary primers (PstI 70 f/PstI 70 r) to yield pQE70*. The coding sequence for GFP3 was excised by *PstI* digestion from pQE9 GFP3 yPheRS (T415G) and inserted into pQE70* at the PstI site to generate pQE70*_GFP3C. Expression cassette of GFP3C isolated from pQE70*_GFP3C inserted was into pQE9_GFP3_lacI_yPheRS_naph to yield pQE70*_GFP3C_lacI_yPheRS_naph. Lys 158 in GFP3C was mutated to Leu (encoded as UUG) by PCR mutagenesis with two complementary (K158 UUG f/K158 UUG r) primers using pQE70*_GFP3C_lacI_yPheRS_naph as a template to yield pQE70*_GFP3C (158UUG)_lacI_yPheRS_naph. Both pQE70*_GFP3C_lacI_yPheRS_naph and (158UUG)_lacI_yPheRS_naph pQE70*_GFP3C co-transformed with were pREP4 ytRNA^{Phe} CAA into **MPC390** cells competent to construct MP [pQE70* GFP3C lacI yPheRS naph/pREP4 ytRNA^{Phe} CAA] MP and [pQE70*_GFP3C (158UUG)_lacI_yPheRS_naph/ pREP4_ytRNA^{Phe} CAA], respectively.

Expression of mDHFR Variants and GFP Variants in Vivo. Both AFW and AFWK expression strains co-transformed with pQE plasmid variants and pREP4 plasmid variants were grown in M9 minimal medium supplemented with 0.4 wt % glucose, 35 mg/L thiamin, 1mM MgSO₄, 1mM CaCl₂, 20 amino acids (at 25 mg/L), 35 mg/L kanamycin, and 200 mg/L ampicillin. The overnight cultures of expression strains were diluted 20-fold in fresh M9 minimal medium and incubated at 37 °C. When cells reached

OD of 0.8 - 1.0, cells were spun down and washed twice with cold 0.9% NaCl. The cultures were resuspended in fresh M9 minimal medium supplemented with 18 amino acids (25 μg/mL), and the indicated concentrations of Phe, Trp, and 2Nal. After 10 min incubation, 1 mM of isopropyl-β-D-thiogalactopyranoside (IPTG) was added to induce expression of GFP at 30 °C. After 4 hours, cells were harvested and either kept at -80 °C or subjected to fluorescence measurement according to the procedures described earlier (see Characterization of the Isolated Clones). Whole cell lysates were analyzed by SDS-PAGE. Due to slow growth of DHF and MPC390 expression hosts co-transformed with pQE plasmid variants and pREP4 plasmid variants, transformants were grown in 2xYT medium to prepare glycerol stocks first. Then glycerol stocks were inoculated into minimal medium supplemented with 20 amino acids (at 25 mg/L) and incubated overnight at 37 °C. The remaining steps were similar to those for AF and AFWK expression hosts.

Quantitative Analysis of Codon Occupancy. Quantitative analysis of codon occupancy was performed by either N-terminal protein sequencing or LC-MS analysis of tryptic digests. The GFP6 (2UUU) and GFP6 (UUC) variants were expressed in minimal medium and purified by Ni-NTA affinity chromatography according to the manufacturer's protocol (Qiagen) under denaturing conditions. The purified GFP variants were subjected to N-terminal protein sequencing using a 492 cLC Procise protein microsequencer (Applied Biosystems, Foster City, CA). Occupancy of Phe codons in mDHFR and Leu codons in GFP was determined by LC-MS analysis. mDHFR expressed in minimal medium were subjected to purification via Ni-NTA affinity chromatography

according to the manufacturer's protocol (Qiagen) under denaturing conditions. After purification, expression levels of GFP and mDHFR were determined by UV absorbance at 280 nm using a calculated extinction coefficient of 20,010 cm⁻¹ M⁻¹ and 24, 750 cm⁻¹ M^{-1} , respectively. The purified proteins were concentrated by ultrafiltration (Millipore). 10 µL of the concentrate was diluted into 90 µL of 75 mM (NH₄)₂CO₃ solution and then 1 µL of modified trypsin (Promega, 0.2 µg/µL) was added. Reaction was carried out for 2-4 hrs at 37 °C and quenched by addition of 13 µL of 5% trifluoroacetic acid (TFA) solution. The solution was then directly subjected to LC-MS analysis conducted on a LCT Premier XE MICROMASS MS system (MS Technologies, Montgomery Village, MD) with Acquity UPLCTM system (Waters, Milford, MA). Tryptic digests were separated by Acquity BEH300 C18 column (1.7 µm, 300 Å, 2.1 x 50mm) using a gradient of 5-95% of solvent B (90% of acetonitrile/10% of 0.1% formic acid solution) and solvent A (2% of acetonitrile/98% of 0.1% formic acid solution) in 10 min. The column eluent was transferred to the electrospray source and mass spectra were recorded.

Results and Discussion

Misincorporation of Trp into Recombinant Proteins in Response to UUU Codons. mDHFR was expressed in AFW [pQE16_mDHFR_yPheRS (T415G)/pREP4_ytRNA^{Phe}_AAA] expression hosts in minimal medium supplemented with 18 amino acids (MM18_FW), 2.5 μ M Phe, 3 mM Trp. Occupancy of the UUU codon sites was determined by LC-MS analysis of tryptic digests of mDHFR expressed with or without 3 mM 2Nal. We focused on Peptide 1 (residues 140-144; LF_{UUU}VTR), which contains a Phe residue at position 141 as UUU. LC-MS analysis of Peptide 1 variants indicated that 10% of position 141 was occupied by Trp (Figure 1b). Trp misincorporation at UUU site can be explained by the relaxed substrate specificity of yPheRS (T415G). yPheRS (T415G) showed 2-fold higher k_{cat}/K_m for Trp than for 2Nal in ATP-PPi exchange assays (Table 1).

Construction of yPheRS Library. In order to enhance the specificity of yPheRS with respect to 2Nal, we explored high-throughput screening of a yPheRS library. On the basis of the crystal structure of the homologous PheRS (tPheRS) from *Thermus thermophilus*⁴⁷ and sequence alignment between tPheRS and yPheRS, four residues (N412, S418, T415, and S437) within 7 Å of the *para* position of the phenyl ring of Phe bound to yPheRS were subjected to saturation mutagenesis (Figure 2). PCR mutagenesis generated 6x10⁶ yPheRS transformants (see Materials and Methods), a population somewhat larger than the theoretical library size (10⁶). By transforming yPheRS library plasmids into DHF expression hosts, 1x10⁷ transformants were obtained and pooled to construct the yPheRS expression library.

High-throughput Screening of yPheRS Library. Screening of the yPheRS library entailed both positive and negative screenings to obtain active and selective yPheRS variants, respectively. Active synthetases would allow incorporation of either 2Nal or natural amino acids at non-permissive sites of GFP6, where amino acids other than Phe would lead to reduction in the fluorescence of cells. Our preliminary study showed that misincorporation of 2Nal at 12 Phe sites of GFP6 resulted in more than 20-fold reduction in fluorescence of cells, likely due to the perturbed folded structure of GFP6 (data not

shown). For positive screening, the yPheRS library was induced with 1 mM IPTG to express GFP6 in minimal medium supplemented with 3 mM 2Nal. The fluorescence histogram of yPheRS library cells expressing GFP6 was obtained by flow cytometry. The gate in the fluorescence channel was set to collect 1% of cells for which fluorescence was slightly above background. The 5×10^4 cells collected were regrown in 2xYT medium and subjected to negative screening to enrich cells containing selective yPheRS variants. The negative screening was based on the fact that synthetases selective for 2Nal should not misincorporate any natural amino acid at Phe sites in GFP6 but E. coli endogenous Phe orthogonal pair will incorporate Phe at the Phe sites. The resulting GFP6 would retain full intensity of fluorescence. For negative screening, GFP6 was expressed in minimal medium in the absence of 2Nal. The top 1% of cells in the fluorescence channel was collected to obtain 5×10^4 cells. The collected cells were regrown in 2xYT medium and entered a next round of screening. After two rounds of positive and negative screening, one more positive screening was carried out to enrich active yPheRS variants toward 2Nal. A portion of the collected cells were spread on agar plates containing suitable antibiotics and ten colonies were subjected to further characterization.

Characterization of the Selective yPheRS Variant. One out of ten clones showed the fluorescence characteristics expected of a selective yPheRS variant. The other clones were false positives that have two peaks coincidently overlapped with the gates for sorting in the histogram of fluorescence. In the absence of 2Nal, cells expressing GFP6 retained full intensity of fluorescence (Figure 4a). However, in the presence of 2Nal, cells expressing GFP6 showed substantially reduced intensity of fluorescence (Figure 4b).

Similar trends were observed in the visual comparison of fluorescence of cell pellets (Figure 4c). The plasmid DNA coding the selective yPheRS variant was isolated from the culture of the clone and subjected to further analysis. DNA sequencing analysis of the plasmid showed that there were mutations of all four positions (N412G, T415G, S418C, and S437F) in the binding pocket of yPheRS. The isolated yPheRS variant was designated yPheRS_naph.

Amino Acid Activation Analysis by yPheRS_naph. Activation of Phe, Trp and 2Nal by both yPheRS (T415G) and yPheRS_naph were examined *in vitro*. The kinetic parameters are listed in Table 1. yPheRS (T415G) showed 2-fold lower activity toward 2Nal than Trp. However, the yPheRS_naph showed 6-fold higher activity toward 2Nal than Trp, which translates into 12-fold enhanced selectivity toward 2Nal (vs. Trp) compared to yPheRS (T415G). The yPheRS_naph also showed 17-fold higher activity toward 2Nal than Phe, while yPheRS (T415G) showed 8.4-fold higher activity. Previously we reported that rationally designed yPheRS (T415A) showed 10-fold higher activity toward pBrF than Trp. However, the activation of Phe by yPheRS (T415A) was comparable to that of pBrF. Therefore, yPheRS_naph is the first yPheRS variant that efficiently discriminates a nonnatural amino acid from all twenty natural amino acids.

The poor binding of Phe and Trp by yPheRS_naph may be understood by loss of favorable interactions between the aromatic rings of the substrate and active-site residues of the synthetase. In a crystal structure of tPheRS,⁴⁷ the side chain of Phe 258 (which corresponds to Asn 412 in yPheRS) makes direct contact with the phenyl ring of the substrate. Since the interaction makes the substrate recognition highly specific and very
favorable energetically, we can readily assume that N412G mutation in yPheRS would lead to poor binding of aromatic amino acid by the PheRS. However, the S418C mutation may not be critical to substrate specificity for 2Nal, because the yPheRS variant containing only three mutations (N412G, T415G and S437F) showed activity toward 2Nal similar to that of yPheRS_naph in ATP-PPi exchange assays. Analysis of fluorescence changes upon incorporation of 2Nal revealed that omission of the S437F mutation leads to a loss of activity toward 2Nal (data not shown). Therefore, S437F mutation was thought to play a key role in discriminating 2Nal from both Phe and Trp.

Elimination of Trp Misincorporation at UUU Codons by yPheRS_naph in Vivo. AFW [pQE16_mDHFR_yPheRS_naph/pREP4_ytRNA^{Phe}_AAA] cells were induced to express mDHFR in MM18_FW medium supplemented with 2.5 μM Phe, 50 μM Trp and 3 mM 2Nal. Peptide 1 (residues 140-144; LF_{UUU}VTR), one of the tryptic fragments of purified mDHFR, contains a Phe residue encoded as UUU. Occupancy of the UUU site in Peptide 1 was investigated by LC-MS analysis. The results revealed that use of yPheRS_naph completely eliminated misincorporation of Trp at the UUU site (Figure 1c), while yPheRS (T415G) allowed misincorporation of Trp at the UUU site (Figure 1b). Similar results were obtained by N-terminal sequencing of purified intact GFP6 (2UUU). DHF [pQE9_GFP6_lacI_yPheRS_naph/pREP4_ytRNA^{Phe}_AAA] cells were induced to express GFP6 (2UUU) in MM18_FW medium supplemented with 50 μM Trp and 3 mM 2Nal. 80% and 20% of the UUU codon at the 2nd position of GFP6 (UUU) were decoded as 2Nal and Phe, respectively; but Trp was not detected at this position (Table 2).

Residue- and Single-Site-Specific Incorporation of 2Nal in Vivo. With an appropriate tRNA, the selective yPheRS_naph variant can be used for residue- and single-sitespecific incorporation of 2Nal into proteins. In order to realize residue-specific incorporation of 2Nal, DHF [pQE9 GFP6 (2UUU) lacI yPheRS naph/pREP4 ytRNA^{Phe} GAA] expression hosts were induced to express GFP6 (2UUU) in MM18_FW medium supplemented with 50 µM Trp and 3 mM 2Nal. N-terminal sequencing of the purified GFP6 (2UUU) showed that 92% of position 2 was occupied by 2Nal (Table 2), slightly higher than the 80% occupancy achieved by multiple-site-specific incorporation. The enhanced 2Nal incorporation may be a consequence of the known 12-fold higher aminoacylation rate for ytRNA^{Phe}_{GAA} by yPheRS as compared to ytRNA^{Phe}AAA.⁴⁸ Single-site-specific incorporation of 2Nal into mDHFR_38Am achieved AFWK was by [pQE16_mDHFR_yPheRS_naph/pREP4_ytRNA^{Phe}_UG] in miminal medium supplemented with 25 µg/mL 17 amino acids (MM17 FWK), 50 µM Phe, 50 µM Trp, 50 µM Lys, and 3 mM 2Nal. MALDI-MS analysis of tryptic digests of mDHFR_38Am revealed that 2Nal was dominant at the amber site. Neither Trp nor Phe was detected, confirming the high selectivity of yPheRS_naph toward 2Nal.

Single-Site-Specific Incorporation of 2Nal into GFP in Vivo. Single-site-specific incorporation of 2Nal into GFP6 was investigated to determine whether a correctly folded fluorescent protein can be obtained upon the addition of a nonnatural amino acid. An AAG (Lys) codon was changed to an amber codon in position 158 in GFP6, which is known to be permissive to replacement of Lys with other natural amino acids or to

circular permutation.⁴⁹ DHF [pQE9_GFP6 (158Am)_lacI_yPheRS_naph/pREP4_ytRNA^{Phe}_ UG] cells were used to express GFP6 (158Am) in MM17_FWK medium supplemented with 50 μ M Phe, 50 μ M Trp, 50 μ M Lys, and 3 mM 2Nal. The fluorescence of cells expressing full length of GFP6 (158Am) was 280-fold higher than that in uninduced conditions, which clearly indicated that 2Nal can be inserted into GFP6 without substantial reduction of fluorescence.

Misincorporation of 2Nal at Unwanted Sites (UUC Codons) in Vivo. Site-specific incorporation involves introduction of a nonnatural amino acid into a target protein only at programmed sites. Introduction of a nonnatural amino acid at unwanted site will diminish its advantages over the residue-specific incorporation method. Although AFW [pQE16 mDHFR yPheRS (T415G)/pREP4 ytRNA^{Phe} AAA] realized incorporation of 2Nal at programmed UUU codons in mDHFR (Figure 1c), misincorporation of 2Nal at UUC codons was also observed (Figure 1f). Similar results were observed in two GFP variants, GFP6 (2UUU) and GFP6 (2UUC), expressed in DHF [pQE9_GFP6 (2UUC)_lacI_yPheRS_naph/pREP4_ytRNA^{Phe}_AAA] and [pQE9_GFP6 (2UUU)_lacI_yPheRS/pREP4_ytRNA^{Phe}_AAA], respectively. Supplementation of 2.5 μM; 5 μM Phe into expression medium lowered 2Nal misincorporation at the unwanted sites (UUC codons) from 34% to 20%; 6%. However, the incorporation level of 2Nal at the programmed sites (UUU codons) also decreased from 80% to 62%; 47% (Figure 6). Incorporation of 2Nal at UUC sites in GFP6 (ca. 20%) with supplementation of 2.5 µM Phe is higher than that in mDHFR (ca. 10%) (Figure 1f), which might be explained by more stable characteristics of GFP6 containing 2Nal as compared to mDHFR. Even 6%

of misincorporation of 2Nal at unwanted sites in a target protein could be unfavorable for the target protein to retain its native properties. As a test case, 2Nal misincorporation into GFP6_WC, which contains 12 unwanted sites, led to 10-fold reduction in the fluorescence of cells (Figure 8).

We reasoned that misincorporation of 2Nal at UUC codon resulted from recognition of UUC codons by the AAA anticodon of ytRNA^{Phe}_{AAA}. According to Crick's wobble rule proposed in 1966,³¹ A in the first position of the anticodon can recognize only U in the third position of the codon. Therefore, UUC codon should not be recognized by the AAA anticodon. The discrepancy between the experimental results and Crick's wobble rule may be explained by the expanded wobble rule proposed by Lim and Curran in 2001.³² According to the expanded wobble rule, A in the first position of the anticodon can recognize all four bases in the third position in codon. The base A in the first position of the anticodon favors bases in the order U > C > G > A, consistent with the codon-biased incorporation of 2Nal observed in this work.

Breaking the Degeneracy of the Leucine Codons. According to the expanded wobble rules, C in the first position of the anticodon will recognize only G in the third position of the codon. Therefore, we hypothesized that $ytRNA^{Phe}_{CAA}$ (containing the modified CAA anticodon) would selectively recognize UUG codons. In order to test this hypothesis, mDHFR was expressed in MP [pQE16_mDHFR2_lacI_yPheRS_naph/pREP4_ytRNA^{Phe}_CAA] cells. mDHFR contains twenty Leu codons, of which are six UUG, two UUA and twelve CUN (N = A/T/G/C). The expression level of mDHFR was 2.9 mg/L. Occupancy of each Leu codon by various

amino acids was determined by LC-MS analysis of tryptic digests of mDHFR expressed with and without 2Nal. We focused on four peptides. Peptide **3** (residues 165-180; $L_{CUU}L_{CUC}PEYPGVL_{CUC}SEVQEEK$) contains three Leu residues, encoded as CUU and CUC codons. Peptide **4** (residues 54-61; QNL_{CUG}VIMGR) contains a Leu residue, encoded as a CUG codon. Peptide **5** (residues 62-70; $L_{CUU}IEQPEL_{UUG}ASK$) contains two Leu residues, encoded as CUU and UUG codons. Peptide **6** (residues 99-105; $SL_{UUG}DDAL_{UUA}R$) contains two Leu residues, encoded as UUG and UUA codons. 2Nal was not detected at any CUN codon in Peptide **3** and **4** (Figure 8a-d). However, 50% of UUG codons in Peptide **5** and **6** were occupied by 2Nal (Figure 8e-h). In order to determine UUA codon occupancy by 2Nal, Peptide **6**_{UUA} was tested. Peptide **6**_{UUA} is the same as Peptide **6** except both Leu residues are encoded as UUA codons. Since the Peptide **6**_{UUA} variant containing 2Nal was not detected, we conclude that 2Nal incorporation is highly specific to the UUG codon.

Codon-Specific Incorporation of 2Nal. Previously we UUG showed that misincorporation of 2Nal at unwanted UUC codons in GFP6_WC led to 10-fold reduction in fluorescence of cells, even though there are no UUU codons in GFP6_WC. Similar to GFP6_WC, we prepared GFP3 containing twenty three Leu codons, of which are no UUG, four UUA and nineteen CUN (N = A/T/G/C). In order to investigate effect of misincorporation of 2Nal at unwanted sites (Leu codons other than UUG), GFP3 was Ε. MP expressed in coli strain [pQE9_GFP3_lacI_yPheRS_naph/pREP4_ytRNA^{Phe}_CAA]. The fluorescence intensities of cells expressing GFP3 without 2Nal or with 2Nal were compared (Figure 7c and d).

There was no detectable difference in the fluorescence of cells prepared under these two conditions, confirming the absence of misincorporation of 2Nal at Leu codons other than UUG.

Minimal Perturbation of Native Properties of GFP upon Incorporation of 2Nal. Ideally introduction of a nonnatural amino acid should impart new properties to a target protein with minimal or no perturbation of native properties. As a test case, we chose fluorescence change of GFP variant upon 2Nal incorporation to evaluate perturbation of GFP structure. A UUG codon, an incorporation site for 2Nal, was introduced at the 158th position of GFP3 to generate GFP3 (158UUG) variant. GFP3 (158UUG) variant was expressed in minimal medium supplemented with 17 amino acids (25 μ g/mL), 50 μ M Phe, 50 μ M Trp, 1.25 μ M Leu, and indicated concentration of 2Nal. There was no significant differences in fluorescence histograms of cells expressing GFP3 (158UUG) in the absence and presence of 2Nal, which meant that GFP3 (158UUG) retained its folded structure even upon 2Nal incorporation at the 158th position.

Conclusions

In this chapter, we have shown that the substrate specificity of yPheRS was altered to selectively recognize a nonnatural amino acid, 2Nal. As far as we know, this is the first report of high-throughput screening of a eukaryotic aminoacyl-tRNA synthetase library to obtain synthetase variants selective toward nonnatural amino acids. Use of the selective yPheRS_naph variant resulted in elimination of misincorporation of Trp at UUU codons, which was detected previously in experiments with yPheRS (T415G). Combined use of yPheRS_naph and ytRNA^{Phe}_{CUA_UG} realized high fidelity (ca. 98%) incorporation of 2Nal into proteins in response to an amber codon.

Misincorporation of 2Nal at unwanted sites resulting from the relaxed codon recognition of the AAA anticodon of ytRNA^{Phe} has been overcome by use of the more codon selective ytRNA^{Phe}_{CAA}. The CAA anticodon of ytRNA^{Phe}_{CAA} completely discriminates UUU codons from other five Leu codons. When both yPheRS_naph and ytRNA^{Phe}_{CAA} were overexpressed in *E. coli* expression hosts, 50% of UUG codon sites were occupied by 2Nal, but no other Leu codon sites were occupied by 2Nal, but no other Leu codon sites were occupied by 2Nal. Combined use of yPheRS_naph and the codon selective ytRNA^{Phe}_{CAA} has realized multiple-site-specific incorporation of 2Nal into proteins. We are working to improve the level of incorporation of 2Nal at programmed UUG sites

Acknowledgments

We thank Dr. M. Shahgholi and Dr. J. Zhou for mass spectrometry studies. We also thank Dr. F. Rusnak for help with N-terminal sequencing. We are grateful Dr. A. J. Link for help with the cell sorter. We appreciate I. C. Tanrikulu and P. M. Kekenes-Huskey for help with analysis of the PheRS crystal structure. This work was supported by National Institutes of Health grant.

References

Bae, J. H.; Rubini, M.; Jung, G.; Wiegand, G.; Seifert, M. H. J.; Azim, M. K.; Kim, J. S.; Zumbusch, A.; Holak, T. A.; Moroder, L.; Huber, R.; Budisa, N. *J. Mol. Biol.* 2003, *328*, 1071-1081.

- Budisa, N.; Minks, C.; Medrano, F. J.; Lutz, J.; Huber, R.; Moroder, L. Proc. Natl. Acad. Sci. U. S. A. 1998, 95, 455-459.
- Budisa, N.; Rubini, M.; Bae, J. H.; Weyher, E.; Wenger, W.; Golbik, R.; Huber, R.; Moroder, L. Angew. Chem. Int. Ed. 2002, 41, 4066-4069.
- Datta, D.; Wang, P.; Carrico, I. S.; Mayo, S. L.; Tirrell, D. A. J. Am. Chem. Soc. 2002, 124, 5652-5653.
- Kiick, K. L.; Saxon, E.; Tirrell, D. A.; Bertozzi, C. R. Proc. Natl. Acad. Sci. U. S. A.
 2002, 99, 19-24.
- 6. Kirshenbaum, K.; Carrico, I. S.; Tirrell, D. A. ChemBioChem 2002, 3, 235-237.
- Link, A. J.; Vink, M. K. S.; Agard, N. J.; Prescher, J. A.; Bertozzi, C. R.; Tirrell, D. A. Proc. Natl. Acad. Sci. U. S. A. 2006, 103, 10180-10185.
- 8. Pratt, E. A.; Ho, C. Fed. Proc. 1974, 33, 1463-1463.
- 9. Sharma, N.; Furter, R.; Kast, P.; Tirrell, D. A. FEBS Lett. 2000, 467, 37-40.
- 10. Tang, Y.; Tirrell, D. A. J. Am. Chem. Soc. 2001, 123, 11089-11090.
- 11. Tang, Y.; Tirrell, D. A. Biochemistry 2002, 41, 10635-10645.
- 12. Wang, P.; Tang, Y.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 6900-6906.
- 13. Kwon, I.; Wang, P.; Tirrell, D. A. J. Am. Chem. Soc. 2006, 128, 11778-11783.
- 14. Anderson, J. C.; Schultz, P. G. Biochemistry 2003, 42, 9598-9608.
- 15.Hohsaka, T.; Ashizuka, Y.; Taira, H.; Murakami, H.; Sisido, M. *Biochemistry* **2001**, *40*, 11060-11064.
- 16.Kowal, A. K.; Kohrer, C.; RajBhandary, U. L. Proc. Natl. Acad. Sci. U. S. A. 2001, 98, 2268-2273.
- 17. Wang, J. Y.; Xie, J. M.; Schultz, P. G. J. Am. Chem. Soc. 2006, 128, 8738-8739.

- 18. Wang, L.; Schultz, P. G. Chem. Commun. 2002, 1-11.
- 19. Zeng, H. Q.; Me, J. M.; Schultz, P. G. Bioorg. Med. Chem. Lett. 2006, 16, 5356-5359.
- Zhang, Z. W.; Gildersleeve, J.; Yang, Y. Y.; Xu, R.; Loo, J. A.; Uryu, S.; Wong, C. H.;
 Schultz, P. G. *Science* 2004, *303*, 371-373.
- 21. Beatty, K. E.; Xie, F.; Wang, Q.; Tirrell, D. A. J. Am. Chem. Soc. 2005, 127, 14150-14151.
- 22. Bose, M.; Groff, D.; Xie, J. M.; Brustad, E.; Schultz, P. G. J. Am. Chem. Soc. 2006, 128, 388-389.
- 23. Budisa, N.; Pal, P. P. Biol. Chem. 2004, 385, 893-904.
- 24. Chin, J. W.; Santoro, S. W.; Martin, A. B.; King, D. S.; Wang, L.; Schultz, P. G. J. Am. Chem. Soc. 2002, 124, 9026-9027.
- 25. Link, A. J.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 11164-11165.
- 26. Wang, L.; Xie, J. M.; Deniz, A. A.; Schultz, P. G. J. Org. Chem. 2003, 68, 174-176.
- 27. Xie, J. M.; Wang, L.; Wu, N.; Brock, A.; Spraggon, G.; Schultz, P. G. Nat. Biotechnol.
 2004, 22, 1297-1301.
- 28. Kwon, I.; Kirshenbaum, K.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 7512-7513.
- Santoro, S. W.; Wang, L.; Herberich, B.; King, D. S.; Schultz, P. G. Nat. Biotechnol.
 2002, 20, 1044-1048.
- 30. Montclare, J. K.; Tirrell, D. A. Angew. Chem. Int. Ed. 2006, 45, 4518-4521.
- 31. Crick, F. H. C. J. Mol. Biol. 1966, 19, 548-555.
- 32. Lim, V. I.; Curran, J. F. RNA 2001, 7, 942-957.
- Chen, P.; Qian, Q.; Zhang, S. P.; Isaksson, L. A.; Bjork, G. R. J. Mol. Biol. 2002, 317, 481-492.

- 34. Watanabe, Y.; Tsurui, H.; Ueda, T.; Furushima, R.; Takamiya, S.; Kita, K.; Nishikawa, K.; Watanabe, K. *Biochim. Biophys. Acta-Gene Struct. Express.* 1997, *1350*, 119-122.
- Inagaki, Y.; Kojima, A.; Bessho, Y.; Hori, H.; Ohama, T.; Osawa, S. J. Mol. Biol.
 1995, 251, 486-492.
- 36. Curran, J. F. Nucleic Acids Res. 1995, 23, 683-688.
- Boren, T.; Elias, P.; Samuelsson, T.; Claesson, C.; Barciszewska, M.; Gehrke, C. W.;
 Kuo, K. C.; Lustig, F. J. Mol. Biol. 1993, 230, 739-749.
- 38. Osawa, S.; Jukes, T. H.; Watanabe, K.; Muto, A. Microbiol. Rev. 1992, 56, 229-264.
- 39. Munz, P.; Leupold, U.; Agris, P.; Kohli, J. Nature 1981, 294, 187-188.
- 40. Lim, V. I.; Aglyamova, G. V. Biol. Chem. 1998, 379, 773-781.
- 41. Lim, V. I. J. Mol. Biol. 1997, 266, 877-890.
- 42. Lim, V. I. J. Mol. Biol. 1995, 252, 277-282.
- 43. Lim, V. I.; Venclovas, C. FEBS Lett. 1992, 313, 133-137.
- 44. Ishii, Y.; Kondo, S. Mutat. Res. 1975, 27, 27-44.
- 45. Tsien, R. Y. Annu. Rev. Biochem. 1998, 67, 509-544.
- 46. Stemmer, W. P. C.; Crameri, A.; Ha, K. D.; Brennan, T. M.; Heyneker, H. L. *Gene* **1995**, *164*, 49-53.
- 47. Fishman, R.; Ankilova, V.; Moor, N.; Safro, M. Acta Crystallogr. D. Biol. Crystallogr.
 2001, 57, 1534-44.
- 48. Sampson, J. R.; Behlen, L. S.; Direnzo, A. B.; Uhlenbeck, O. C. *Biochemistry* **1992**, *31*, 4161-4167.
- 49. Ghosh, I.; Hamilton, A. D.; Regan, L. J. Am. Chem. Soc. 2000, 122, 5658-5659.



Figure 1: LC-MS chromatogram of tryptic digests of mDHFR. Peptide **1** (residues 140-144; $LF_{UUU}VTR$) contains a Phe residue encoded as UUU. Peptide **2** (residues 63-70; TWF_{UUC}SIPEK) contains a Phe residue encoded as UUC codon. Peptide **1** variants containing Phe, Trp, and 2Nal were designated **1**F, **1**W, and **1**Z, respectively. Peptide **2** variants containing Phe and 2Nal were designated **2**F and **2**Z, respectively. These peptides were separated by LC and detected by MS. Unmodified mDHFR was synthesized in a Phe/Trp auxotrophic expression host (a and d) in 2xYT media. Modified mDHFRs were synthesized in a Phe/Trp auxotrophic expression host outfitted with ytRNA^{Phe}_{AAA} and yPheRS (T415G) (b and e) or yPheRS_naph (c and f). The minimal expression media were supplemented with 18 amino acids (25 µg/mL), 2.5 µM Phe, 3 mM Trp, and 3 mM 2Nal.



Figure 2: Phe substrate (red) and four residues (blue) within 7 Å of the *para*-position of the phenyl ring of the substrate inside the binding pocket of a homology model of yPheRS. These four residues were subjected to mutagenesis to generate a yPheRS library. The homology model of yPheRS was prepared by P. M. Kekenes-Huskey at Caltech.



Figure 3: A screening scheme for yPheRS library. GFP6 in yPheRS expression library *E. coli* cells outfitted with ytRNA^{Phe}_{AAA} and yPheRS library was expressed in the presence of 2Nal (a). Weakly fluorescent cells that contain active yPheRS variants were enriched by FACS (b). GFP6 in the collected cells was expressed in the absence of 2Nal (c). Highly fluorescent cells that contain selective yPheRS variants were enriched by FACS (d). After two rounds of screening, ten colonies were isolated from the enriched cells and characterized (e).



Figure 4: Fluorescence intensities of cells containing GFP6 expressed in minimal medium without 2Nal (a); with 2Nal (b). Visual comparison of fluorescence of cells containing GFP6 expressed in the absence of 2Nal (c left); in the presence of 2Nal (c right).



Figure 5: Fluorescence intensities of cells under uninduced conditions (a); cells expressing full length GFP6 (158Am) in the presence of 2Nal (b). The DHF *E. coli* strains were outfitted with $ytRNA^{Phe}_{CUA_UG}$ and $yPheRS_naph$. The expression media were supplemented with 18 amino acids (25 µg/mL), 50 µM Phe, 50 µM Trp, and 3 mM 2Nal.



Figure 6: UUC and UUU codon occupancy by Phe and 2Nal. Both GFP6 (2UUC) and GFP6 (2UUU) were expressed in DHF expression hosts outfitted with yPheRS_naph and ytRNA^{Phe}_{AAA} in minimal medium supplemented with 18 amino acids (25 μ g/mL), 50 μ M Trp, 3 mM 2Nal, and 0; 2.5 μ M; 5.0 μ M Phe. The UUC (a) and UUU (b) codon occupancy by Phe and 2Nal were determined by N-terminal sequencing.



Figure 7: Fluorescence intensities of cells expressing GFP variants. GFP3_WC was expressed in DHF expression hosts outfitted with yPheRS_naph and ytRNA^{Phe}_{AAA} in minimal medium supplemented with 18 amino acids, 5.0 μ M Phe, 50 μ M Trp, and no 2Nal (a); 3 mM 2Nal (b). GFP3 was expressed in MPC390 expression hosts outfitted with yPheRS_naph and ytRNA^{Phe}_{CAA} in minimal medium supplemented with 17 amino acids, 1.25 μ M Leu, 5.0 μ M Phe, 50 μ M Trp, and no 2Nal (c); 3 mM 2Nal (d).



184

Retention time (min)

Figure 8: LC-MS chromatogram of tryptic digests of mDHFR. Peptide 3 (residues 165-180; L_{CUU}L_{CUC}PEYPGVL_{CUC}SEVQEEK) contains three Leu residues encoded as CUU and CUC codons. Peptide 4 (residues 54-61; QNL_{CUG}VIMGR) contains a Leu residue encoded as CUG codon. Peptide 5 (residues 62-70; L_{CUU}IEQPEL_{UUG}ASK) contains two Leu residues encoded as CUU and UUG codons. Peptide 6 (residues 99-105; SL_{UUG}DDAL_{UUA}R) contains two Leu residues encoded as UUG and UUA codons. Peptide $\mathbf{6}_{\text{UUA}}$ is the same as Peptide $\mathbf{6}$ except both Leu residues are encoded as UUA codon. Peptide 3; 4; 5; 6; 6_{UUA} variants containing Leu and 2Nal were designated 3L and 3Z; 4L and 4Z; 5L and 5Z; 6L and 6Z; $6_{UUA}L$ and $6_{UUA}Z$, respectively. These peptides were separated by LC and detected by MS. Unmodified mDHFR was synthesized in the absence of 2Nal in a Phe/Leu auxotrophic expression host (a, c, e, and g) in 2xYT media. Modified mDHFRs were synthesized in a Phe/Leu auxotrophic expression host outfitted with ytRNA^{Phe}_{CAA} and yPheRS_naph (b, d, f, h and i). The expression minimal media were supplemented with 17 amino acids (25 µg/mL), 1.25 µM Leu, 50 µM Phe, 50 µM Trp, and 3 mM 2Nal. No 3Z, 4Z, or $6_{UUA}Z$ was detected by LC-MS analysis.



Figure 9: Fluorescence intensities of cells containing GFP3 (158UUG) expressed in minimal medium supplemented with 17 amino acids (25 μ g/mL), 1.25 μ M Leu, 5.0 μ M Phe, and 50 μ M Trp without 2Nal (a); with 3 mM 2Nal (b). MPC390 expression hosts were outfitted with yPheRS_naph and ytRNA^{Phe}_{CAA}.

 Table 1:
 Kinetic parameters for ATP-PPi exchange by yPheRS (T415G) and yPheRS_naph

$\begin{array}{c ccccccccccccccccccccccccccccccccccc$						
PheT415G 55 ± 14 0.202 ± 0.11 $3,500 \pm 1,100$ 1^{a} TrpT415G 2.83 ± 1.6 0.153 ± 0.003 $63,200 \pm 34,600$ 18^{a} 2NalT415G 7.03 ± 0.14 0.208 ± 0.04 $29,500 \pm 5,800$ 8.4^{a} Phenaph $11,000 \pm 2,700$ 0.0095 ± 0.0021 0.855 ± 0.007 1^{b} Trpnaph $1,400 \pm 600$ 0.0035 ± 0.0009 2.52 ± 0.44 2.9^{b} 2Nalnaph $2,000 \pm 700$ 0.030 ± 0.018 14.54 ± 4.22 17^{b}	Amino Acid	Enzyme	$K_{m}\left(\mu M\right)$	k_{cat} (s ⁻¹)	$k_{cat}/K_m (M^{-1}s^{-1})$	k_{cat}/K_{m} (rel)
TrpT415G 2.83 ± 1.6 0.153 ± 0.003 $63,200 \pm 34,600$ 18^a 2NalT415G 7.03 ± 0.14 0.208 ± 0.04 $29,500 \pm 5,800$ 8.4^a Phenaph $11,000 \pm 2,700$ 0.0095 ± 0.0021 0.855 ± 0.007 1^b Trpnaph $1,400 \pm 600$ 0.0035 ± 0.0009 2.52 ± 0.44 2.9^b 2Nalnaph $2,000 \pm 700$ 0.030 ± 0.018 14.54 ± 4.22 17^b	Phe	T415G	55 ± 14	0.202 ± 0.11	$3,500 \pm 1,100$	1 ^a
2NalT415G 7.03 ± 0.14 0.208 ± 0.04 $29,500 \pm 5,800$ 8.4^{a} Phenaph $11,000 \pm 2,700$ 0.0095 ± 0.0021 0.855 ± 0.007 1^{b} Trpnaph $1,400 \pm 600$ 0.0035 ± 0.0009 2.52 ± 0.44 2.9^{b} 2Nalnaph $2,000 \pm 700$ 0.030 ± 0.018 14.54 ± 4.22 17^{b}	Trp	T415G	2.83 ± 1.6	0.153 ± 0.003	$63,200 \pm 34,600$	18 ^a
Phenaph $11,000 \pm 2,700$ 0.0095 ± 0.0021 0.855 ± 0.007 1^b Trpnaph $1,400 \pm 600$ 0.0035 ± 0.0009 2.52 ± 0.44 2.9^b 2Nalnaph $2,000 \pm 700$ 0.030 ± 0.018 14.54 ± 4.22 17^b	2Nal	T415G	7.03 ± 0.14	0.208 ± 0.04	$29{,}500\pm5{,}800$	8.4 ^a
Trpnaph $1,400 \pm 600$ 0.0035 ± 0.0009 2.52 ± 0.44 2.9^{b} 2Nalnaph $2,000 \pm 700$ 0.030 ± 0.018 14.54 ± 4.22 17^{b}	Phe	naph	$11,000 \pm 2,700$	0.0095 ± 0.0021	0.855 ± 0.007	1 ^b
2Nal naph $2,000 \pm 700$ 0.030 ± 0.018 14.54 ± 4.22 17^{b}	Trp	naph	$1{,}400\pm600$	0.0035 ± 0.0009	2.52 ± 0.44	2.9^b
	2Nal	naph	$2,\!000\pm700$	0.030 ± 0.018	14.54 ± 4.22	17 ^b

^a Relative to k_{cat}/K_m for Phe by yPheRS (T415G).

^b Relative to k_{cat}/K_m for Phe by yPheRS_naph.

Codon	ytRNA	Occupancy of codon (%) ^b		
		2Nal	Phe	Lys
UUU ^a	ytRNA ^{Phe} AAA	80	20	ND ^c
UUU	ytRNA ^{Phe} GAA	92	8	ND
UAG ^d	ytRNA ^{Phe} CUA_UG	98	ND	2

Table 2: Occupancy of UUU and UAG codons by various amino acids

^a The second position in the amino acid sequence of GFP6.

^b UUU and UAG codon occupancy was determined by N-terminal protein sequencing and

LC-MS analysis, respectively.

^c Not detected.

^d The 38th position in the amino acid sequence of mDHFR_38Am.

Appendix A

Plasmid Information

190

Following are sequences and information on plasmids described in the thesis.

1. pQE16_mDHFR_yPheRS (T415G) (7765 bp)



T5 promoter and lac operon	7 -	- 87 bp
----------------------------	-----	---------

mDHFR_His	115 -	660 bp
-----------	-------	--------

- yPheRS_alpha subunit 1226 2787 bp
- yPheRS_beta subunit 2802 4589 bp
- ColE1 origin 5972 5997 bp
- beta-lactamase 7560 6700 bp

Full sequence

CTC GAG AAA TCA TAA AAA ATT TAT TTG CTT TGT GAG CGG ATA ACA ATT ATA ATA GAT TCA ATT GTG AGC GGA TAA CAA TTT CAC ACA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG AGA GGA TCC GGC ATC ATG GTT CGA CCA TTG AAC GCC TCG ATC GTC GTG TCC CAA AAT ATG GGG ATT GGC AAG AAC GGA GAC CTA CCC TGG CCT CTC CCG AGG AAC GAG TTC AAG TAC TTC CAA AGA ATG ACC ACA ACC TCT TCA GTG GAA GGT AAA CAG AAT CTG GTG ATT ATG GGT AGG AAA ACC TGG TTC TCC ATT CCT GAG AAG AAT CGA CCT TTA AAG GAC AGA ATT AAT ATA GTT CTC AGT AGA GAA CTC AAA GAA

CCA CCA CGA GGA GCT CAT TTT CTT GCC AAA AGT TTG GAT GAT GCC TTA AGA CTT ATT GAA CAA CCG GAA TTG GCA AGT AAA GTA GAC ATG GTT TGG ATA GTC GGA GGC AGT TCT GTT TAC CAG GAA GCC ATG AAT CAA CCA GGC CAC CTT AGA CTC TTT GTG ACA AGG ATC ATG CAG GAA TTT GAA AGT GAC ACG TTT TTC CCA GAA ATT GAT TTG GGG AAA TAT AAA CTT CTC CCA GAA TAC CCA GGC GTC CTC TCT GAG GTC CAG GAG GAA AAA GGC ATC AAG TAT AAG TTT GAA GTC TAC GAG AAG AAA GGT TCC AGA TCT CAT CAC CAT CAC CAT CAC TAA GCT TAA TTA GCT GAG CTT GGA CTC CTG TTG ATA GAT CCA GTA ATG ACC TCA GAA CTC CAT CTG GAT TTG TTC AGA ACG CTC GGT TGC CGC CGG GCG TTT TTT ATT GGT GAG AAT CCA AGC TAG CTT GGC GAG ATT TTC AGG AGC TAA GGA AGC TAA AAT GGA GAA AAA AAT CAC TGG ATA TAC CAC CGT TGA TAT ATC CCA ATG GCA TCG TAA AGA ACA TTT TGA GGC ATT TCA GTC AGT TGC TCA ATG TAC CTA TAA CCA GAC CGT TCA GCT GGC ACG ACA GGT TTC CCG ACT GGA AAG CGG GCA GTG AGC GCA ACG CAA TTA ATG TGA GTT AGC TCA CTC ATT AGG CAC CCC AGG CTT TAC ACT TTA TGC TTC CGG CTC GTA TGT TGT GTG GAA TTG TGA GCG GAT AAC AAT TTC ACA CAG GAA ACA GCT ATG ACC ATG ATT ACG CCA AGC TTG CAT GCC TGC AGT TGA CAA TTA ATC ATC GGC TCG TAT AAT GGA TCC AAT TGT GAG CGG AAT CGA TTT TCA CAC AGG AAA CAG ACC ATG AAT CTA GAG ATG TCT GAC TTC CAA TTA GAA ATT CTA AAG AAA CTA GAT GAA TTG GAT GAG ATC AAG TCC ACA CTG GCA ACT TTC CCT CAG CAC GGC TCT CAA GAT GTT CTT TCC GCT TTG AAC TCT TTG AAA GCC CAC AAC AAG TTA GAG TTT TCC AAG GTC GAC ACG GTT ACG TAT GAC TTG ACC AAA GAA GGT GCT CAA ATT TTG AAT GAA GGT TCG TAC GAA ATT AAA CTA GTC AAG CTC ATC CAA GAG TTG GGT CAA CTT CAA ATC AAA GAT GTG ATG TCC AAA CTA GGC CCT CAA GTT GGT AAG GTC GGT CAG GCT AGA GCT TTC AAG AAC GGC TGG ATC GCC AAA AAC GCC TCA AAC GAG CTT GAA CTC TCC GCA AAA TTG CAA AAT ACC GAT TTA AAT GAG CTT ACT GAT GAA ACG CAA TCT ATT CTA GCG CAA ATC AAG AAC ACC TCG CAT CTG GAT AGC ATT GAC GCC AAG ATT TTG AAC GAC TTG AAG AAA AGA AAG TTA ATT GCT CAA GGT AAA ATC ACA GAT TTC AGT GTC ACC AAA GGG CCA GAG TTC TCG ACC GAC CTC ACC AAA TTG GAA ACC GAT CTT ACC TCC GAC ATG GTC TCC ACC AAT GCA TAC AAG GAC TTG AAG TTC AAG CCT TAC AAT TTC AAT TCT CAA GGT GTG CAA ATA TCT TCA GGT GCT CTT CAC CCC TTA AAC AAA GTC AGA GAG GAA TTT AGA CAA ATT TTC TTT TCC ATG GGA TTC ACA GAG ATG CCC TCG AAC CAA TAC GTC GAG ACA GGT TTC TGG AAC TTC GAT GCC CTT TAC GTC CCA CAA CAG CAT CCT GCT CGT GAC CTG CAA GAC ACT TTC TAC ATC AAG GAC CCA CTA ACC GCT GAG TTG CCC GAT GAC AAG ACA TAC ATG GAC AAT ATC AAA GCC GTT CAC GAA CAG GGG AGA TTC GGG TCC ATC GGT TAT CGT TAC AAC TGG AAG CCA GAA GAA TGT CAA AAA TTG GTC TTG AGA ACT CAC TCC ACA GCC ATC TCT GCC AGA ATG CTG CAC GAT TTG GCC AAA GAT CCA AAG CCC ACC AGA TTG TTT TCT ATC GAC CGT GTT TTC CGT AAC GAA GCA GTT GAC GCC ACC CAT TTG GCC GAA TTC CAC CAG GTG GAA GGT GTT CTT GCC GAC TAC AAC ATT ACT CTG GGT GAC CTG ATC AAG TTC ATG GAA GAG TTT TTC GAA AGA ATG GGT GTC ACC GGT TTG AGA TTC AAG CCT ACC TAC AAT CCT TAC GGC GAG CCA TCA ATG GAA ATC TTT TCT TGG CAC GAA GGT TTG CAA AAA TGG GTC GAA ATC GGT AAC TCT GGT ATG TTC AGA CCA GAA ATG CTC GAG TCC ATG GGT CTA CCA AAG GAT CTA AGA GTC CTT GGT TGG GGG TTA TCC TTG GAA AGA CCT ACC ATG ATC AAA TAT AAG GTT CAA AAC ATC AGA GAA CTG TTA GGT CAT AAA GTC TCT TTG GAC TTT ATC GAA ACC AAT CCT GCT GCT AGA TTG GAC GAA GAC TTG TAC GAA TAA GGC AGG AAT AGA TTA TGC CTA CCG TCT CCG TGA ACA AGC AGC AAT TAT TTG ATC TTC TAG GCA AAG ACT ACA CTT CCC AAG AGT TCG ACG AAT TAT GTT TTG AAT TCG GTA TGG AAA TGG ACG AAG ACA CCA CAG AAG AGG CCT TGA AAA CCG GGG AGG AGC CGG AAT TGA AGC TTG ATA TCA GTG CCA ATC GTT ACG ATT TGC TTT GTA TCG AAG GTA TTT CAC AGT CGC TGA ACG AAT ACT TGG AAC GTA AAG AAA GAC CTG ACT ATA AAT TAA GCA AGC CAA CCA CTA AGT TGA TCA TCG ACA AAT CAA CGG AGC AAA TTA GAC CTT TTG CTA CCG CTG CTG TAT TGA GAA ATA TCA AGC TTA ACG AAA AAT CTT ACG CTT CTT TTA TTG CCT TGC AAG ATA AAT TAC ATG CCA ATC TAT GTA GAA ACA GAA GCT TGG TTG CCA TGG GTA CTC ACG ATT TAG ATT CAA TTG AAG GTC CAT TCC ATT ACA GAG CTC TAC CAC CAA AGG ACA TCA AGT TCG TAC CAT TGA ATC AAA CCC AAG AGT TTA CTG GTG ACA AAT TGA TCG AGT TTT ATA AAT CTC CAG AAC AGA AAA ACA ACA TAG GGA GAT ACG TTC ACA TTA TTG AGG ATT CTC CAG TCT TCC CAG TTA TTA TGG ACA GCA AAG ATC GTG

TTT GCT CCC TGC CAC CAT TAA TCA ATA GTG AAC ATT CGA AGA TCT CTG TGA ACA CCC GTA ACA TTT TGA TTG ATA TAA CCG CCA CCG ATA AGA CCA AAG CCG AGA TCG TTT TGA ACA TAT TAA CTA CAA TGT TCT CAC GTT ATT GTG ACG AAC CAT TCA CGG TTG AGC CTG TAG AAA TTG TCT CTG AAC ACA ATG GCC AAT CCC GTT TGG CGC CAA ACT TCA ACG ATA GAA TTA TGG ATG TCT CCA TTA AGT ACA TCA ACT CCT GTC TTG GCC TAG ATC AAT CCG CTG ATG AAA TTG CTC ATT GTC TAA AAA AGA TGT CGT TGC ATG CCG TTC AAT CAA AGG AAG ACA AGG ACA TCT TGC ACG TTG ACA TTC CGG TAA CTA GAC CTG ATA TTT TGC ACG CTT GTG ATA TAA TGG AAG ATG CCG CTG TCG GTT ATG GTT TCA ATA ATT TGC CAA AGG GTG AGA AAT TAT CCA ATG CCA ACT TCA TTG CCA AAC CAT TAC CAA TCA ACA AGG TTT CTG ATA TTT TCA GAG TTG CAT CCT CTC AAG CCA CGT GGG TTG AGG TTT TAC CAT TGA CCT TAT GTT CGC ACG ATG AAA ACT TTA AAT TTC TAA GAC AAT CCG ACA ATG ATG ATT TAG CTG TCA AAT TGG CCA ACC CAA AGA CTT TGG AAT ACC AAG TTG TTA GAA CCA CTT TAT TGC CTG GTA TCT TAA AGA CAG TCA AGG AAA ACA GAA AAC ATT CCT TAC CAA TCA AAG TCT TTG AAA CCG GTG ACG TTG TAT TTA AAG ACG ACA AAC TAG AAA GGA AGG CGT ACA ATG AAC GTC ACT GGG CTG CCA TCT ACG TGG GTA AGA ATT CTG GGT TTG AAA TCA TTC AAG GGT TAT TGG GTA AAA TCA TGC AAA CTT TTA GAA CAG AGT GGA TTG CAG ACT ACG GTG CTG CTG CTT CTG GCA GAG GTT ACT GGA TTG AAG AAG ACG ATT CTG TGA AAA CCT ATT TCC CAG GTA GAG GTG CCA AGG TCA TGT TCA GAT CCA AAG AAG GCG CTG AGC CAA AGC AAA TCG GCC ACT TGG GTG TCT TGC ATC CTG AAG TCA TGA TGA ATT TCG ACG TTC CAT TCG CTG CAT CCT TTG TAG AGG TTA ATG CCG AAG TCT TCC TAT AAT GTA ATG TTC TAA CAA AAA TTT TTA CTG ATT TAT AAA ACT TAT ATA GAT AGA TAG ACA TAT ATA TAT GGG TAC CGA GCT CGA ATT CAC TGG CCG TCG TTT TAC AAC GTC GTG ACT GGG AAA ACC CTG GCG TTA CCC AAC TTA ATC GCC TTG CAG CAC ATC CCC CTT TCG CCA GCT GGA TAT TAC GGC CTT TTT AAA GAC CGT AAA GAA AAA TAA GCA CAA GTT TTA TCC GGC CTT TAT TCA CAT TCT TGC CCG CCT GAT GAA TGC TCA TCC GGA ATT TCG TAT GGC AAT GAA AGA CGG TGA GCT GGT GAT ATG GGA TAG TGT TCA CCC TTG TTA CAC CGT TTT CCA TGA GCA AAC TGA AAC GTT TTC ATC GCT CTG GAG TGA ATA CCA CGA CGA TTT CCG GCA GTT TCT ACA CAT ATA TTC GCA AGA TGT GGC GTG TTA CGG TGA AAA CCT GGC CTA TTT CCC TAA AGG GTT TAT TGA GAA TAT GTT TTT CGT CTC AGC CAA TCC CTG GGT GAG TTT CAC CAG TTT TGA TTT AAA CGT GGC CAA TAT GGA CAA CTT CTT CGC CCC CGT TTT CAC CAT GGG CAA ATA TTA TAC GCA AGG CGA CAA GGT GCT GAT GCC GCT GGC GAT TCA GGT TCA TCA TGC CGT CTG TGA TGG CTT CCA TGT CGG CAG AAT GCT TAA TGA ATT ACA ACA GTA CTG CGA TGA GTG GCA GGG CGG GGC GTA ATT TTT TTA AGG CAG TTA TTG GTG CCC TTA AAC GCC TGG GGT AAT GAC TCT CTA GCT TGA GGC ATC AAA TAA AAC GAA AGG CTC AGT CGA AAG ACT GGG CCT TTC GTT TTA TCT GTT GTT TGT CGG TGA ACG CTC TCC TGA GTA GGA CAA ATC CGC CGC TCT AGA GCT GCC TCG CGC GTT TCG GTG ATG ACG GTG AAA ACC TCT GAC ACA TGC AGC TCC CGG AGA CGG TCA CAG CTT GTC TGT AAG CGG ATG CCG GGA GCA GAC AAG CCC GTC AGG GCG CGT CAG CGG GTG TTG GCG GGT GTC GGG GCG CAG CCA TGA CCC AGT CAC GTA GCG ATA GCG GAG TGT ATA CTG GCT TAA CTA TGC GGC ATC AGA GCA GAT TGT ACT GAG AGT GCA CCA TAT GCG GTG TGA AAT ACC GCA CAG ATG CGT AAG GAG AAA ATA CCG CAT CAG GCG CTC TTC CGC TTC CTC GCT CAC GGC GGT AAT ACG GTT ATC CAC AGA ATC AGG GGA TAA CGC AGG AAA GAA CAT GTG AGC AAA AGG CCA GCA AAA GGC CAG GAA CCG TAA AAA GGC CGC GTT GCT GGC GTT TTT CCA TAG GCT CCG CCC CCC TGA CGA GCA TCA CAA AAA TCG ACG CTC AAG TCA GAG GTG GCG AAA CCC GAC AGG ACT ATA AAG ATA CCA GGC GTT TCC CCC TGG AAG CTC CCT CGT GCG CTC TCC TGT TCC GAC CCT GCC GCT TAC CGG ATA CCT GTC CGC CTT TCT CCC TTC GGG AAG CGT GGC GCT TTC TCA ATG CTC ACG CTG TAG GTA TCT CAG TTC GGT GTA GGT CGT TCG CTC CAA GCT GGG CTG TGT GCA CGA ACC CCC CGT TCA GCC CGA CCG CTG CGC CTT ATC CGG TAA CTA TCG TCT TGA GTC CAA CCC GGT AAG ACA CGA CTT ATC GCC ACT GGC AGC AGC CAC TGG TAA CAG GAT TAG CAG AGC GAG GTA TGT AGG CGG TGC TAC AGA GTT CTT GAA GTG GTG GCC TAA CTA CGG CTA CAC TAG AAG GAC AGT ATT TGG TAT CTG CGC TCT GCT GAA GCC AGT TAC CTT CGG AAA AAG AGT TGG TAG CTC TTG ATC CGG CAA ACA AAC CAC CGC TGG TAG CGG TGG TTT TTT TGT TTG CAA GCA GCA GAT TAC GCG CAG AAA AAA AGG ATC TCA AGA AGA

TCC TTT GAT CTT TTC TAC GGG GTC TGA CGC TCA GTG GAA CGA AAA CTC ACG TTA AGG GAT TTT GGT CAT GAG ATT ATC AAA AAG GAT CTT CAC CTA GAT CCT TTT AAA TTA AAA ATG AAG TTT TAA ATC AAT CTA AAG TAT ATA TGA GTA AAC TTG GTC TGA CAG TTA CCA ATG CTT AAT CAG TGA GGC ACC TAT CTC AGC GAT CTG TCT ATT TCG TTC ATC CAT AGC TGC CTG ACT CCC CGT CGT GTA GAT AAC TAC GAT ACG GGA GGG CTT ACC ATC TGG CCC CAG TGC TGC AAT GAT ACC GCG AGA CCC ACG CTC ACC GGC TCC AGA TTT ATC AGC AAT AAA CCA GCC AGC CGG AAG GGC CGA GCG CAG AAG TGG TCC TGC AAC TTT ATC CGC CTC CAT CCA GTC TAT TAA TTG TTG CCG GGA AGC TAG AGT AAG TAG TTC GCC AGT TAA TAG TTT GCG CAA CGT TGT TGC CAT TGC TAC AGG CAT CGT GGT GTC ACG CTC GTC GTT TGG TAT GGC TTC ATT CAG CTC CGG TTC CCA ACG ATC AAG GCG AGT TAC ATG ATC CCC CAT GTT GTG CAA AAA AGC GGT TAG CTC CTT CGG TCC TCC GAT CGT TGT CAG AAG TAA GTT GGC CGC AGT GTT ATC ACT CAT GGT TAT GGC AGC ACT GCA TAA TTC TCT TAC TGT CAT GCC ATC CGT AAG ATG CTT TTC TGT GAC TGG TGA GTA CTC AAC CAA GTC ATT CTG AGA ATA GTG TAT GCG GCG ACC GAG TTG CTC TTG CCC GGC GTC AAT ACG GGA TAA TAC CGC GCC ACA TAG CAG AAC TTT AAA AGT GCT CAT CAT TGG AAA ACG TTC TTC GGG GCG AAA ACT CTC AAG GAT CTT ACC GCT GTT GAG ATC CAG TTC GAT GTA ACC CAC TCG TGC ACC CAA CTG ATC TTC AGC ATC TTT TAC TTT CAC CAG CGT TTC TGG GTG AGC AAA AAC AGG AAG GCA AAA TGC CGC AAA AAA GGG AAT AAG GGC GAC ACG GAA ATG TTG AAT ACT CAT ACT CTT CCT TTT TCA ATA TTA TTG AAG CAT TTA TCA GGG TTA TTG TCT CAT GAG CGG ATA CAT ATT TGA ATG TAT TTA GAA AAA TAA ACA AAT AGG GGT TCC GCG CAC ATT TCC CCG AAA AGT GCC ACC TGA CGT CTA AGA AAC CAT TAT TAT CAT GAC ATT AAC CTA TAA AAA TAG GCG TAT CAC GAG GCC CTT TCG TCT TCA C

2. pQE9_GFP6_lacI_yPheRS_naph (8,294 bp)



T5 promoter and lac operon	7 -	87 bp
His_GFP6	115 -	921 bp

yPheRS_alpha subunit	1449 -	2972 bp
----------------------	--------	---------

vPheRS beta subunit	2987 - 4774 bp
yr netto_beta subunit	2707 +77+0p

ColE1 origin	6501 -	6526 bp

beta-lactamase 8089 - 7229 bp

Full sequence

CTC GAG AAA TCA TAA AAA ATT TAT TTG CTT TGT GAG CGG ATA ACA ATT ATA ATA GAT TCA ATT GTG AGC GGA TAA CAA TTT CAC ACA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG AGA GGA TCG CAT CAC CAT CAC CAT CAC GGA TCC GTC GAC CTG CAG CCC CGG GTA CCG GTA GAA AAA ATG AGT AAA GGA GAA GAA CTT TTTACT GGA GTT GTC CCA ATT CTT GTT GAA TTA GAT GGT GAT GTT AAT GGG CAC AAA TTTTCT GTC AGT GGA GAG GGT GAA GGT GAT GCA ACA TAC GGA AAA CTT ACC CTT AAA TTT ATT TGC ACT ACT GGA AAA CTA CCT GTT CCA TGG CCA ACA CTT GTC ACT ACT TTT ACC TAT GGT GTT CAA TGC TTT TCC CGT TAT CCG GAT CAT ATG AAA CGG CAT GAC TTT TTT AAG AGT GCC ATG CCC GAA GGT TAT GTA CAG GAA CGC ACT ATA TTT TTT AAA GAT GAC GGG AAC TAC AAG ACG CGT GCT GAA GTC AAG TTT GAA GGT GAT ACC CTT GTT AAT CGT ATC GAG TTA AAA GGT ATT GAT TTT AAA GAA GAT GGA AAC ATT CTC GGA CAC AAA CTC GAG TAC AAC TAT AAC TCA CAC AAT GTA TAC ATC ATG GCA GAC

194

AAA CAA AAG AAT GGA ATC AAA GCT AAC TTT AAA ATT CGC CAC AAC ATT GAA GAT GGA TCC GTT CAA CTA GCA GAC CAT TAT CAA CAA AAT ACT CCA ATT GGC GAT GGC CCT GTC CTT TTA CCA GAC AAC CAT TAC CTG TCG ACA CAA TCT GCC CTT TCG AAA GAT CCC AAC GAA AAG CGT GAC CAC ATG GTC CTT CTT GAG TTT GTA ACT GCT GCT GGG ATT ACA CAT GGC ATG GAT GAG CTC TAC AAA CTG CAG CCA AGC TTA ATT AGC TGA AGC TTG GAC TCC TGT TGA TAG ATC CAG TAA TGA CCT CAG AAC TCC ATC TGG ATT TGT TCA GAA CGC TCG GTT GCC GCC GGG CGT TTT TTA TTG GTG AGA ATC CAA GCT AGC TTG GCG AGA TTT TCA GGA GCT AAG GAA GCT AAA ATG GAG AAA AAA ATC ACT GGA TAT ACC ACC GTT GAT ATA TCC CAA TGG CAT CGT AAA GAA CAT TTT GAG GCA TTT CAG TCA GTT GCT CAA TGT ACC TAT AAC CAG ACC GTT CAG CTG GCA CGA CAG GTT TCC CGA CTG GAA AGC GGG CAG TGA GCG CAA CGC AAT TAA TGT GAG TTA GCT CAC TCA TTA GGC ACC CCA GGC TTT ACA CTT TAT GCT TCC GGC TCG TAT GTT GTG TGG AAT TGT GAG CGG ATA ACA ATT TCA CAC AGG AAA CAG CTA TGA CCA TGA TTA CGC CAA GCT TGC ATG CCT GCA GTT GAC AAT TAA TCA TCG GCT CGT ATA ATG GAT CCA ATT GTG AGC GGA ATC GAT TTT CAC ACA GGA AAC AGA CCA TGA ATC TAG AGA TGT CTG ACT TCC AAT TAG AAA TTC TAA AGA AAC TAG ATG AAT TGG ATG AGA TCA AGT CCA CAC TGG CAA CTT TCC CTC AGC ACG GCT CTC AAG ATG TTC TTT CCG CTT TGA ACT CTT TGA AAG CCC ACA ACA AGT TAG AGT TTT CCA AGG TCG ACA CGG TTA CGT ATG ACT TGA CCA AAG AAG GTG CTC AAA TTT TGA ATG AAG GTT CGT ACG AAA TTA AAC TAG TCA AGC TCA TCC AAG AGT TGG GTC AAC TTC AAA TCA AAG ATG TGA TGT CCA AAC TAG GCC CTC AAG TTG GTA AGG TCG GTC AGG CTA GAG CTT TCA AGA ACG GCT GGA TCG CCA AAA ACG CCT CAA ACG AGC TTG AAC TCT CCG CAA AAT TGC AAA ATA CCG ATT TAA ATG AGC TTA CTG ATG AAA CGC AAT CTA TTC TAG CGC AAA TCA AGA ACA ACT CGC ATC TGG ATA GCA TTG ACG CCA AGA TTT TGA ACG ACT TGA AGA AAA GAA AGT TAA TTG CTC AAG GTA AAA TCA CAG ATT TCA GTG TCA CCA AAG GGC CAG AGT TCT CGA CCG ACC TCA CCA AAT TGG AAA CCG ATC TTA CCT CCG ACA TGG TCT CCA CCA ATG CAT ACA AGG ACT TGA AGT TCA AGC CTT ACA ATT TCA ATT CTC AAG GTG TGC AAA TAT CTT CAG GTG CTC TTC ACC CCT TAA ACA AAG TCA GAG AGG AAT TTA GAC AAA TTT TCT TTT CCA TGG GAT TCA CAG AGA TGC CCT CGA ACC AAT ACG TCG AGA CAG GTT TCT GGA ACT TCG ATG CCC TTT ACG TCC CAC AAC AGC ATC CTG CTC GTG ACC TGC AAG ACA CTT TCT ACA TCA AGG ACC CAC TAA CCG CTG AGT TGC CCG ATG ACA AGA CAT ACA TGG ACA ATA TCA AAG CCG TTC ACG AAC AGG GGA GAT TCG GGT CCA TCG GTT ATC GTT ACA ACT GGA AGC CAG AAG AAT GTC AAA AAT TGG TCT TGA GAA CTC ACT CCA CAG CCA TCT CTG CCA GAA TGC TGC ACG ATT TGG CCA AAG ATC CAA AGC CCA CCA GAT TGT TTT CTA TCG ACC GTG TTT TCC GTA ACG AAG CAG TTG ACG CCA CCC ATT TGG CCG AAT TCC ACC AGG TGG AAG GTG TTC TTG CCG ACT ACA ACA TTA CTC TGG GTG ACC TGA TCA AGT TCA TGG AAG AGT TTT TCG AAA GAA TGG GTG TCA CCG GTT TGA GAT TCA AGC CTA CCT ACG GGC CTT ACG GGG AGC CAT CTA TGG AAA TCT TTT CTT GGC ACG AAG GTT TGC AAA AAT GGG TCG AAA TCG GTA ACT TTG GTA TGT TCA GAC CAG AAA TGC TCG AGT CCA TGG GTC TAC CAA AGG ATC TAA GAG TCC TTG GTT GGG GGT TAT CCT TGG AAA GAC CTA CCA TGA TCA AAT ATA AGG TTC AAA ACA TCA GAG AAC TGT TAG GTC ATA AAG TCT CTT TGG ACT TTA TCG AAA CCA ATC CTG CTG CTA GAT TGG ACG AAG ACT TGT ACG AAT AAG GCA GGA ATA GAT TAT GCC TAC CGT CTC CGT GAA CAA GCA GCA ATT ATT TGA TCT TCT AGG CAA AGA CTA CAC TTC CCA AGA GTT CGA CGA ATT ATG TTT TGA ATT CGG TAT GGA AAT GGA CGA AGA CAC CAC AGA AGA GGC CTT GAA AAC CGG GGA GGA GCC GGA ATT GAA GCT TGA TAT CAG TGC CAA TCG TTA CGA TTT GCT TTG TAT CGA AGG TAT TTC ACA GTC GCT GAA CGA ATA CTT GGA ACG TAA AGA AAG ACC TGA CTA TAA ATT AAG CAA GCC AAC CAC TAA GTT GAT CAT CGA CAA ATC AAC GGA GCA AAT TAG ACC TTT TGC TAC CGC TGC TGT ATT GAG AAA TAT CAA GCT TAA CGA AAA ATC TTA CGC TTC TTT TAT TGC CTT GCA AGA TAA ATT ACA TGC CAA TCT ATG TAG AAA CAG AAG CTT GGT TGC CAT GGG TAC TCA CGA TTT AGA TTC AAT TGA AGG TCC ATT CCA TTA CAG AGC TCT ACC ACC AAA GGA CAT CAA GTT CGT ACC ATT GAA TCA AAC CCA AGA GTT TAC TGG TGA CAA ATT GAT CGA GTT TTA TAA ATC TCC AGA ACA GAA AAA CAA CAT AGG GAG ATA CGT TCA CAT TAT TGA GGA TTC TCC AGT CTT CCC AGT TAT TAT GGA CAG CAA AGA TCG TGT TTG CTC CCT GCC ACC ATT AAT CAA TAG TGA ACA TTC GAA GAT CTC TGT GAA CAC CCG TAA CAT TTT GAT TGA TAT AAC CGC CAC

CGA TAA GAC CAA AGC CGA GAT CGT TTT GAA CAT ATT AAC TAC AAT GTT CTC ACG TTA TTG TGA CGA ACC ATT CAC GGT TGA GCC TGT AGA AAT TGT CTC TGA ACA CAA TGG CCA ATC CCG TTT GGC GCC AAA CTT CAA CGA TAG AAT TAT GGA TGT CTC CAT TAA GTA CAT CAA CTC CTG TCT TGG CCT AGA TCA ATC CGC TGA TGA AAT TGC TCA TTG TCT AAA AAA GAT GTC GTT GCA TGC CGT TCA ATC AAA GGA AGA CAA GGA CAT CTT GCA CGT TGA CAT TCC GGT AAC TAG ACC TGA TAT TTT GCA CGC TTG TGA TAT AAT GGA AGA TGC CGC TGT CGG TTA TGG TTT CAA TAA TTT GCC AAA GGG TGA GAA ATT ATC CAA TGC CAA CTT CAT TGC CAA ACC ATT ACC AAT CAA CAA GGT TTC TGA TAT TTT CAG AGT TGC ATC CTC TCA AGC CAC GTG GGT TGA GGT TTT ACC ATT GAC CTT ATG TTC GCA CGA TGA AAA CTT TAA ATT TCT AAG ACA ATC CGA CAA TGA TGA TTT AGC TGT CAA ATT GGC CAA CCC AAA GAC TTT GGA ATA CCA AGT TGT TAG AAC CAC TTT ATT GCC TGG TAT CTT AAA GAC AGT CAA GGA AAA CAG AAA ACA TTC CTT ACC AAT CAA AGT CTT TGA AAC CGG TGA CGT TGT ATT TAA AGA CGA CAA ACT AGA AAG GAA GGC GTA CAA TGA ACG TCA CTG GGC TGC CAT CTA CGT GGG TAA GAA TTC TGG GTT TGA AAT CAT TCA AGG GTT ATT GGG TAA AAT CAT GCA AAC TTT TAG AAC AGA GTG GAT TGC AGA CTA CGG TGC TGC TGC TTC TGG CAG AGG TTA CTG GAT TGA AGA AGA CGA TTC TGT GAA AAC CTA TTT CCC AGG TAG AGG TGC CAA GGT CAT GTT CAG ATC CAA AGA AGG CGC TGA GCC AAA GCA AAT CGG CCA CTT GGG TGT CTT GCA TCC TGA AGT CAT GAT GAA TTT CGA CGT TCC ATT CGC TGC ATC CTT TGT AGA GGT TAA TGC CGA AGT CTT CCT ATA ATG TAA TGT TCT AAC AAA AAT TTT TAC TGA TTT ATA AAA CTT ATA TAG ATA GAT AGA CAT ATA TAT ATA TCT ATA TAT AGA AAC ACA ACT AAA GTT TAC CAT GTT TTA TAT AGG GTA CCG AGC TCG AAT TCA CTG GCC GTC GTT TTA CAA CGT CGT GAC TGG GAA AAC CGC GGT TAC GTG CAG TCG ATG ATA AGC TGT CAA ACA TGA GAA TTG TGC CTA ATG AGT GAG CTA ACT TAC ATT AAT TGC GTT GCG CTC ACT GCC CGC TTT CCA GTC GGG AAA CCT GTC GTG CCA GCT GCA TTA ATG AAT CGG CCA ACG CGC GGG GAG AGG CGG TTT GCG TAT TGG GCG CCA GGG TGG TTT TTC TTT TCA CCA GTG AGA CGG GCA ACA GCT GAT TGC CCT TCA CCG CCT GGC CCT GAG AGA GTT GCA GCA AGC GGT CCA CGC TGG TTT GCC CCA GCA GGC GAA AAT CCT GTT TGA TGG TGG TTA ACG GCG GGA TAT AAC ATG AGC TGT CTT CGG TAT CGT CGT ATC CCA CTA CCG AGA TAT CCG CAC CAA CGC GCA GCC CGG ACT CGG TAA TGG CGC GCA TTG CGC CCA GCG CCA TCT GAT CGT TGG CAA CCA GCA TCG CAG TGG GAA CGA TGC CCT CAT TCA GCA TTT GCA TGG TTT GTT GAA AAC CGG ACA TGG CAC TCC AGT CGC CTT CCC GTT CCG CTA TCG GCT GAA TTT GAT TGC GAG TGA GAT ATT TAT GCC AGC CAG CCA GAC GCA GAC GCG CCG AGA CAG AAC TTA ATG GGC CCG CTA ACA GCG CGA TTT GCT GGT GAC CCA ATG CGA CCA GAT GCT CCA CGC CCA GTC GCG TAC CGT CTT CAT GGG AGA AAA TAA TAC TGT TGA TGG GTG TCT GGT CAG AGA CAT CAA GAA ATA ACG CCG GAA CAT TAG TGC AGG CAG CTT CCA CAG CAA TGG CAT CCT GGT CAT CCA GCG GAT AGT TAA TGA TCA GCC CAC TGA CGC GTT GCG CGA GAA GAT TGT GCA CCG CCG CTT TAC AGG CTT CGA CGC CGC TTC GTT CTA CCA TCG ACA CCA CGC TGG CAC CCA GTT GAT CGG CGC GAG ATT TAA TCG CCG CGA CAA TTT GCG ACG GCG CGT GCA GGG CCA GAC TGG AGG TGG CAA CGC CAA TCA GCA ACG ACT GTT TGC CCG CCA GTT GTT GTG CCA CGC GGT TGG GAA TGT AAT TCA GCT CCG CCA TCG CCG CTT CCA CTT TTT CCC GCG TTT TCG CAG AAA CGT GGC TGG CCT GGT TCA CCA CGC GGG AAA CGG TCT GAT AAG AGA CAC CGG CAT ACT CTG CGA CAT CGT ATA ACG TTA CTG GTT TCA CAT TCA CCA CCC TGA ATT GAC TCT CTT CCG GGC GCT ATC ATG CCA TAC CGC GAA AGG TTT TGC ACC ATT CGA TGG TGT CGG AAT TTC GGG CAG CGT TGG GTC CTG GCC ACG GGT GCG CAT GAC TTA AGC GGT GTG AAA TAC CGC ACA GAT GCG TAA GGA GAA AAT ACC GCA TCA GGC GCT CTT CCG CTT CCT CGC TCA CTG ACT CGC TGC GCT CGG TCT GTC GGC TGC GGC GAG CGG TAT CAG CTC ACT CAA AGG CGG TAA TAC GGT TAT CCA CAG AAT CAG GGG ATA ACG CAG GAA AGA ACA TGT GAG CAA AAG GCC AGC AAA AGG CCA GGA ACC GTA AAA AGG CCG CGT TGC TGG CGT TTT TCC ATA GGC TCC GCC CCC CTG ACG AGC ATC ACA AAA ATC GAC GCT CAA GTC AGA GGT GGC GAA ACC CGA CAG GAC TAT AAA GAT ACC AGG CGT TTC CCC CTG GAA GCT CCC TCG TGC GCT CTC CTG TTC CGA CCC TGC CGC TTA CCG GAT ACC TGT CCG CCT TTC TCC CTT CGG GAA GCG TGG CGC TTT CTC AAT GCT CAC GCT GTA GGT ATC TCA GTT CGG TGT AGG TCG TTC GCT CCA AGC TGG GCT GTG TGC ACG AAC CCC CCG TTC AGC CCG ACC GCT GCG CCT TAT CCG GTA ACT ATC GTC TTG AGT CCA ACC CGG TAA GAC ACG ACT

TAT CGC CAC TGG CAG CAG CCA CTG GTA ACA GGA TTA GCA GAG CGA GGT ATG TAG GCG GTG CTA CAG AGT TCT TGA AGT GGT GGC CTA ACT ACG GCT ACA CTA GAA GGA CAG TAT TTG GTA TCT GCG CTC TGC TGA AGC CAG TTA CCT TCG GAA AAA GAG TTG GTA GCT CTT GAT CCG GCA AAC AAA CCA CCG CTG GTA GCG GTG GTT TTT TTG TTT GCA AGC AGA TTA CGC GCA GAA AAA AAG GAT CTC AAG AAG ATC CTT TGA TCT TTT CTA CGG GGT CTG ACG CTC AGT GGA ACG AAA ACT CAC GTT AAG GGA TTT TGG TCA TGA GAT TAT CAA AAA GGA TCT TCA CCT AGA TCC TTT TAA ATT AAA AAT GAA GTT TTA AAT CAA TCT AAA GTA TAT ATG AGT AAA CTT GGT CTG ACA GTT ACC AAT GCT TAA TCA GTG AGG CAC CTA TCT CAG CGA TCT GTC TAT TTC GTT CAT CCA TAG CTG CCT GAC TCC CCG TCG TGT AGA TAA CTA CGA TAC GGG AGG GCT TAC CAT CTG GCC CCA GTG CTG CAA TGA TAC CGC GAG ACC CAC GCT CAC CGG CTC CAG ATT TAT CAG CAA TAA ACC AGC CAG CCG GAA GGG CCG AGC GCA GAA GTG GTC CTG CAA CTT TAT CCG CCT CCA TCC AGT CTA TTA ATT GTT GCC GGG AAG CTA GAG TAA GTA GTT CGC CAG TTA ATA GTT TGC GCA ACG TTG TTG CCA TTG CTA CAG GCA TCG TGG TGT CAC GCT CGT CGT TTG GTA TGG CTT CAT TCA GCT CCG GTT CCC AAC GAT CAA GGC GAG TTA CAT GAT CCC CCA TGT TGT GCA AAA AAG CGG TTA GCT CCT TCG GTC CTC CGA TCG TTG TCA GAA GTA AGT TGG CCG CAG TGT TAT CAC TCA TGG TTA TGG CAG CAC TGC ATA ATT CTC TTA CTG TCA TGC CAT CCG TAA GAT GCT TTT CTG TGA CTG GTG AGT ACT CAA CCA AGT CAT TCT GAG AAT AGT GTA TGC GGC GAC CGA GTT GCT CTT GCC CGG CGT CAA TAC GGG ATA ATA CCG CGC CAC ATA GCA GAA CTT TAA AAG TGC TCA TCA TTG GAA AAC GTT CTT CGG GGC GAA AAC TCT CAA GGA TCT TAC CGC TGT TGA GAT CCA GTT CGA TGT AAC CCA CTC GTG CAC CCA ACT GAT CTT CAG CAT CTT TTA CTT TCA CCA GCG TTT CTG GGT GAG CAA AAA CAG GAA GGC AAA ATG CCG CAA AAA AGG GAA TAA GGG CGA CAC GGA AAT GTT GAA TAC TCA TAC TCT TCC TTT TTC AAT ATT ATT GAA GCA TTT ATC AGG GTT ATT GTC TCA TGA GCG GAT ACA TAT TTG AAT GTA TTT AGA AAA ATA AAC AAA TAG GGG TTC CGC GCA CAT TTC CCC GAA AAG TGC CAC CTG ACG TCT AAG AAA CCA TTA TTA TCA TGA CAT TAA CCT ATA AAA ATA GGC GTA TCA CGA GGC CCT TTC GTC TTC AC

198

3. pQE16_mDHFR_lacI_yPheRS_naph (8109 bp)



T5 promoter and lac operon 7 - 87 bp

- mDHFR_His 115 660 bp
- yPheRS_alpha subunit 1264 2787 bp
- yPheRS_beta subunit 2802 4589 bp
- lacI^q 5926 4842 bp
- ColE1 origin 6501 6526 bp
- beta-lactamase 7904 7044 bp

Full sequence

CTC GAG AAA TCA TAA AAA ATT TAT TTG CTT TGT GAG CGG ATA ACA ATT ATA ATA GAT TCA ATT GTG AGC GGA TAA AAA ATT TTG CTT TGT GAG CGG ATA ACA ATT ATA ATA GAT TCA ATT GTG AGC GGA TAA CAA TTT CAC ACA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG AGG AGA GGA TCC GGC ATC ATG GTT CGA CCA TTG AAC TCG ATC GTC GCC GCC GTG TCC CAA AAT ATG GGG ATT GGC AAG AAC GGA GAC CTA CCC TGG CCT CCG CTC AGG AAC GAG ATT CATG GTG ATT ATG GGT AGG AAA ACC TGG TTC TCC ATT CCT GAG AAG AAT CGA CCT TTA AAG GAC AGA ATT AAT ATA GTT CTC AGT AGA GAA CTC AAA GAA

CCA CCA CGA GGA GCT CAT TTT CTT GCC AAA AGT TTG GAT GAT GCC TTA AGA CTT ATT GAA CAA CCG GAA TTG GCA AGT AAA GTA GAC ATG GTT TGG ATA GTC GGA GGC AGT TCT GTT TAC CAG GAA GCC ATG AAT CAA CCA GGC CAC CTT AGA CTC TTT GTG ACA AGG ATC ATG CAG GAA TTT GAA AGT GAC ACG TTT TTC CCA GAA ATT GAT TTG GGG AAA TAT AAA CTT CTC CCA GAA TAC CCA GGC GTC CTC TCT GAG GTC CAG GAG GAA AAA GGC ATC AAG TAT AAG TTT GAA GTC TAC GAG AAG AAA GGT TCC AGA TCT CAT CAC CAT CAC CAT CAC TAA GCT TAA TTA GCT GAG CTT GGA CTC CTG TTG ATA GAT CCA GTA ATG ACC TCA GAA CTC CAT CTG GAT TTG TTC AGA ACG CTC GGT TGC CGC CGG GCG TTT TTT ATT GGT GAG AAT CCA AGC TAG CTT GGC GAG ATT TTC AGG AGC TAA GGA AGC TAA AAT GGA GAA AAA AAT CAC TGG ATA TAC CAC CGT TGA TAT ATC CCA ATG GCA TCG TAA AGA ACA TTT TGA GGC ATT TCA GTC AGT TGC TCA ATG TAC CTA TAA CCA GAC CGT TCA GCT GGC ACG ACA GGT TTC CCG ACT GGA AAG CGG GCA GTG AGC GCA ACG CAA TTA ATG TGA GTT AGC TCA CTC ATT AGG CAC CCC AGG CTT TAC ACT TTA TGC TTC CGG CTC GTA TGT TGT GTG GAA TTG TGA GCG GAT AAC AAT TTC ACA CAG GAA ACA GCT ATG ACC ATG ATT ACG CCA AGC TTG CAT GCC TGC AGT TGA CAA TTA ATC ATC GGC TCG TAT AAT GGA TCC AAT TGT GAG CGG AAT CGA TTT TCA CAC AGG AAA CAG ACC ATG AAT CTA GAG ATG TCT GAC TTC CAA TTA GAA ATT CTA AAG AAA CTA GAT GAA TTG GAT GAG ATC AAG TCC ACA CTG GCA ACT TTC CCT CAG CAC GGC TCT CAA GAT GTT CTT TCC GCT TTG AAC TCT TTG AAA GCC CAC AAC AAG TTA GAG TTT TCC AAG GTC GAC ACG GTT ACG TAT GAC TTG ACC AAA GAA GGT GCT CAA ATT TTG AAT GAA GGT TCG TAC GAA ATT AAA CTA GTC AAG CTC ATC CAA GAG TTG GGT CAA CTT CAA ATC AAA GAT GTG ATG TCC AAA CTA GGC CCT CAA GTT GGT AAG GTC GGT CAG GCT AGA GCT TTC AAG AAC GGC TGG ATC GCC AAA AAC GCC TCA AAC GAG CTT GAA CTC TCC GCA AAA TTG CAA AAT ACC GAT TTA AAT GAG CTT ACT GAT GAA ACG CAA TCT ATT CTA GCG CAA ATC AAG AAC AAC TCG CAT CTG GAT AGC ATT GAC GCC AAG ATT TTG AAC GAC TTG AAG AAA AGA AAG TTA ATT GCT CAA GGT AAA ATC ACA GAT TTC AGT GTC ACC AAA GGG CCA GAG TTC TCG ACC GAC CTC ACC AAA TTG GAA ACC GAT CTT ACC TCC GAC ATG GTC TCC ACC AAT GCA TAC AAG GAC TTG AAG TTC AAG CCT TAC AAT TTC AAT TCT CAA GGT GTG CAA ATA TCT TCA GGT GCT CTT CAC CCC TTA AAC AAA GTC AGA GAG GAA TTT AGA CAA ATT TTC TTT TCC ATG GGA TTC ACA GAG ATG CCC TCG AAC CAA TAC GTC GAG ACA GGT TTC TGG AAC TTC GAT GCC CTT TAC GTC CCA CAA CAG CAT CCT GCT CGT GAC CTG CAA GAC ACT TTC TAC ATC AAG GAC CCA CTA ACC GCT GAG TTG CCC GAT GAC AAG ACA TAC ATG GAC AAT ATC AAA GCC GTT CAC GAA CAG GGG AGA TTC GGG TCC ATC GGT TAT CGT TAC AAC TGG AAG CCA GAA GAA TGT CAA AAA TTG GTC TTG AGA ACT CAC TCC ACA GCC ATC TCT GCC AGA ATG CTG CAC GAT TTG GCC AAA GAT CCA AAG CCC ACC AGA TTG TTT TCT ATC GAC CGT GTT TTC CGT AAC GAA GCA GTT GAC GCC ACC CAT TTG GCC GAA TTC CAC CAG GTG GAA GGT GTT CTT GCC GAC TAC AAC ATT ACT CTG GGT GAC CTG ATC AAG TTC ATG GAA GAG TTT TTC GAA AGA ATG GGT GTC ACC GGT TTG AGA TTC AAG CCT ACC TAC GGG CCT TAC GGG GAG CCA TCT ATG GAA ATC TTT TCT TGG CAC GAA GGT TTG CAA AAA TGG GTC GAA ATC GGT AAC TTT GGT ATG TTC AGA CCA GAA ATG CTC GAG TCC ATG GGT CTA CCA AAG GAT CTA AGA GTC CTT GGT TGG GGG TTA TCC TTG GAA AGA CCT ACC ATG ATC AAA TAT AAG GTT CAA AAC ATC AGA GAA CTG TTA GGT CAT AAA GTC TCT TTG GAC TTT ATC GAA ACC AAT CCT GCT GCT AGA TTG GAC GAA GAC TTG TAC GAA TAA GGC AGG AAT AGA TTA TGC CTA CCG TCT CCG TGA ACA AGC AGC AAT TAT TTG ATC TTC TAG GCA AAG ACT ACA CTT CCC AAG AGT TCG ACG AAT TAT GTT TTG AAT TCG GTA TGG AAA TGG ACG AAG ACA CCA CAG AAG AGG CCT TGA AAA CCG GGG AGG AGC CGG AAT TGA AGC TTG ATA TCA GTG CCA ATC GTT ACG ATT TGC TTT GTA TCG AAG GTA TTT CAC AGT CGC TGA ACG AAT ACT TGG AAC GTA AAG AAA GAC CTG ACT ATA AAT TAA GCA AGC CAA CCA CTA AGT TGA TCA TCG ACA AAT CAA CGG AGC AAA TTA GAC CTT TTG CTA CCG CTG CTG TAT TGA GAA ATA TCA AGC TTA ACG AAA AAT CTT ACG CTT CTT TTA TTG CCT TGC AAG ATA AAT TAC ATG CCA ATC TAT GTA GAA ACA GAA GCT TGG TTG CCA TGG GTA CTC ACG ATT TAG ATT CAA TTG AAG GTC CAT TCC ATT ACA GAG CTC TAC CAC CAA AGG ACA TCA AGT TCG TAC CAT TGA ATC AAA CCC AAG AGT TTA CTG GTG ACA AAT TGA TCG AGT TTT ATA AAT CTC CAG AAC AGA AAA ACA ACA TAG GGA GAT ACG TTC ACA TTA TTG AGG ATT CTC CAG TCT TCC CAG TTA TTA TGG ACA GCA AAG ATC GTG

TTT GCT CCC TGC CAC CAT TAA TCA ATA GTG AAC ATT CGA AGA TCT CTG TGA ACA CCC GTA ACA TTT TGA TTG ATA TAA CCG CCA CCG ATA AGA CCA AAG CCG AGA TCG TTT TGA ACA TAT TAA CTA CAA TGT TCT CAC GTT ATT GTG ACG AAC CAT TCA CGG TTG AGC CTG TAG AAA TTG TCT CTG AAC ACA ATG GCC AAT CCC GTT TGG CGC CAA ACT TCA ACG ATA GAA TTA TGG ATG TCT CCA TTA AGT ACA TCA ACT CCT GTC TTG GCC TAG ATC AAT CCG CTG ATG AAA TTG CTC ATT GTC TAA AAA AGA TGT CGT TGC ATG CCG TTC AAT CAA AGG AAG ACA AGG ACA TCT TGC ACG TTG ACA TTC CGG TAA CTA GAC CTG ATA TTT TGC ACG CTT GTG ATA TAA TGG AAG ATG CCG CTG TCG GTT ATG GTT TCA ATA ATT TGC CAA AGG GTG AGA AAT TAT CCA ATG CCA ACT TCA TTG CCA AAC CAT TAC CAA TCA ACA AGG TTT CTG ATA TTT TCA GAG TTG CAT CCT CTC AAG CCA CGT GGG TTG AGG TTT TAC CAT TGA CCT TAT GTT CGC ACG ATG AAA ACT TTA AAT TTC TAA GAC AAT CCG ACA ATG ATG ATT TAG CTG TCA AAT TGG CCA ACC CAA AGA CTT TGG AAT ACC AAG TTG TTA GAA CCA CTT TAT TGC CTG GTA TCT TAA AGA CAG TCA AGG AAA ACA GAA AAC ATT CCT TAC CAA TCA AAG TCT TTG AAA CCG GTG ACG TTG TAT TTA AAG ACG ACA AAC TAG AAA GGA AGG CGT ACA ATG AAC GTC ACT GGG CTG CCA TCT ACG TGG GTA AGA ATT CTG GGT TTG AAA TCA TTC AAG GGT TAT TGG GTA AAA TCA TGC AAA CTT TTA GAA CAG AGT GGA TTG CAG ACT ACG GTG CTG CTG CTT CTG GCA GAG GTT ACT GGA TTG AAG AAG ACG ATT CTG TGA AAA CCT ATT TCC CAG GTA GAG GTG CCA AGG TCA TGT TCA GAT CCA AAG AAG GCG CTG AGC CAA AGC AAA TCG GCC ACT TGG GTG TCT TGC ATC CTG AAG TCA TGA TGA ATT TCG ACG TTC CAT TCG CTG CAT CCT TTG TAG AGG TTA ATG CCG AAG TCT TCC TAT AAT GTA ATG TTC TAA CAA AAA TTT TTA CTG ATT TAT AAA ACT TAT ATA GAT AGA TAG ACA TAT ATA TAT ATC TAT ATA TAG AAA CAC AAC TAA AGT TTA CCA TGT TTT ATA TAG GGT ACC GAG CTC GAA TTC ACT GGC CGT CGT TTT ACA ACG TCG TGA CTG GGA AAA CCG CGG TTA CGT GCA GTC GAT GAT AAG CTG TCA AAC ATG AGA ATT GTG CCT AAT GAG TGA GCT AAC TTA CAT TAA TTG CGT TGC GCT CAC TGC CCG CTT TCC AGT CGG GAA ACC TGT CGT GCC AGC TGC ATT AAT GAA TCG GCC AAC GCG CGG GGA GAG GCG GTT TGC GTA TTG GGC GCC AGG GTG GTT TTT CTT TTC ACC AGT GAG ACG GGC AAC AGC TGA TTG CCC TTC ACC GCC TGG CCC TGA GAG AGT TGC AGC AAG CGG TCC ACG CTG GTT TGC CCC AGC AGG CGA AAA TCC TGT TTG ATG GTG GTT AAC GGC GGG ATA TAA CAT GAG CTG TCT TCG GTA TCG TCG TAT CCC ACT ACC GAG ATA TCC GCA CCA ACG CGC AGC CCG GAC TCG GTA ATG GCG CGC ATT GCG CCC AGC GCC ATC TGA TCG TTG GCA ACC AGC ATC GCA GTG GGA ACG ATG CCC TCA TTC AGC ATT TGC ATG GTT TGT TGA AAA CCG GAC ATG GCA CTC CAG TCG CCT TCC CGT TCC GCT ATC GGC TGA ATT TGA TTG CGA GTG AGA TAT TTA TGC CAG CCA GCC AGA CGC AGA CGC GCC GAG ACA GAA CTT AAT GGG CCC GCT AAC AGC GCG ATT TGC TGG TGA CCC AAT GCG ACC AGA TGC TCC ACG CCC AGT CGC GTA CCG TCT TCA TGG GAG AAA ATA ATA CTG TTG ATG GGT GTC TGG TCA GAG ACA TCA AGA AAT AAC GCC GGA ACA TTA GTG CAG GCA GCT TCC ACA GCA ATG GCA TCC TGG TCA TCC AGC GGA TAG TTA ATG ATC AGC CCA CTG ACG CGT TGC GCG AGA AGA TTG TGC ACC GCC GCT TTA CAG GCT TCG ACG CCG CTT CGT TCT ACC ATC GAC ACC ACC ACG CTG GCA CCC AGT TGA TCG GCG CGA GAT TTA ATC GCC GCG ACA ATT TGC GAC GGC GCG TGC AGG GCC AGA CTG GAG GTG GCA ACG CCA ATC AGC AAC GAC TGT TTG CCC GCC AGT TGT TGT GCC ACG CGG TTG GGA ATG TAA TTC AGC TCC GCC ATC GCC GCT TCC ACT TTT TCC CGC GTT TTC GCA GAA ACG TGG CTG GCC TGG TTC ACC ACG CGG GAA ACG GTC TGA TAA GAG ACA CCG GCA TAC TCT GCG ACA TCG TAT AAC GTT ACT GGT TTC ACA TTC ACC ACC CTG AAT TGA CTC TCT TCC GGG CGC TAT CAT GCC ATA CCG CGA AAG GTT TTG CAC CAT TCG ATG GTG TCG GAA TTT CGG GCA GCG TTG GGT CCT GGC CAC GGG TGC GCA TGA CTT AAG CGG TGT GAA ATA CCG CAC AGA TGC GTA AGG AGA AAA TAC CGC ATC AGG CGC GCG GTA TCA GCT CAC TCA AAG GCG GTA ATA CGG TTA TCC ACA GAA TCA GGG GAT AAC GCA GGA AAG AAC ATG TGA GCA AAA GGC CAG CAA AAG GCC AGG AAC CGT AAA AAG GCC GCG TTG CTG GCG TTT TTC CAT AGG CTC CGC CCC CCT GAC GAG CAT CAC AAA AAT CGA CGC TCA AGT CAG AGG TGG CGA AAC CCG ACA GGA CTA TAA AGA TAC CAG GCG TTT CCC CCT GGA AGC TCC CTC GTG CGC TCT CCT GTT CCG ACC CTG CCG CTT ACC GGA TAC CTG TCC GCC TTT CTC CCT TCG GGA AGC GTG GCG CTT TCT CAA TGC TCA CGC TGT AGG TAT CTC AGT TCG GTG TAG GTC GTT CGC TCC AAG CTG GGC

TGT GTG CAC GAA CCC CCC GTT CAG CCC GAC CGC TGC GCC TTA TCC GGT AAC TAT CGT CTT GAG TCC AAC CCG GTA AGA CAC GAC TTA TCG CCA CTG GCA GCA GCC ACT GGT AAC AGG ATT AGC AGA GCG AGG TAT GTA GGC GGT GCT ACA GAG TTC TTG AAG TGG TGG CCT AAC TAC GGC TAC ACT AGA AGG ACA GTA TTT GGT ATC TGC GCT CTG CTG AAG CCA GTT ACC TTC GGA AAA AGA GTT GGT AGC TCT TGA TCC GGC AAA CAA ACC ACC GCT GGT AGC GGT GGT TTT TTT GTT TGC AAG CAG CAG ATT ACG CGC AGA AAA AAA GGA TCT CAA GAA GAT CCT TTG ATC TTT TCT ACG GGG TCT GAC GCT CAG TGG AAC GAA AAC TCA CGT TAA GGG ATT TTG GTC ATG AGA TTA TCA AAA AGG ATC TTC ACC TAG ATC CTT TTA AAT TAA AAA TGA AGT TTT AAA TCA ATC TAA AGT ATA TAT GAG TAA ACT TGG TCT GAC AGT TAC CAA TGC TTA ATC AGT GAG GCA CCT ATC TCA GCG ATC TGT CTA TTT CGT TCA TCC ATA GCT GCC TGA CTC CCC GTC GTG TAG ATA ACT ACG ATA CGG GAG GGC TTA CCA TCT GGC CCC AGT GCT GCA ATG ATA CCG CGA GAC CCA CGC TCA CCG GCT CCA GAT TTA TCA GCA ATA AAC CAG CCA GCC GGA AGG GCC GAG CGC AGA AGT GGT CCT GCA ACT TTA TCC GCC TCC ATC CAG TCT ATT AAT TGT TGC CGG GAA GCT AGA GTA AGT AGT TCG CCA GTT AAT AGT TTG CGC AAC GTT GTT GCC ATT GCT ACA GGC ATC GTG GTG TCA CGC TCG TCG TTT GGT ATG GCT TCA TTC AGC TCC GGT TCC CAA CGA TCA AGG CGA GTT ACA TGA TCC CCC ATG TTG TGC AAA AAA GCG GTT AGC TCC TTC GGT CCT CCG ATC GTT GTC AGA AGT AAG TTG GCC GCA GTG TTA TCA CTC ATG GTT ATG GCA GCA CTG CAT AAT TCT CTT ACT GTC ATG CCA TCC GTA AGA TGC TTT TCT GTG ACT GGT GAG TAC TCA ACC AAG TCA TTC TGA GAA TAG TGT ATG CGG CGA CCG AGT TGC TCT TGC CCG GCG TCA ATA CGG GAT AAT ACC GCG CCA CAT AGC AGA ACT TTA AAA GTG CTC ATC ATT GGA AAA CGT TCT TCG GGG CGA AAA CTC TCA AGG ATC TTA CCG CTG TTG AGA TCC AGT TCG ATG TAA CCC ACT CGT GCA CCC AAC TGA TCT TCA GCA TCT TTT ACT TTC ACC AGC GTT TCT GGG TGA GCA AAA ACA GGA AGG CAA AAT GCC GCA AAA AAG GGA ATA AGG GCG ACA CAG GGT TAT TGT CTC ATG AGC GGA TAC ATA TTT GAA TGT ATT TAG AAA AAT AAA CAA ATA GGG GTT CCG CGC ACA TTT CCC CGA AAA GTG CCA CCT GAC GTC TAA GAA ACC ATT ATT ATC ATG ACA TTA ACC TAT AAA AAT AGG CGT ATC ACG AGG CCC TTT CGT CTT CAC
4. pQE32_yPheRS (T415G) (6,895 bp)



T5 promoter and lac operon	7 - 87 bp
yPheRS_alpha subunit	115 - 1680 bp
yPheRS_beta subunit	1695 - 3482 bp
ColE1 origin	5103 - 5127 bp
beta-lactamase	6690 - 5830 bp

Full sequence

CTC GAG AAA TCA TAA AAA ATT TAT TTG CTT TGT GAG CGG ATA ACA ATT ATA ATA GAT TCA ATT GTG AGC GGA TAA CAA TTT CAC ACA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG AGA GGA TCT CAC CAT CAC CAT CAC CAT GGG ATC CAG ACC ATG AAT CTA GAG ATG TCT GAC TTC CAA TTA GAA ATT CTA AAG AAA CTA GAT GAA TTG GAT GAG ATC AAG TCC ACA CTG GCA ACT TTC CCT CAG CAC GGC TCT CAA GAT GTT CTT TCC GCT TTG AAC TCT TTG AAA GCC CAC AAC AAG TTA GAG TTT TCC AAG GTC GAC ACG GTT ACG TAT GAC TTG ACC AAA GAA GGT GCT CAA ATT TTG AAT GAA GGT TCG TAC GAA ATT AAA CTA GTC AAG CTC ATC CAA GAG TTG GGT CAA CTT CAA ATC AAA GAT GTG ATG TCC AAA CTA GGC CCT CAA GTT GGT AAG GTC GGT CAG GCT AGA GCT TTC AAG AAC GGC TGG ATC GCC AAA AAC GCC TCA AAC GAG CTT GAA CTC TCC GCA AAA TTG CAA AAT ACC GAT TTA AAT GAG CTT ACT GAT GAA ACG CAA TCT ATT CTA GCG CAA ATC AAG AAC AAC TCG CAT CTG GAT AGC ATT GAC GCC AAG ATT TTG AAC GAC TTG AAG AAA AGA AAG TTA ATT GCT CAA GGT AAA ATC ACA GAT TTC AGT GTC

ACC AAA GGG CCA GAG TTC TCG ACC GAC CTC ACC AAA TTG GAA ACC GAT CTT ACC TCC GAC ATG GTC TCC ACC AAT GCA TAC AAG GAC TTG AAG TTC AAG CCT TAC AAT TTC AAT TCT CAA GGT GTG CAA ATA TCT TCA GGT GCT CTT CAC CCC TTA AAC AAA GTC AGA GAG GAA TTT AGA CAA ATT TTC TTT TCC ATG GGA TTC ACA GAG ATG CCC TCG AAC CAA TAC GTC GAG ACA GGT TTC TGG AAC TTC GAT GCC CTT TAC GTC CCA CAA CAG CAT CCT GCT CGT GAC CTG CAA GAC ACT TTC TAC ATC AAG GAC CCA CTA ACC GCT GAG TTG CCC GAT GAC AAG ACA TAC ATG GAC AAT ATC AAA GCC GTT CAC GAA CAG GGG AGA TTC GGG TCC ATC GGT TAT CGT TAC AAC TGG AAG CCA GAA GAA TGT CAA AAA TTG GTC TTG AGA ACT CAC TCC ACA GCC ATC TCT GCC AGA ATG CTG CAC GAT TTG GCC AAA GAT CCA AAG CCC ACC AGA TTG TTT TCT ATC GAC CGT GTT TTC CGT AAC GAA GCA GTT GAC GCC ACC CAT TTG GCC GAA TTC CAC CAG GTG GAA GGT GTT CTT GCC GAC TAC AAC ATT ACT CTG GGT GAC CTG ATC AAG TTC ATG GAA GAG TTT TTC GAA AGA ATG GGT GTC ACC GGT TTG AGA TTC AAG CCT ACC TAC AAT CCT TAC GGC GAG CCA TCA ATG GAA ATC TTT TCT TGG CAC GAA GGT TTG CAA AAA TGG GTC GAA ATC GGT AAC TCT GGT ATG TTC AGA CCA GAA ATG CTC GAG TCC ATG GGT CTA CCA AAG GAT CTA AGA GTC CTT GGT TGG GGG TTA TCC TTG GAA AGA CCT ACC ATG ATC AAA TAT AAG GTT CAA AAC ATC AGA GAA CTG TTA GGT CAT AAA GTC TCT TTG GAC TTT ATC GAA ACC AAT CCT GCT GCT AGA TTG GAC GAA GAC TTG TAC GAA TAA GGC AGG AAT AGA TTA TGC CTA CCG TCT CCG TGA ACA AGC AGC AAT TAT TTG ATC TTC TAG GCA AAG ACT ACA CTT CCC AAG AGT TCG ACG AAT TAT GTT TTG AAT TCG GTA TGG AAA TGG ACG AAG ACA CCA CAG AAG AGG CCT TGA AAA CCG GGG AGG AGC CGG AAT TGA AGC TTG ATA TCA GTG CCA ATC GTT ACG ATT TGC TTT GTA TCG AAG GTA TTT CAC AGT CGC TGA ACG AAT ACT TGG AAC GTA AAG AAA GAC CTG ACT ATA AAT TAA GCA AGC CAA CCA CTA AGT TGA TCA TCG ACA AAT CAA CGG AGC AAA TTA GAC CTT TTG CTA CCG CTG CTG TAT TGA GAA ATA TCA AGC TTA ACG AAA AAT CTT ACG CTT CTT TTA TTG CCT TGC AAG ATA AAT TAC ATG CCA ATC TAT GTA GAA ACA GAA GCT TGG TTG CCA TGG GTA CTC ACG ATT TAG ATT CAA TTG AAG GTC CAT TCC ATT ACA GAG CTC TAC CAC CAA AGG ACA TCA AGT TCG TAC CAT TGA ATC AAA CCC AAG AGT TTA CTG GTG ACA AAT TGA TCG AGT TTT ATA AAT CTC CAG AAC AGA AAA ACA ACA TAG GGA GAT ACG TTC ACA TTA TTG AGG ATT CTC CAG TCT TCC CAG TTA TTA TGG ACA GCA AAG ATC GTG TTT GCT CCC TGC CAC CAT TAA TCA ATA GTG AAC ATT CGA AGA TCT CTG TGA ACA CCC GTA ACA TTT TGA TTG ATA TAA CCG CCA CCG ATA AGA CCA AAG CCG AGA TCG TTT TGA ACA TAT TAA CTA CAA TGT TCT CAC GTT ATT GTG ACG AAC CAT TCA CGG TTG AGC CTG TAG AAA TTG TCT CTG AAC ACA ATG GCC AAT CCC GTT TGG CGC CAA ACT TCA ACG ATA GAA TTA TGG ATG TCT CCA TTA AGT ACA TCA ACT CCT GTC TTG GCC TAG ATC AAT CCG CTG ATG AAA TTG CTC ATT GTC TAA AAA AGA TGT CGT TGC ATG CCG TTC AAT CAA AGG AAG ACA AGG ACA TCT TGC ACG TTG ACA TTC CGG TAA CTA GAC CTG ATA TTT TGC ACG CTT GTG ATA TAA TGG AAG ATG CCG CTG TCG GTT ATG GTT TCA ATA ATT TGC CAA AGG GTG AGA AAT TAT CCA ATG CCA ACT TCA TTG CCA AAC CAT TAC CAA TCA ACA AGG TTT CTG ATA TTT TCA GAG TTG CAT CCT CTC AAG CCA CGT GGG TTG AGG TTT TAC CAT TGA CCT TAT GTT CGC ACG ATG AAA ACT TTA AAT TTC TAA GAC AAT CCG ACA ATG ATG ATT TAG CTG TCA AAT TGG CCA ACC CAA AGA CTT TGG AAT ACC AAG TTG TTA GAA CCA CTT TAT TGC CTG GTA TCT TAA AGA CAG TCA AGG AAA ACA GAA AAC ATT CCT TAC CAA TCA AAG TCT TTG AAA CCG GTG ACG TTG TAT TTA AAG ACG ACA AAC TAG AAA GGA AGG CGT ACA ATG AAC GTC ACT GGG CTG CCA TCT ACG TGG GTA AGA ATT CTG GGT TTG AAA TCA TTC AAG GGT TAT TGG GTA AAA TCA TGC AAA CTT TTA GAA CAG AGT GGA TTG CAG ACT ACG GTG CTG CTG CTT CTG GCA GAG GTT ACT GGA TTG AAG AAG ACG ATT CTG TGA AAA CCT ATT TCC CAG GTA GAG GTG CCA AGG TCA TGT TCA GAT CCA AAG AAG GCG CTG AGC CAA AGC AAA TCG GCC ACT TGG GTG TCT TGC ATC CTG AAG TCA TGA TGA ATT TCG ACG TTC CAT TCG CTG CAT CCT TTG TAG AGG TTA ATG CCG AAG TCT TCC TAT AAT GTA ATG TTC TAA CAA AAA TTT TTA CTG ATT TAT AAA ACT TAT ATA GAT AGA TAG ACA CAT AAA TGT ATC TAT ATA TAG AAA CAC AAC TAA AGT TTA CCA TCT TAA AGG TAA TGG GTA CCC CGG GTC GAC CTG CAG CCA AGC TTA ATT AGC TGA GCT TGG ACT CCT GTT GAT AGA TCC AGT AAT GAC CTC AGA ACT CCA TCT GGA TTT GTT CAG AAC GCT CGG TTG CCG CCG GGC GTT TTT TAT TGG TGA GAA TCC AAG CTA GCT TGG CGA GAT TTT CAG GAG CTA AGG AAG CTA AAA

TGG AGA AAA AAA TCA CTG GAT ATA CCA CCG TTG ATA TAT CCC AAT GGC ATC GTA AAG AAC ATT TTG AGG CAT TTC AGT CAG TTG CTC AAT GTA CCT ATA ACC AGA CCG TTC AGC TGG ATA TTA CGG CCT TTT TAA AGA CCG TAA AGA AAA ATA AGC ACA AGT TTT ATC CGG CCT TTA TTC ACA TTC TTG CCC GCC TGA TGA ATG CTC ATC CGG AAT TTC GTA TGG CAA TGA AAG ACG GTG AGC TGG TGA TAT GGG ATA GTG TTC ACC CTT GTT ACA CCG TTT TCC ATG AGC AAA CTG AAA CGT TTT CAT CGC TCT GGA GTG AAT ACC ACG ACG ATT TCC GGC AGT TTC TAC ACA TAT ATT CGC AAG ATG TGG CGT GTT ACG GTG AAA ACC TGG CCT ATT TCC CTA AAG GGT TTA TTG AGA ATA TGT TTT TCG TCT CAG CCA ATC CCT GGG TGA GTT TCA CCA GTT TTG ATT TAA ACG TGG CCA ATA TGG ACA ACT TCT TCG CCC CCG TTT TCA CCA TGG GCA AAT ATT ATA CGC AAG GCG ACA AGG TGC TGA TGC CGC TGG CGA TTC AGG TTC ATC ATG CCG TTT GTG ATG GCT TCC ATG TCG GCA GAA TGC TTA ATG AAT TAC AAC AGT ACT GCG ATG AGT GGC AGG GCG GGG CGT AAT TTT TTT AAG GCA GTT ATT GGT GCC CTT AAA CGC CTG GGG TAA TGA CTC TCT AGC TTG AGG CAT CAA ATA AAA CGA AAG GCT CAG TCG AAA GAC TGG GCC TTT CGT TTT ATC TGT TGT TTG TCG GTG AAC GCT CTC CTG AGT AGG ACA AAT CCG CCC TCT AGA GCT GCC TCG CGC GTT TCG GTG ATG ACG GTG AAA ACC TCT GAC ACA TGC AGC TCC CGG AGA CGG TCA CAG CTT GTC TGT AAG CGG ATG CCG GGA GCA GAC AAG CCC GTC AGG GCG CGT CAG CGG GTG TTG GCG GGT GTC GGG GCG CAG CCA TGA CCC AGT CAC GTA GCG ATA GCG GAG TGT ATA CTG GCT TAA CTA TGC GGC ATC AGA GCA GAT TGT ACT GAG AGT GCA CCA TAT GCG GTG TGA AAT ACC GCA CAG ATG CGT AAG GAG AAA ATA CCG CAT CAG GCG CTC TTC CGC TTC CTC GCT CAC TGA CTC GCT GCG CTC GGT CGT TCG GCT GCG GCG AGC GGT ATC AGC TCA CTC AAA GGC GGT AAT ACG GTT ATC CAC AGA ATC AGG GGA TAA CGC AGG AAA GAA CAT GTG AGC AAA AGG CCA GCA AAA GGC CAG GAA CCG TAA AAA GGC CGC GTT GCT GGC GTT TTT CCA TAG GCT CCG CCC CCC TGA CGA GCA TCA CAA AAA TCG ACG CTC AAG TCA GAG GTG GCG AAA CCC GAC AGG ACT ATA AAG ATA CCA GGC GTT TCC CCC TGG AAG CTC CCT CGT GCG CTC TCC TGT TCC GAC CCT GCC GCT TAC CGG ATA CCT GTC CGC CTT TCT CCC TTC GGG AAG CGT GGC GCT TTC TCA TAG CTC ACG CTG TAG GTA TCT CAG TTC GGT GTA GGT CGT TCG CTC CAA GCT GGG CTG TGT GCA CGA ACC CCC CGT TCA GCC CGA CCG CTG CGC CTT ATC CGG TAA CTA TCG TCT TGA GTC CAA CCC GGT AAG ACA CGA CTT ATC GCC ACT GGC AGC AGC CAC TGG TAA CAG GAT TAG CAG AGC GAG GTA TGT AGG CGG TGC TAC AGA GTT CTT GAA GTG GTG GCC TAA CTA CGG CTA CAC TAG AAG GAC AGT ATT TGG TAT CTG CGC TCT GCT GAA GCC AGT TAC CTT CGG AAA AAG AGT TGG TAG CTC TTG ATC CGG CAA ACA AAC CAC CGC TGG TAG CGG TGG TTT TTT TGT TTG CAA GCA GCA GAT TAC GCG CAG AAA AAA AGG ATC TCA AGA AGA TCC TTT GAT CTT TTC TAC GGG GTC TGA CGC TCA GTG GAA CGA AAA CTC ACG TTA AGG GAT TTT GGT CAT GAG ATT ATC AAA AAG GAT CTT CAC CTA GAT CCT TTT AAA TTA AAA ATG AAG TTT TAA ATC AAT CTA AAG TAT ATA TGA GTA AAC TTG GTC TGA CAG TTA CCA ATG CTT AAT CAG TGA GGC ACC TAT CTC AGC GAT CTG TCT ATT TCG TTC ATC CAT AGT TGC CTG ACT CCC CGT CGT GTA GAT AAC TAC GAT ACG GGA GGG CTT ACC ATC TGG CCC CAG TGC TGC AAT GAT ACC GCG AGA CCC ACG CTC ACC GGC TCC AGA TTT ATC AGC AAT AAA CCA GCC AGC CGG AAG GGC CGA GCG CAG AAG TGG TCC TGC AAC TTT ATC CGC CTC CAT CCA GTC TAT TAA TTG TTG CCG GGA AGC TAG AGT AAG TAG TTC GCC AGT TAA TAG TTT GCG CAA CGT TGT TGC CAT TGC TAC AGG CAT CGT GGT GTC ACG CTC GTC GTT TGG TAT GGC TTC ATT CAG CTC CGG TTC CCA ACG ATC AAG GCG AGT TAC ATG ATC CCC CAT GTT GTG CAA AAA AGC GGT TAG CTC CTT CGG TCC TCC GAT CGT TGT CAG AAG TAA GTT GGC CGC AGT GTT ATC ACT CAT GGT TAT GGC AGC ACT GCA TAA TTC TCT TAC TGT CAT GCC ATC CGT AAG ATG CTT TTC TGT GAC TGG TGA GTA CTC AAC CAA GTC ATT CTG AGA ATA GTG TAT GCG GCG ACC GAG TTG CTC TTG CCC GGC GTC AAT ACG GGA TAA TAC CGC GCC ACA TAG CAG AAC TTT AAA AGT GCT CAT CAT TGG AAA ACG TTC TTC GGG GCG AAA ACT CTC AAG GAT CTT ACC GCT GTT GAG ATC CAG TTC GAT GTA ACC CAC TCG TGC ACC CAA CTG ATC TTC AGC ATC TTT TAC TTT CAC CAG CGT TTC TGG GTG AGC AAA AAC AGG AAG GCA AAA TGC CGC AAA AAA GGG AAT AAG GGC GAC ACG GAA ATG TTG AAT ACT CAT ACT CTT CCT TTT TCA ATA TTA TTG AAG CAT TTA TCA GGG TTA TTG TCT CAT GAG CGG ATA CAT ATT TGA ATG TAT TTA GAA AAA TAA ACA AAT AGG GGT TCC GCG CAC ATT TCC CCG AAA AGT GCC ACC TGA CGT CTA AGA AAC CAT TAT TAT CAT GAC ATT AAC CTA TAA AAA TAG GCG TAT CAC GAG GCC CTT TCG TCT TCA C

5. pREP4_ytRNA^{Phe}_AAA (4084 bp)



Vector size	4084 bp
agpt promoter	259-260 bp
RBS	336-344 bp
Aminoglycosie phosphotransferase	357-1151 bp
<i>lacI</i> promoter	2638-2639 bp
<i>lacI</i> repressor	1529-2611 bp
lpp promoter	3029-3083 bp
ytRNA	2945-3020 bp
transcriptional terminator(rrnC)	2938-2909 bp
p15A origin of replication	3408-3411 bp

Full sequence

AAG CTT CAC GCT GCC GCA AGC ACT CAG GGC GCA AGG GCT GCT AAA GGA AGC GGA ACA CGT AGA AAG CCA GTC CGC AGA AAC GGT GCT GAC CCC GGA TGA ATG TCA GCT

206

GGG	CTA	TCT	GGA	CAA	GGG	AAA	ACG	CAA	GCG	CAA	AGA	GAA	AGC	AGG	TAG	CTT
GTG	GGC	TTA	CAT	GGC	GAT	AGC	TAG	ACT	GGG	CGG	TTT	TAT	GGA	CAG	CAA	GCG
CGG	ΔΔΤ	TGC	CAG	CTG	GGG	CGC	ССТ	CTG	GTA	AGG	TTG	GGA	AGC	ССТ	GCA	AAG
	CCA	TCC	Стт		TCC	CGC	CZZ	CCA		CAT	CCC	CCA	CCC	CAT	CZZ	CAT
			CAC	CAT		CCT	CCT		CCA	TCC		AAC		ATC	Слт	TCC
AIC			CAG	CCC		CCC	TG0	707	CCC		TIG TIG				CCC	C A C
	GII		CGG	CCG		DDD 7 mg	TGG	AGA	GGC	TAI	add	GCI	AIG	ACI	DDD 7 CC	CAC
AGA	CAA	TCG	GCT	GCT	CTG	ATG	CCG	CCG	TGT	TCC	GGC	TGT	CAG	CGC	AGG	GGC
CGG	TTC	.111.	TTG	TCA	AGA	CCG	ACC	ΊĠΊ	CCG	GTG	CCC	ΊGΑ	ATG	AAC	TGC	AGG
AGG	CAG	CGC	GGC	TAT	CGT	GGC	TGG	CCA	CGA	CGG	GCG	TTC	CTT	GCG	CAG	CTG
TCG	ACG	TTG	TCA	CTG	AAG	CGG	GAA	GGG	ACT	GGC	TGC	TAT	TGG	GCG	AAG	TGC
GGC	AGG	ATC	TCC	TGT	CAT	CTC	ACC	TTG	CTC	CTG	CCG	AGA	AAG	TAT	CCA	TCA
CTG	ATG	CAA	TGC	GGC	GGC	TGC	ATA	CGC	TTG	ATC	CGG	CTA	CCT	GCC	CAT	TCG
ACC	AAG	CGA	AAC	ATC	GCA	TCG	AGC	GAG	CAC	GTA	CTC	GGA	TGG	AAG	CCG	GTC
TCG	ATC	AGG	ATG	ATC	TGG	ACG	AAG	AGC	ATC	AGG	GGC	TCG	CGC	CAG	CCG	AAC
TCG	CCA	GGC	TCA	AGG	CGC	GCA	TGC	CCG	ACG	GCG	AGG	ATC	TCG	TCG	TGA	CCC
CCC	ATC	CCT	CCT	TGC	CGA			TCC	TCC		ATC	CCC	CCT	<u>т</u> тт	CTC	ССС
		CTC	CCC	CCC	TCC	CTC	TCA	CCC			ATC				CCT	TCC
CCG	ACI	SIG Ama		CTTC	1GG	J C C		CGG	ACC	GCI NNTT	AIC	AGG	ACA	COT	TGC TGC	TGG
	J CC		TIG maa	CIG		AGC		GCG	GCG 7 dd	AAI	UUU maa	CIG	ACC	GCI MIC	add	TCG
	ACG	GIA	TCG	CCG	CIC	CCG	AII	CGC	AGC	GCA	TCG	CCI		AIC	GCC	TIC
ACG	AG'I'	TCT	TCT	GAG	CGG	GAC	TCT	GGG	G.II.	CGA	AA.I.	GAC	CGA	CCA	AGC	GAC
CAA	CCT	GCC	ATC	ACG	AGA	TTT	CGA	TTC	CAC	CGC	CGC	CTT	CTA	TGA	AAG	GTT
CTT	CGG	AAT	CGT	TTT	CCG	GGA	CGC	CGG	CTG	GAT	GAT	CCT	CCA	GCG	CGG	GGA
CAT	GCT	GGA	GTT	CTT	CGC	CCA	CCC	CGG	GCT	CGA	TCC	CCT	CGC	GAG	TTG	GTT
CTG	CTG	CCT	GAG	GCT	GGA	CGA	CCT	CGC	GGA	GTT	CTA	CCG	GCA	GTG	CAA	ATC
CGG	CAT	CCA	GGA	AAC	CAG	CAG	CGG	CTA	TCC	GCG	CAT	CCA	TGC	CCC	CGA	ACT
GGA	GTG	GGG	AGG	CAC	GAT	GGC	CGC	TTT	GGT	CGA	CAA	TTC	GCG	CTA	ACT	TAC
AAT	TGC	GTT	GCG	CTC	ACT	GCC	CGC	TTT	CCA	GTC	GGG	AAA	CCT	GTC	GTG	CCA
GCA	TTA	ATG	AAT	CGG	CCA	ACG	CGC	GGG	GAG	AGG	CGG	TTT	GCG	TAT	TGG	GCG
GGG	TGG	TTT	TTC	TTT	TCA	CCA	GTG	AGA	CGG	GCA	ACA	GCT	GAT	TGC	CCT	TCA
CCT	GGC	CCT	GAG	AGA	GTT	GCA	GCA	AGC	GGT	CCA	CGC	TGG	TTT	GCC	CCA	GCA
GAA	AAT	CCT	GTT	TGA	TGG	TGG	TTA	ACG	GCG	GGA	TAT	AAC	ATG	AGC	TGT	CTT
TAT	CGT	CGT	ATC	CCA	CTA	CCG	AGA	TAT	CCG	CAC	CAA	CGC	GCA	GCC	CGG	ACT
TAA	TGG	CGC	GCA	TTG	CGC	CCA	GCG	CCA	TCT	GAT	CGT	TGG	CAA	CCA	GCA	TCG
TGG	GAA	CGA	TGC	CCT	CAT	TCA	GCA	TTT	GCA	TGG	TTT	GTT	GAA	AAC	CGG	ACA
CAC	TCC	AGT	CGC	СТТ	CCC	GTT	CCG	СТА	TCG	GCT	GAA	TTT	GAT	TGC	GAG	TGA
ATT	ТАТ	GCC	AGC	CAG	CCA	GAC	GCA	GAC	GCG	CCG	AGA	CAG	AAC	TTA	ATG	GGC
CTA	ACA	GCG	CGA	TTT	GCT	GGT	GAC	CCA	ATG	CGA	CCA	GAT	GCT	CCA	CGC	CCA
CCC	TAC	ССТ	Стт	CAT	CCC	AGA			TAC	TGT	TGA	TCC	CTC		ССТ	CAG
СУД		COI	CII	0111	000	11011	1 11 11 1	T T TT T	1110	TOT	1011	100				0110
CAI	1.111	CAA	ለጥለ	ACC	CCC		$C \Lambda T$	TAC	TCC	ACC	CAC	CTTT	CCA	CNC	C777	тсс
CCT	CAA	GAA	ATA	ACG	CCG	GAA	CAT	TAG TGA	TGC TCA	AGG	CAG	CTT	CCA	CAG	CAA	TGG
CCT	GGT	GAA CAT	ATA CCA	ACG GCG	CCG GAT	GAA AGT	CAT TAA	TAG TGA	TGC TCA	AGG GCC	CAG CAC	CTT TGA	CCA CGC	CAG GTT	CAA GCG	TGG CGA
CCT GAT	GGT TGT	GAA CAT GCA	ATA CCA CCG	ACG GCG CCG	CCG GAT CTT	GAA AGT TAC	CAT TAA AGG	TAG TGA CTT	TGC TCA CGA	AGG GCC CGC	CAG CAC CGC	CTT TGA TTC	CCA CGC GTT	CAG GTT CTA	CAA GCG CCA	TGG CGA TCG
CCT GAT CCA	GGT TGT CCA	GAA CAT GCA CGC	ATA CCA CCG TGG	ACG GCG CCG CAC	CCG GAT CTT CCA	GAA AGT TAC GTT	CAT TAA AGG GAT	TAG TGA CTT CGG	TGC TCA CGA CGC	AGG GCC CGC GAG	CAG CAC CGC ATT	CTT TGA TTC TAA	CCA CGC GTT TCG	CAG GTT CTA CCG	CAA GCG CCA CGA	TGG CGA TCG CAA
CCT GAT CCA GCG	GGT TGT CCA ACG	GAA CAT GCA CGC GCG	ATA CCA CCG TGG CGT	ACG GCG CCG CAC GCA	CCG GAT CTT CCA GGG	GAA AGT TAC GTT CCA	CAT TAA AGG GAT GAC	TAG TGA CTT CGG TGG	TGC TCA CGA CGC AGG	AGG GCC CGC GAG TGG	CAG CAC CGC ATT CAA	CTT TGA TTC TAA CGC	CCA CGC GTT TCG CAA	CAG GTT CTA CCG TCA	CAA GCG CCA CGA GCA	TGG CGA TCG CAA ACG
CCT GAT CCA GCG GTT	GGT TGT CCA ACG TGC	GAA CAT GCA CGC GCG CCG	ATA CCA CCG TGG CGT CCA	ACG GCG CCG CAC GCA GTT	CCG GAT CTT CCA GGG GTT	GAA AGT TAC GTT CCA GTG	CAT TAA AGG GAT GAC CCA	TAG TGA CTT CGG TGG CGC	TGC TCA CGA CGC AGG GGT	AGG GCC CGC GAG TGG TGG	CAG CAC CGC ATT CAA GAA	CTT TGA TTC TAA CGC TGT	CCA CGC GTT TCG CAA AAT	CAG GTT CTA CCG TCA TCA	CAA GCG CCA CGA GCA GCT	TGG CGA TCG CAA ACG CCG
CCT GAT CCA GCG GTT TCG	GGT TGT CCA ACG TGC CCG	GAA CAT GCA CGC GCG CCG CTT	ATA CCA CCG TGG CGT CCA CCA	ACG GCG CCG CAC GCA GTT CTT	CCG GAT CTT CCA GGG GTT TTT	GAA AGT TAC GTT CCA GTG CCC	CAT TAA AGG GAT GAC CCA GCG	TAG TGA CTT CGG TGG CGC TTT	TGC TCA CGA CGC AGG GGT TCG	AGG GCC CGC GAG TGG TGG CAG	CAG CAC CGC ATT CAA GAA AAA	CTT TGA TTC TAA CGC TGT CGT	CCA CGC GTT TCG CAA AAT GGC	CAG GTT CTA CCG TCA TCA TGG	CAA GCG CCA CGA GCA GCT CCT	TGG CGA TCG CAA ACG CCG GGT
CCT GAT CCA GCG GTT TCG CCA	GGT TGT CCA ACG TGC CCG CGC	GAA CAT GCA CGC GCG CCG CCG CTT GGG	ATA CCA CCG TGG CGT CCA CCA AAA	ACG GCG CCG CAC GCA GTT CTT CGG	CCG GAT CTT CCA GGG GTT TTT TCT	GAA AGT TAC GTT CCA GTG CCC GAT	CAT TAA AGG GAT GAC CCA GCG AAG	TAG TGA CTT CGG TGG CGC TTT AGA	TGC TCA CGA CGC AGG GGT TCG CAC	AGG GCC CGC GAG TGG TGG CAG CGG	CAG CAC CGC ATT CAA GAA AAA CAT	CTT TGA TTC TAA CGC TGT CGT ACT	CCA CGC GTT TCG CAA AAT GGC CTG	CAG GTT CTA CCG TCA TCA TGG CGA	CAA GCG CCA CGA GCA GCT CCT CAT	TGG CGA TCG CAA ACG CCG GGT CGT
CCT GAT CCA GCG GTT TCG CCA ACG	GGT TGT CCA ACG TGC CCG CCG CGC TTA	GAA CAT GCA CGC GCG CCG CTT GGG CTG	ATA CCA CCG TGG CGT CCA CCA AAA GTT	ACG GCG CCG CAC GCA GTT CTT CGG TCA	CCG GAT CTT CCA GGG GTT TTT TCT CAT	GAA AGT TAC GTT CCA GTG CCC GAT TCA	CAT TAA AGG GAT GAC CCA GCG AAG CCA	TAG TGA CTT CGG TGG CGC TTT AGA CCC	TGC TCA CGA CGC AGG GGT TCG CAC TGA	AGG GCC CGC GAG TGG TGG CAG CAG CGG ATT	CAG CAC CGC ATT CAA GAA AAA CAT GAC	CTT TGA TTC TAA CGC TGT CGT ACT TCT	CCA CGC GTT TCG CAA AAT GGC CTG CTT	CAG GTT CTA CCG TCA TCA TGG CGA CCG	CAA GCG CCA CGA GCA GCT CCT CAT GGC	TGG CGA TCG CAA ACG CCG GGT GGT GCT
CCT GAT CCA GCG GTT TCG CCA ACG ATG	GGT TGT CCA ACG TGC CCG CCG CGC TTA CCA	GAA CAT GCA CGC GCG CCG CCG CTT GGG CTG TAC	ATA CCA CCG TGG CGT CCA CCA AAA GTT CGC	ACG GCG CCG GCA GTT CTT CGG TCA GAA	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG	GAA AGT TAC GTT CCA GTG CCC GAT TCA TTT	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT	AGG GCC CGC GAG TGG TGG CAG CGG ATT CGA	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG	CTT TGA TTC TAA CGC TGT CGT ACT TCT TGT	CCA CGC GTT TCG CAA AAT GGC CTG CTT CAA	CAG GTT CTA CCG TCA TCA TCA TGG CGA CCG CGT	CAA GCG CCA CGA GCA GCT CCT CAT GGC AAA	TGG CGA TCG CAA ACG CCG GGT CGT GCT TGC
CCT GAT CCA GCG GTT TCG CCA ACG ATG CCG	GGT TGT CCA ACG TGC CCG CCG CGC TTA CCA CTT	GAA CAT GCA CGC GCG CCG CTT GGG CTG TAC CGC	ATA CCA CCG TGG CGT CCA CCA AAA GTT CGC CTT	ACG GCG CCG GCA GTT CTT CGG TCA GAA CGC	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG GCG	GAA AGT TAC GTT CCA GTG CCC GAT TCA TTT CGA	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GTC	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC	AGG GCC CGC GAG TGG TGG CAG CGG ATT CGA CCT	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GTC	CTT TGA TTC TAA CGC TGT CGT ACT TCT TGT CCT	CCA CGC GTT TCG CAA AAT GGC CTG CTT CAA CCT	CAG GTT CTA CCG TCA TCA TCA TGG CGA CCG CGT GTT	CAA GCG CCA CGA GCA GCT CCT CAT GGC AAA CAG	TGG CGA TCG CAA ACG CCG GGT CGT GCT TGC CTA
CCT GAT CCA GCG GTT TCG CCA ACG ATG CCG ACG	GGT TGT CCA ACG TGC CCG CGC TTA CCA CTT GGG	GAA CAT GCA CGC GCG CTT GGG CTT GGG CTG TAC CGC TGG	ATA CCA CCG TGG CGT CCA AAA GTT CGC CTT TGC	ACG GCG CAC GCA GTT CTT CGG TCA GAA CGC GTA	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG GCG ACG	GAA AGT TAC GTT CCA GTG CCC GAT TCA TTT CGA GCA	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT AAA	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GTC GCA	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG	AGG GCC CGC GAG TGG TGG CAG CGG ATT CGA CCT CCG	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GTC GAC	CTT TGA TTC TAA CGC TGT CGT ACT TCT TGT CCT ATC	CCA CGC GTT TCG CAA AAT GGC CTG CTG CTT CAA CCT AGC	CAG GTT CTA CCG TCA TCA TCA TCG CGA CCG CGT GTT GCT	CAA GCG CCA GCA GCT CCT CAT GGC AAA CAG AGC	TGG CGA TCG CAA ACG CCG GGT CGT GCT TGC CTA GGA
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG TAC	GGT TGT CCA ACG TGC CCG CCG CCG CCG CCA CCA CTT GGG TGG	GAA CAT GCA CGC CCG CTG GGG CTG TAC CGC TGG CGA	ATA CCA CCG CGT CCA CCA AAA GTT CGC CTT TGC AAG	ACG GCG CCG GCA GTT CTT CGG TCA GAA CGC GTA GGG	CCG GAT CTT CCA GGG GTT TTT CAT AGG GCG ACG GAG	GAA AGT TAC GTT CCA GTG CCC GAT TCA TTT CGA GCA GTG	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT AAA CTG	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC GCA CAA	TGC TCA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC	AGG GCC CGC GAG TGG CAG CGG ATT CGA CCT CCG GAT	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GTC GAC TAA	CTT TGA TTC TAA CGC TGT CGT TCT TGT CCT ATC GTT	CCA CGC GTT TCG CAA AAT GGC CTG CTG CTT CAA CCT AGC GGG	CAG GTT CTA CCG TCA TCA TCA TCA CCG CGT GCT GCT TAA	CAA GCG CCA GCA GCT CCT CAT GGC AAA CAG AGC CGC	TGG CGA TCG CAA ACG CCG GGT CGT GCT GGA CAG
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG ACG TAC TTT	GGT TGT CCA ACG TGC CCG CCG CCG CCA CCA CTT GGG TGG CCC	GAA CAT GCA CGC CCG CTT GGG CTG TAC CGC TGG CGA AGT	ATA CCA CCG CGT CCA AAA GTT CGC CTT TGC AAG CAC	ACG GCG CCG GCA GTT CTT CCG TCA GAA CGC GTA GGG GAC	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG GCG ACG GAG GTT	GAA AGT TAC GTT CCA GTG CCC GAT TCA TTT CGA GCA GTG GTA	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT AAA CTG AAA	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC GCA CAA CGA	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG	AGG GCC CGC GAG TGG CAG CGG ATT CGA CCT CCG GAT CCA	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GTC GAC TAA GTG	CTT TGA TTC TAA CGC TGT CGT TCT TGT CCT ATC GTT CCA	CCA GTT TCG CAA AAT GGC CTG CTT CAA CCT AGC GGG AGC	CAG GTT CTA CCG TCA TCA TCA TCA TCA CCG CGT GCT TAA TTA	CAA GCG CCA GCA GCT CCT CAT GGC AAA CAG AGC CGC AAA	TGG CGA TCG CAA ACG GGT CGT GCT GCT GGA CAG AAA
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG ACG TAC TTT CTT	GGT TGT CCA ACG TGC CCG CCG CGC TTA CCA CTT GGG TGG CCC AGC	GAA CAT GCA CGC GCG CCG CTT GGG CTG TAC CGC TGG CGA AGT TTT	ATA CCA CCG TGG CGT CCA CCA AAA GTT CGC CTT TGC AAG CAC CGC	ACG GCG CCG GCA GCA GTT CGG TCA GGA CGC GTA GGG GAC TAA	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG GCG ACG GAG GTT GGA	GAA AGT TAC GTT CCA GTG CCC GAT TCA TTT CGA GCA GTG GTA TCT	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT AAA CTG AAA GCA	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC GCA CAA CGA GTG	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG GTG	AGG GCC CGC GAG TGG CAG CGG ATT CCA CCT CCG GAT CCA CGA	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GTC GAC TAA GTG ACT	CTT TGA TTC TAA CGC TGT CGT CGT TCT TGT CCT ATC GTT CCA CTG	CCA GTT TCG CAA AAT GGC CTG CTT CAA CCT AGC GGG AGC TGG	CAG GTT CTA CCG TCA TCA TCA TCA TCA CCG CGT GCT TAA TTA ATC	CAA GCG CCA GCA GCT CCT CAT GGC AAA CAG AGC CGC AAA GAA	TGG CGA TCG CAA ACG GGT CGT GCT GGA CAG AAA CAC
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG TAC TTT CTT ACC	GGT TGT CCA ACG TGC CCG CGC TTA CCA CTT GGG TGG CCC AGC TCC	GAA CAT GCA CGC GCG CTG GGG CTG TGC CGC TGG CGA AGT TTT AGA	ATA CCA CCG TGG CGT CCA CCA AAA GTT CGC CTT TGC AAG CAC CGC TTT	ACG GCG CCG GCA GCA GTT CGG TCA GGA GGC GAC TAA TTA	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG GCG ACG GAG GTT GGA GTC	GAA AGT TAC GTT CCA GTG CCC GAT TCA TTT CGA GCA GTG GTA TCT TGG	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT AAA CTG AAA GCA CGC	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC GCC CAA CGA GTG TCT	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG GTG CCC	AGG GCC CGC GAG TGG CAG CGG ATT CCA CCT CCG GAT CCA CGA AAC	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GTC GAC TAA GTG ACT TGA	CTT TGA TTC TAA CGC TGT CGT TCT TGT CCT ATC GTT CCA CTG GCT	CCA CGC GTT CCA AAT CCTG CTG CTG CCT AGC CCT AGC AGC TGG AAG	CAG GTT CTA CCG TCA TCG CCG CCG CCG CCG TCA CCG TTAA TTA ATC TCC	CAA GCG CCA GCA GCT CCT CAT GGC AAA CAG AGC CGC AAA GCG	TGG CGA TCG CAA ACG CCG GGT CGT GCT TGC CTA GGA CAG AAA CAC AAT
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG TAC TTT CTT ACC GCG	GGT TGT CCA ACG CCC CCC CCC CCC CCC CCC CCC AGC TCC TCC TTA	GAA CAT GCA CGC GCG CTT GGG CTG TGC CGC TGG CGA AGT TTT AGA CAA	ATA CCA CCG TGG CGT CCA CCA AAA GTT CGC CTT TGC AAG CAC CGC TTT GTA	ACG GCG CCG GCA GTT CTT CGG TCA GCG GCA GCC GTA GGG GAC TAA TTA	CCG GAT CTT CCA GGG GTT TTT CAT AGG GCG GAG GTT GGA GTC CAC	GAA AGT TAC GTT CCA GTG CCC GAT TCT CGA GCA GTG GTA TCT TGG AAA	CAT TAA AGG GAT GAC CCA GCG AAG TGC AAA CTG AAA GCA CGC GTT	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC GCC CAA CGA GTG TCT TTT	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG GTG CCC TAT	AGG GCC CGC GAG TGG CAG CGG ATT CCA CCT CCG GAT CCA CGA AAC GTT	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GTC GAC TAA GTG ACT TGA GAG	CTT TGA TTC TAA CGC TGT CGT TGT CCT CCT GCT CCA CTG GCT AAT	CCA CGC GTT TCG CAA AAT CCTG CTG CTT CAA CCT AGC GGG AGC TGG AAG ATT	CAG GTT CTA CCG TCA TCG CGA CCG CCT GCT TAA TTA ATC TCC TTT	CAA GCG CCA GCA GCT CCT CAT GGC AAA CAG AGC CGC AAA GCG TTG	TGG CGA TCG CAA ACG CCG GGT CGT GGT GGA CAG AAA CAC AAT ATG
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG TAC TTT CTT ACC GCG CGC	GGT TGT CCA ACG CCC CCC CCC CCC CCC CCC CCC CCC	GAA CAT GCA CGC GCG CCG CCG CTT GGG CGG CGG CGG AGT TTT AGA CAA TTA	ATA CCA CCG TGG CGT CCA CCA AAA GTT CGC CTT TGC AAG CAC CGC TTT GTA TTT	ACG GCG CCG GCA GTT CTT CGG TCA GAA GGG GAC TTA TTA TTA	CCG GAT CTT CCA GGG GTT TTT CAT AGG GAG GAG GTT GGA GTC CAC ATC	GAA AGT TAC GTT CCA GTG CCC GAT TCA GCA GTG GTA TCT TGG AAA GTT	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC AAA CTG AAA CTG AAA CCC GCT CGC	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC CAA CGA CGA CGA CGA	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG GTG CCC TAT AAG	AGG GCC CGC GAG TGG CAG CGG ATT CCA CCT CCG GAT CCA CGA AAC GTT AAG	CAG CAC CGC ATT CAA GAA AAA CAT GAC GAC GAC GAC TAA GTG GAC TGA GAG CGG	CTT TGA TTC CGC TGT CGT CGT TCT TCT CCT GTT CCTG GCT CCTG GCT AAT CGC	CCA CGC GTT TCG CAA AAT CCTG CTT CAA CCT AGC CCT AGC AGC TGG AAG ATT CAG	CAG GTT CTA CCG TCA TCG CCG CCG CCG TCA CCG GTT GCT TTA ATC TCC TTT GGT	CAA GCG CCA GCA GCT CCT CAT GGC AAA CAG AGC CGC AAA GCG TTG GGT	TGG CGA TCG CAA ACG CCG GGT CGT TGC CTA GGA CAG AAA CAC AAT ATG TTT
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG TAC TTT CTT ACC GCG CGC TTT	GGT TGT CCA ACG CCG CGC CGC CGC CGC CGC CGC CGC	GAA CAT GCA CGC GCG CCG CTT GGG CTG CGC TGC CGC TGC AGT TTA AGA TTA CAA	ATA CCA CCG TGG CGT CCA CCA AAA GTT CGC CTT TGC CGC CTT TGA	ACG GCG CCG CAC GCA GTT CTT CGG TCA GAA CGCA GAA GGA GAC TTA TTA TTG GAC	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG GCG GCG GCG GTC CAC ATC GGG	GAA AGT TAC GTT CCA GTG CCC GAT TCA TCA TCA GCA GTG GTA TCG AAA GTT CAA	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC AAA CTG AAA GCA CGC GTT CGC CAG	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC CAA CGA GTG TCT TCA CTG	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CCG GTG CCC TAT AAG GCT	AGG GCC CGC GAG TGG CAG CGG ATT CGA CCT CCG GAT CCA CGA AAC GTT AAG TAC	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GAC TGG GAC TCA GAG CGG TAT	CTT TGA TTC CGC TGT CGT CGT TCT TCT CCT CCT CCT	CCA CGC GTT TCG CAA AAT CCTG CTT CAA CCT AGC CCT AGC AGC AGC AGC CAG AGC CCT AGC CCT CCA CCT CCA CCT CCA CCC CCC CCC CC	CAG GTT CTA CCG TCA TCG CCG TCA CCG CCT GCT TCA CCG TTA ATC TTT GGT ACT	CAA GCG CCA GCA GCT CAT GGC AAA CAG CAG CGC AAA GCG TTG GGT GAT	TGG CGA TCG CAA ACG CCG GGT CGT TGC CTA GGA CAG AAA CAC AAT ATG TTT GAG
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG TAC TTT CTT ACC GCG CGC TTT GTC	GGT TGT CCA ACG CCG CGC CGC CGC CGC CGC CGC CGC	GAA CAT GCA CGC GCG CTT GGG CTG CGC TGC CGC TGC AGA TTA CAA CAA CAA GAA	ATA CCA CCG TGG CGT CCA AAA GTT CGC CTT TGC AAG CAC CGC TTT TGA GTG	ACG GCG CCG CAC GCA GTT CTT CGG TCA GAA CGC GTA GAA CGCA TTA TTG GAC CTT	CCG GAT CCA GGG GTT TTT CAT AGG GCG GCG GCG GTC CAC ATC GGG CAT	GAA AGT TAC GTT CCA GTG CCC GAT TCA TCA TCA GCA GTG GTA TCG AAA GTT CAA GTT CAA GTG	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC AAA CTG AAA GCA CGC CAG GCA	TAG TGA CTT CGG CGC TTT AGA CCC GCC GCC GCC CAA CAA CGA GTG TCT TCA CTG GGA	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CGG GGG GCC CGG GCC TAAG GCT GAA	AGG GCC CGC GAG TGG CAG CGG ATT CGA CCT CCA CGA CGA CGA CGA CGA CGA AAC TAC AAA	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GAC TGG GAC TAA GAG CGG TAT AGG	CTT TGA TTC TGA CGC TGT CGT TCT TGT CTC CTC GCT CCTG CCTG CCTG CTG	CCA CGC GTT TCG CAA GGC CTG CCT AGC CCT AGC CCT AGC AGC AGC CAC CAC CCC CCC CCA	CAG GTT CTA CCG TCA TCG CCG TCA TCG GCT TCC TTT GCT ACT CGG	CAA GCG CCA GCA GCT CAT GGC AAA CAG CAG CAC CAC AAA GCG TTG GGT GAT TGC	TGG CGA TCG CAA ACG CCG GGT CGT TGC CTA GGA CAG AAA CAA ATG TTT GAG GTC
CCT GAT CCA GCG GTT TCG CCA ACG ACG TAC TTT CCT CTT ACC GCG CGC TTT GTC AGA	GGT TGT CCA ACG CCA CCG CGC CGC CGC CGC CGC CGC	GAA CAT GCA CGC GCG CTG CGC TGG CGC TGG AGT TTA CAA TTA CAA TGT	ATA CCA CCG TGG CGT CCA AAA GTT CGC CTT TGC AAG CAC CGC TTT GTA TGA GTG GAT	ACG GCG CCG GCA GCA GCA GCA CCT CCG GAA CGC GAA CGC GGA GGA TTA TTA TTG GAC CTT ACA	CCG GAT CCA GGG GTT TTT CAT AGG GCG GCG GCG GTA GCA GTA GCA CATC GGA GGA GGA CATC	GAA AGT TAC GTT CCA GTG CCC GAT TCA TCA TCA GCA GTA GCA GTA GTA GTG CAA GTG TCA TCA TCA TCA TCA TCA TCA TCA TCA TCA	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC AAA CCA CCA CCA CCA CCA CCA CCA CCA C	TAG TGA CTT CGG CGC TGG CGC GCC GCC GCC GCC GCC	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG GCC CAC AAG CTT	AGG GCC CGC GAG TGG CAG CGG ATT CGA CCT CCA CGA CCA CCA CCA AAC AAC AAA CCT	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GAC TGG GTC GAC TAA GAG CGG TAT AGG CGC	CTT TGA TTC TGA CGC TGT CGT TCT TGT CCTC GTCA CCTG GCT CCTG GCT CCTG CCTG	CGC GTT TCG CAA GGC CTT CAA CCT AGC CCT AGC CCT AGC CCT GGC CAC CAC CCT GGC CCC CCC CCC CCC CCC CCC CCC CC	CAG GTT CTA CCG TCA TCA TCA TCA CCG CGT GCT TCA CCG GTT GCT TTA ATC TCT GGT ACT CCG ACT	CAA GCG CCA GCA GCT CAT GGC AAA CAG CAG CAG CAA GAA GCG GGT GGT GGT CGC	TGG CGA TCG CAA ACG CCG GGT CGT GGC TGC CTA GGA CAG AAA CAA TTT GAG GTC TAC
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG TAC TTT CTT ACC GCG CGC TTT GTC AGA CCG	GGT TGT CCA ACG TGC CCG CGC TTA CCA CTT GGG CCC TTA CAC TCC AGC TCA ACG TCC AGC TCA ACG TCC ACG CCC CCA CCA CCA CCA CCA CCA CCA C	GAA CAT GCA CGC GCG CTG GGG CTG TGC CGC TGG CGC TGG CGA AGT TTC CAA TTC	ATA CCA CCG TGG CGT CCA AAA GTT CGC CTT TGC AAG CAC CTT TGA GTT GTT GTT GAT GAC	ACG GCG CCG GCA GCA GCA CTT CGG TCA GAA CGC GAA GGA GGA TTA TTA TTA GAC CTT ACA TGC	CCG GAT CCA GGG GTT TTT CAT AGG GCG GCG GCG GCG GTC CAC GGA GGC GGA GGC GGC	GAA AGT TAC GTT CCA GTG CCC GAT TCA TCA TCA GCA GTA TCGA GCA GTA TCGA AGT CAA GTA CAA GTA CAA GTA CAA CCA CCC GTA CCA CCA CCA CCA CCA CCA CCA CCA CCA C	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT AAA CTG ACC CTG CTG CCC GCT CCG GCA GCA CCA CCA TGC AAG CCA TGC AAG CCA CCA CCA CCA CCA CCA CCA CCA C	TAG TGA CTT CGG CGC TGG CGC GCC GCC GCC GCC GCC	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG GTC CAC TAA GCT AAG CTT TGG	AGG GCC CGC GAG TGG CAG CGG ATT CCA CCT CCA CCA CCA CCA AAC AAC AAA CCT CCT	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GAC TGG GTC GAC TAA GAG CGC ACG CAC CAC	CTT TGA TTC CGC TGT CGT CGT CGT CCT CCT CCT CCT	CGC GTT CAA GGC CTT CAA GGC CTT CAA CCT GGC CAG GGC CAG GGC CTG GGC CAG GGC CCTG GGC CCTG GGC CCC CCC CCC CCC	CAG GTT CTA CCG TCA TCA TCA TCA CCG CGT GCT TCA CCG GCT TTA ATC TCT GGT ACT CCG ACT CCG ACT CCG CCG CCG CCG CCG CCG CCG CCG CCG C	CAA GCG CCA GCA GCT CCT CAT GGC AAA CAG AGC CGC AAA GCG GAT TGC GAC CGC GAG	TGG CGA TCG CAA ACG CGT GGT TGC CTA GGA CAG AAA CAC AAT GAG GTC TAC AAT
CCT GAT CCA GCG GTT TCG CCA ACG ACG CCG ACG TAC TTT CTT ACC GCGC TTT GTC AGA CGG TGC	GGT TGT CCA ACG TGC CCG CGC TTA CCA CTT GGG CGC TCA CCT TCA CAC TCA ACG TCC ACG TCA ACG TCC ACG CCA CCA CCA ACG CCA ACG CCA ACG CCA ACG CCA ACG CCA ACG CCA ACG CCA ACG CCA CCA	GAA CAT GCA CGC GCG CCG CTG GGG CTG CGC TGG CGA AGT TTC AGA CAA TTC CAA CAA CAA	ATA CCA CCG TGG CGT CCA AAA GTT CGC CTT TGC AAG CAC CTT TTA GTG GAT GAC CCA	ACG GCG CCG GCA GCA GCA CCT CCGG TCA GCA CGC GCA CGC GGA TTA TTA GAC CTT ACA CGC GGA	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG GCG ACG GAG GTC CATC GGA GGC ACG GGC ACG GGC ACG GGC ACG GGC ACA CAT	GAA AGT TAC GTT CCA GTG CCC GAT TCA TCA TCA TCA GCA GTG GTA TCGA GTG GTA TCGA GTG GTA TCA TCA TCA TCA TCA TCA TCA CCA CCC GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTT CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC CCA GTC GTC GTC GTC GTC CCA GTC GTC GTC GTC GTC GTC GTC GTC GTC GTC	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT AAA CTG ACC CTG CCG GCC CAG GCC CCA TGC ATT CCG TTA	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC GCA CGA CGA CGA CGA	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG GTG CCC TAT GAA CTT GAA CTT GAA	AGG GCC CGC GAG TGG CAG CGG ATT CCA CCT CCA CCA CCA CCA CCA CCA CCA	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GAC TGG GAC TGA GAG CGC ACG CGC ACG CGC ACG	CTT TGA TTC CGC TGT CGT CGT CCT CCT CCT CCT CCT	CCCA CGC GTT TCG CAA AAT GGC CTG CTT CAA CCT AGC CCT AGC CCTG AAG CCTG CAG CCC CAG CCC CCC CCC CCC CCC CCC CC	CAG GTT CTA CCG TCA TCA TCA TCA CCG CGT GCT TCA CCG GTT GCT TTA CCG TCA CCG CGT CCA CCG CGT CCA CCG CCG CCG CCG CCG CCA CCG CCG	CAA GCG CCA GCA GCT CAT GGC AAA CAG AGC CGC AAA GCG GGT TGG GGT TGC CGC GAG GGC	TGG CGA TCG CAA ACG CGT GGT TGC CTA GGA CAG AAA CAC AAT GAG GTC TAC GAC AAT AAA
CCT GAT CCA GCG GTT TCG CCA ACG ACG ACG TAC TTT CTT ACC GCG CGC TTC GCG CGC TTC GCG GTT	GGT TGT CCA ACG TGC CCG CGC TTA CCA CTT GGG TGC CCC TTA CAC TCA CCA TCA CTT ACG TCC ACG TCA CCA TCA CCA TCC CCA TCC CCA TCC TCC	GAA CAT GCA CGC GCG CTT GGG CTG CGC TGG CGA AGT TTT AGA CAA TTC CAA CCA	ATA CCA CCG TGG CGT CCA AAA GTT CGC CTT TGC AAG CAC CGC TTT GTA GTA GTA GAC GAC CCA	ACG GCG CCG GCA GCA GCA CCT CCGG TCA GCA CGC GCA CGC GCA CGC GCA CGC TCA GGC CGC CGC CCC CCG CCA CCG CCG CAC CGC CAC CGC CCC CC	CCG GAT CCA GGG GTT TTT TCT CAT AGG GCG ACG GCG ACG GGC ACG GGC ACG GGC ACG CAC CCA	GAA AGT TAC GTT CCA GTG CCC GAT TCA TCA TCA TCA GCA GTG GTA TCG AAT CGA GTG GTA TCG GCA GTG GTA TCG CCC GCA CCC CCC GTG CCC CCC GTG CCC CCC GTG CCC CCC	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC AAG CCA TGC ATT AAA GCA CGC GCC CAG GCC CAG GCA CTG CCA CCA TGC ACC CCA TGC ACC CCA CCA CCA CCA CCA CCA CCA CCA C	TAG TGA CTT CGG TGG CGC TTT AGA CCC GCC GCC GCC GCC GCC GCA CGA CGA	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CGG GTG CCC TAT GAC CCG GGC CTAT GGA CTA GCA	AGG GCC CGC GAG TGG CAG CGG ATT CCA CCA CCA CCA CCA CCA CCA CCA CCA	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GAC TGG GTC GAC TGA GAG CGG CAC CAC ACG CAC ACT CAA CAT CAA CAT CAA CAA CAA CAA	CTT TGA TTC CGC TGT CGT CGT CCT CCT CCT CCT CCT	CCCA CGC GTT TCG CAA AAT GGC CTG CTT CAA CCT AGC GGG AGC CAG GGG CAG GGC CAG GGC CAG CCTG CAA	CAG GTT CTA CCG TCA TCA TCA TCA TCA CCG CGT GCT TTA ATC TTT GGT ACT CGG ACT CGG CGC CGT CGG CGC CGC CGC CCG TCA	CAA GCG CCA GCA GCT CAT GGC AAA CAG CGC AAA GCG CGC CGC GAA GCG TTG GGT TGC CGC GAG GGC CAC CGC ACA CGC ACA CAC CCA CCA	TGG CGA TCG CAA ACG CGT GGT CGT TGC CTA GGA CAG AAA CAC AAT TTT GAG GTC TAC AAT TTA GAC CTC
CCT GAT CCA GCG GTT TCG CCA ACG ACG CCG TTT CCTT ACC GCG CGC TTT GTC ACG CGC TTT GTC ACG CGC TTT CTT	GGT TGT CCA ACG TGC CCG CGC TTA CCA CTT GGG CGC TTA CAC TCA CAC TCA CCA TTA CCA TCA CCA TCA CCA TCA CCA TCC CCA TCC CCA TCC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TGC CCA TCC TCC	GAA CAT GCA CGC GCG CTT GGG CTG CGC TGG CGA AGT TTT AGA CAA TTA CAG GAA TTT CAG GAA CCA CGC CGC CGC CGC CGC CGC CGC	ATA CCA CCG TGG CGT CCA AAA GTT CGC CTT TGC AAG CAC CGC TTT GTA TTT GTA GTA GCG GCC CCA	ACG GCG CCG GCA GCA GCA CTT CGG TCA GAA CGC GCA GGA GGC TAA TTA TTA TTA TTA GAC CCG GAC CGC GCA CGC CAC CGG CAC CGC CAC CA	CCG GAT CTT CCA GGG GTT TTT TCT CAT AGG GCG ACG GCG ACG GGT GGA GGA GGA GGA GGA GGA CCC	GAA AGT TAC GTT CCA GTG CCC GAT TCA TCA TCA TCA GCA GTG GTA TCT GAA GTA GTA GTA GTA GTA GTA GTA CCA GCA GCA GCA CCC GAT CCA CCA CCA CCA CCA CCA CCA CCA CCA C	CAT TAA AGG GAT GAC CCA GCG AAG CCA TGC ATT AAA CTG ACT CCA GCA CTG CAG GCA CCA TGC ATT AAA CTG ACC CCA TGC ACC CCA CCA CCA CCA CCA CCA CCA CCA C	TAG TGA CTT CGG CGC TTT AGA CCC GCC GCC GCC GCC GCC CAA CGA CGA	TGC TCA CGA CGC AGG GGT TCG CAC TGA ATT GAC CCG GGC CCG GGC CCC TAT GGA CTT GGG GCA ATA	AGG GCC CGC GAG TGG CAG CGG ATT CGA CCT CCG GAT CCA CGA AAC GTT AAG CCT CAA CCT CAA CCT AAA CCT CAA AAC	CAG CAC CGC ATT CAA GAA AAA CAT GAC TGG GAC TGG GTC GAC TGA GAG CGG TAA AGG CACG TCA AGG CAC TCA	CTT TGA TTC CGC TGT CGT CGT CCT CCT GTT CCT GTT CCT CC	CCCA CGC GTT TCG CAA AAT GGC CTG CTT CAA CCT AGC GGG CTT CAA CCT AGC GGG CAC CAG GGC CAC CAG GGC CAC CAG CAA AGC CCT CAA CCT CAA CCT CCAA CCT CCAA CTG CAA CTG CAA AAT GGC CTG CCAA CTG CCAA CTG CCAA CTG CCAA CCTG CCCG CCGC CCTG CCCG CCGC CCGC	CAG GTT CCA CCG TCA TCA TCA TCA TCA CCG CGT GCT TAA ATCC TTT ACCT TCA CCG CGT GCT TAA CCG CGT GCT CCA CCG CCG TCA CCG CCA CCG TCA CCG CCG	CAA GCG CCA GCA GCT CAT GGC AAA CAG CGC AAA GCG CGC CGC GAT CGC GAT CGC GAT CGC GAT CGC CGC GAC CGC CA CGC CA CCA CCA CCA C	TGG CGA TCG CAA ACG CGT GGT TGC CTA GGA CAG AAA CAC AAT TTT GAG GTC TAC AAT AAA CTC CT
	CGGT ACTC ACGG ACCG ACGG ACCG ACCG ACCG	CIGG AAT ACT GGA ATC AAG CAG GTT AGA CAA CGG TTC AGA CAA CGG TTC AGG CAG TCG ACG GGC AGG CTG ATG ACC AAG TCG ATC TCG CAA GCG ATG TCG ACT CCC GTG TTT ACG ACT CGT CTT CGG CAT GCT CTG CTG CAA CCT CTG CTG CAA CCT CTG CAT GGA TGC GCA TGC GCA TGC GCA TGC GCA TGC CAA CCT CCG ACT CTA CGT TAA TGG TGG GAA CAC TCC ATT TAT CTA ACA GCG TAC	CIG GAAT TGC ACT GGA TGG ATC AAG AGA CAG GTT CTC AGA CAA TCG CGG TTC TTT AGG CAG CGC TTC ATG CAG CGG TTC TTT AGG CAG CGC TCG ACG TTG GGC AGG ATC CTG ATG CAA ACC AAG CGA TCG CCA GGC GCG ATG CCT TCG ACT GTG CCC GTG ATA TTT ACG GTA ACG AGT TCT CAA CCT GCC CTT CGG AAT CTT CGG AAT CAT GCT GGA CTG CTG CCT CGG CAT CCA GGA GTG GGG AAT TGC GTT GCA TTA ATG GGG TGG TTT CCT GGC CCT GAA AAT CCT TAA CGT CGT TAA TGG CGC TAA TGC CGT TAA TGG CGC TAA TGC GAA CGG GAA CGA CAC TCC AGT ATT TAT GCC CTA ACA GCG GCG TAC CGT	CIGG AAT TGC CAG ACT GGA TGG CTT ATC AAG AGA CAG CAG GTT CTC CGG AGA CAA TCG GCT CGG TTC TTT TTG AGG CAG CGC GGC TCG ACG TTG TCA GGC AGG ATC TCC CTG ATG CAA TGC ACC AAG CGA AAC TCG ATC AGG ATG TCG CCA GGC TCA GCG ATG CCT GTG GCC AGG CTC GTC TCG ACT GTG GCC CCC GTG ATA TTG TTT ACG GTA TCG ACG AGT TCT TCT CAA CCT GCC ATC CTT CGG AAT CGT CTT CGG CCT GAG GGA ATA ATG AAT GGG TGG TTT TTC CCT GGC CCT GAG GAA AAT CCT GTT TAT CGT CGT ATC TAT CGT CGT ATC TAA TGC GTT GCG GAA AAT CCT GTT TAT CGT CGT ATC TAA TGG CGC GCA TGG GAA CGA TGC CAC TCC AGT CGC ATT TAT GCC AGC CAC TCC AGT CGC ATT TAT GCC AGC CAC TCC AGT CGC	CIGG AATTGCCAGCTGACTGGATGGCTTTCTATCAAGAGACAGGATCAGGTTCTCCGGCGGAGACAATCGGCTGCTCGGTTCTTTTTGTCAAGGCAGCGCGGCTATTCGACGTTGTCACTGGGCAGGCGCGGCTATTCGACGCGCGGCTATTCGACGCGATCCTGTCTGATGCAATGCGGCACCAAGCGAAACATCTCGCCAGGCGCCAGGGCGATGCCTGCTGCGCCGGTGATATTGCTGTTTACGGTATCGCCGACGAGTTCTTCTGAGCCCGTGGAAATCCCGCAACCTGCAATCCCGCAACCTGCAAATCGTCTGCTGCCTGAGGCTCTGCTGCCTGAGGCTCGGCATCCAGCATCTCTGGCGCCTGAGGCTCTGCGCCCTGAGGCTCGGCATCCTGCATCCCAATGCCCTGAGGCTCTGGCGCCTGAGGCTCTGGCGCCT	CIGGGCIIACAIGGCGGACGGAATTGCCAGCTGGGGACTGGATGGCTTTTCTGCATCAAGAAATCGGCTCTTAGACAATCGGCTGCTCTGAGACAATCGGCTGCTTTGAGACAGCAGTTTTTGTCAAGAAGGCAGCTCTTTTTGTCAAGAAGGCAGCTCTCTTCTAAGGGCAGGATCTCCTGTCATCTGAAGCGAAACATCGCATCGAAGCGAAACATCGCATCGAAGCGAAACAACAAGTCGAAGGCTGCTGCAAGGTCGAAGGCTGCTGCAAAGTCGAATGTGGCTGCACCGCCGGTGATATTGCTGCAGCCGACTGCGAATCTTCGGCAACCTGCAGCAAACAAGCTTCGGAATCGTCTTCGCCAACCTGCTGCAGCAGCGCAACCTGCTGCACCTGCACAACCTGCTGCACCTCATACGCATCCTGAACCTGCACAACCTGCTGCACCACA	GIG GGC 11ACAT GGC GAG CGCCGG AAT TGC CAG CTG GGG CGCACT GGA TGG CTT TCT TGC CGCATC AAG AGA CAG GAT GAC GGTCAG GTT CTC CGG CCG CTT GGGAGA CAA TCG GCT GCT CTG ATGCGG TTC TTT TTG TCA AGA CCGAGG CAG CGC GGC TAT CGT GGCAGG CAG CGC GGC TAT CGT GAGGGC AGG ATC TCC TGT CAT CTCCTG ATG CAA TGC GGC GGC TGCACC AAG CGA AAC ATC GCA TCGACC AAG CGA AAC ATC GCA TCGTCG ATC AGG ATG ATC TGC CGA ATATCG CCA GGC TCA AGG CGC GCC GCCACC AAG CGA AAC ATC GCA TCGTCG ATC AGG ATG ATC TGC CGA ATATCG CCA GGC TCA AGG CGC GCC GCAGCG ATG CCT GCT TGC CGA ATATCG ACT GTG GCC GCC TGG GTGCCC GTG ATA TTG CTG AAG AGCTTT ACG GTA TCG CCG CTC CCGACG AGT TCT TCT GAG CGG GACCAA CCT GCC ATC ACG AGA TTTCTT CGG AAT CGT TTT CCG GGACAA CCT GCC ATC ACG AGA TTTCTT CGG CAT CCA GGA AAC CAGGGA GTG GGG AGG CAC GAT GGCAAT TGC GTT GCG CTC ACT GCCACG ATA CCT GCG CTC ACT GCCGGA TTA ATG AAT CGG CAA ACGGGG TGG TTT TCC TTT TCA CCACCT GGC CCT GAG AGA GTT GCAGAA AAT CCT GTT TGA TGG TGGTAT CGT CGT ATC CCA CTA CCGGAA AAT CCT GTT TGA TGG TGGTAT CGT CGT ATC CCA CTA CCGTAT GG CGC GCA TTG CCC CAT TCACGG GAA CGA TGC CCT CAT TCACAC TCC AGT CGC CTT CCC GTTATA TGG CGC GCA TTG CCC CTT CCCATT TAT GCC AGC CAG CCA GACCAA CCC AGC CAG CCA GACCTA ACA CGG CGA TTT CCC GTT	GIGGGCITACATGGCGATACCITAACTGGATGCCAGGCTGGCCCCCAAATCAAGAGACAGGATGACGGTCGTCAGGTTCTCCGGCGCGCTGCGGGTCAGCTTTTTTTGTCAAGACCGACCAGGCAGCGCGGCTTTTTGTCAAGACCGCGGTTCTTTTTGTCAAGACCGACCAGGCAGCGCGGCTATCGTGGCTGGTCGACGTTGTCACTGAAGCCGGAAGGCAGGATCTCCTGTCATACCACCACGAAGCGAAACATCGCATCGACGACGAAGCGAAACATCGCAACGAAGTCGATGCCAGCCGCCGCATGCGCACCGATGCCTGCTGCAACGAAGTCACCGATGGCTGCTGCGGCATGCGCATCGATGGCTGCTGCGGCGGCATCACCGGCAGCTGCTGCGGCATCACCGATGGCTGCTGCGGCATCATCGATGGCTGCGGCGGCATCTCCGGCAGCTGCTGCG <td< td=""><td>GIGGGCIIACATGGCGATAGCIACIACCGGAATTGGCTTTCTTGCCGCCAAGGAATCAAGAGACAGGATGACGGTCCTTTCCAGGTTCTCCGGCCGCCTGGGTGCAGAAGACAATCGGCTGCTCTGAGGCCGCCGCGGTTCTTTTTGTCAAGACCGACCTGTAGGCAGCGCGGCTATCGTGGCAGAGGGGGCAGGTTGTCACTGAAGCGGCCATGGAGGCAGTTGTCACTGAAGCGCAGGGGGGGCAGGATCTCCTGTCATCTCACCTTGCTGAAGCCAGCCGCCGCATGCCCAGGCACCAAGCCAGCCGCCGCAACCTGGCCGGCAATCAGGAACATCTGGAAGAGCCCGGCGATGCCAGCCGCCGCAAGCCCGGGGGCGATGCCTGCTGCCGCCGCGGGGGGGGCGATGCCTGCTGCCGCCGCGGGGGCGATGCCTGCTGCGGCGGCGGGGCCGGCAATCTCTGCAAGC<!--</td--><td>CIG AAT TGC CAG CTG GGG CGC CCT CTG GTA ACT GGA TGG CTT TCT TGC CGC CAA GGA TCT ATC AAG AGA CAG GAT GAC GGT CGT TTC GCA CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC AGA CAA TCG GCT GCT CTG ATG CCG CCG TGT CGG TTC TTT TTG TCA AGA CCG ACC TGT CCG AGG CAG CGC GGC TAT CGT GGC TGG CCA CGA TCG ACG TTG TCA CTG AAG CGG GAA GGG ACT GGC AGG GTC TCC TGT CAT CTC ACC TTG CTG ATG CAA TGC GGC GGC TGC ATA CGC TTG ACC AAG CGA AAC ATC GAA CGA AGC GAG CAC TCG ATG CAA TGC GGC GGC TGC ATC CTC ACC TTG CCG ATG CAA TGC GGC GGC TGC ATA CGC TTG ACC AAG CGA AAC ATC GCA TCG AGC GAG CAC TCG ATC AGG ATG ATC TGG ACG AAG AGC ATC TCG CAA GGC TCA AGG CGC GCA TGC CCG ACG GCC ATG CCT GCT TGC CGA ATA TCA TGG TGG TCG ACT GTG GCC GCC TGG GTG TGG CGG ACC CCC GTG ATA TCG CGG CTC CCG ATT CGC AGC AGC AAT TCT TCT GAA GAG AGC TTG GCG GCG TTT ACG GTA TCG CCG CTC CCG ATT CGC AGC AGC AGT TCT TCT GAG GGG GAC TTC GGG GTT CAA CCT GCC ATC ACG AGA TTT CGA TCC CAC CTT CGG AAT CGT TTT CGG CGA CCC CGG CTG CTT CGG AAT CGT TTT CGG CGA CCC CGG CTG CTT CGG AAT CGT TTT CCG GGA CGC CGG CTG CTT CGG AAT CGT TTT CGC GGA CGC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTG CTG CCT GAG GCT GGA CGA CGC CGG CTG CAA CCT GCC ATC ACG AGA TTT CGA TCC CAC CTG CTG CCT GAG GCT CGA CAG CGC CGG CTG CAT GCT GGA ATC CTT TC CGC CAA CCC CGG GAA CGG CAT CCA GGA AAC CAG CAG CGC CGC TTT CCA GGA GTG GGG AGG CAC GAT GGC CGC TTT CCA GGA TTA ATG AAT CGG CCC ACT GCC GGC TAT CCC GGA GTG GGG AGG CAC CA CTA CCG GGG GAG CTT CGT CGT ATC CCA CTA CCG AAG CGG CGG TAT CGT CGT ATC CCA CTA CCG AAG CGG GGG TAT CGT CGT ATC CCA CTA CCG AGA TAT CCG TAA TGG CGC GCA TTG CCC CTA TCA GCA AAC CGG CTA AAT CCT GTT TGA TGG CCC AAG CGC CGA TCT TGG GAA CGA TCC CCT CAT TCA GCA AAC CGA CGC TAA CAG CGC CCA CTT CCC GTT CCG CTA TCT GGG AAA CGA TGC CCT CAT TCA GCA AAC CAA GCG CTA ACA CGG CGA TTG CCG CTA CCG CTA TCT GGG TAC CGT CCT CCT CTT CCC GTT CCG CTA TCG ATT TAT GCC AGC CAT CCC GTT CCC GTT CCG CTA TCT</td><td>GIGGACTIACAGCAGCGGCAGCAGCAGCAGACTGGATGGCTTGTAAGGACTGGATGGCTTTTTTGCCGCCAAGGATGCGATTGCGGATGCGAGTGCGAGTGCGAGTGCGAGTGCGAGGGCTATAGAAGAAGGCGTTTTGGCGGCTGTCCGGCGGGCTATAGAAGAGGCGGCTGTCCGGGCGGCTGTCCGGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGAGCCGGA<!--</td--><td>GIGGACTIAGACGAGCACCCTCTGGAGTTGCACGGATGGCAGCTTTTCTTGCCGCCAAGGATTGGAGGGTTTGCAGGTTCTCCGGCGCCCTGGAGGAGGCTTTCGAGGCTTGCAGGTTCTCCGGCGCCCTGGCGGCTATTCGGGCGGCTATTCGAGACAATCGGCTGCTGCTGGCGGCTATTCGGGCGGCTATTCGAGACAATCGGCTGCTGCTGGCGGCTATTCGGGCGGCGGCTATTCGAGGTTTTTGGCCGCGGCGTGCTCGCCCGGGG</td></td></td></td<> <td>GIG GGA TIA CAI GGC GAG GGC CCT CTG GTA AGG TTG GGA ACT GGA TGG CTT TCT TGC CGC CCA GGA TCT GAT GGC GCA ATC AAG AGA CAG GAT GAC GGT CGT TTC GCA TGC TTG AAC CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC TAT TCG GCT AGA CAA TCG GCT GCT CTG ATG CCG CCG TGT TCC GGC TGT CGG TTC TTT TTG TCA AGA CCG ACC TGT CCC GGC GCT AGG CAG CGC GGC TAT CGT GGC GAC GGA CGG GCG TTC TCG ACG TTG TCA CTG AAG CCG ACC TGT CCC GGC GCG TTC GCG AGG ATC TCC TGT CAT CTC ACC TTG CCC GCG GC TTC ACC AGG CAG CGA AAC ATC GCA TCC ACC TTG CCC GCG CTA ACC AAG CGA AAC ATC GCA TCG AGC GAA GGC ACC GTA CTC GGA TCG ATG CAA TGC GGC GGC TGC ATA CGC TGG ACA GGG CAC GCG ATG CAA GC GGC GCA TGC CCG ACG GCG ACG CCG AGG CTC AAG CCG ACA GCG GCA AGC GCA AGC CCG ATG CAA GC GGC GCA TGC CCG ACG GCG ACG CCG ATG CAA GC GGC CGA TGC CCG ACG GCG AGG CTG CCG ACG GTC AAG CGG GCA TGC CGG ACC GCA ATG GCC TCG ACT GTG GCC GGC TGG GTG TGG CGG ACC GCT ATC AGG CCC GTG ATA TTG CTG AAG AGC TTG GCG ACC GCA ATG GCC TTT ACG GTA TCG CGG CCA TTC CGG ACC GCA ATG GCC TTT ACG GTA TCG CGG CCA TTT CGA GCG GCG AAT GGG CTG TTT ACG GTA TCG CGG CCA CTC GGG GTT CGA AAT GAC CCA CCT GCC ATC CGG GCA TTT CGA GCG GCA ATT GGC CTT CTT CGG AAT CGT TTT CCG GGA CGC CGC CTT CCA CGC CCT TC CTT GGG AAT CGT TTT CCG GGA CGC CGG CTA GAT CCC CAA CCT GCC ATC ACG AGA TTT CAA CGC GCG CTA TCC AGG CCA CT GCC ATC ACG AGA CTT CGG GGT CGA AAT GAC CCAA CCT GCC ATC CGG GAC CGC CGG CTA TCC ACC CCC CGG CAT CAA GGA AAC CAG CAG CGC CGG CTA TCC ACC CGG CAT CAA GGA AAC CAG CAG CCC CGG GAG AGA CAC TCC CAT GCT GGA AGT CTT TCA CAG GCG CGC TTT CCA GCG CAT CCA GGA GTG GGG AGG CAC CAA CAG CGC CGC TTT CCA GCG CAT CCA GGA GTG GGG AGG CAC CAA CGG CGC CTA TCC GGG CAA ACA GCT CCT GGC CCT GAA GAA GTG GCA CGC GCG CTA TCC GCG CAA CAA TTC AAT TGC GTT TCC TTT TA CCA GTG GGG CAA ACA GCT GCA AAT CCT GTT TGA TGG TGG TTA ACG GCG GGA TAT AAC TAT ATG GCT GCC CTT CCC CTT CCG CTA TCG GCC ACA CCA CAA CGC TAA AGA CGA TGC CCT CAT TCA GCA GAC ACC ACG CGG CTA ACA AGC CGC CTT CCC CTT CCG CTA CCG CCA ACG CGA CCA GAT TT ATT TAT GCC AGC CAG CCA CGC CCA ACC AATG CGA CAA</td> <td>GIG GAT TIG CAG GGG GGG CGC CT CTG GTA AGG TTG GGA AGG ACT GGA TGG CTT TCT TGC CGC CAA GGA TCT GAT GGC GCA GGG ATC AAG AGA CAG GAT GAC GGT CGT TTC GCA TGC TTG AAC AAG CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC TAT TCG GCT ATG AGA CAA TCG GCT GCT CTG ATG CCG CG TGT TCC GGC TGT CAG CGG TTC TTT TTG TCA AGA CCG ACC TGT CCG GTG CCC TGA ATG AGG CAG CGC GGC TAT CGT GAT GGC GGC CAC CGA CGG GGC TTC CTT TCG ACG TTG TCA CTG AAG CGG GAA GGG ACT GGC TGC CTT GGC AGG ATC TCC CTG ATG CTC ACC TTG CTC GGC GAG AAG CTG ATG CAA TGC GGC GGC TGC ATA CGC TTG ATC CGG CTA CCT ACC AAG CGA AAC ATC GCA CGA AGG GAC CAC GTA CTC GGA TGG TCG ATG CAA AGA CCG GCA TGC ATA CGC TTG ATC CGG CTA CCT ACC AAG CGA AAC ATC GCA CGA AGA GAG CAC GTA CTC GGA TGG CGG ATG CCT GCT TGC CGA ATA CGC AGG GGC AGG ATC GG GCG AGG ATC TCG CGC TGG CAC ATA CGC GTA CTC GGA CGG CGG ATG CCT GCT TGC CGA ATA TCA TGG TGG AAA ATG GCC GCT CGG ACT AGG ATG TGG CGG GCG TGC GAT CGG CGC AGG AGG ATC AGG ACC GTG ATA TTG CTG AAG AGC TTG GCG ACC GCA TCC GCC TTC TAC AGG ATA TTG CTG AAG AGC TTG GGG AAT GGC CTT CTT ACG GTA TTG CTG AAG AGC TTG GGG GCG AAT GGC CTT CTT ACG GTA TTG CTG AAG AGC TTC GGG GCG AAT GGC CTT CTT ACG ATT ATG CTG AAG AGC TTC GGG GCG AAT GGC CTT CTT ACG AGT TCT TCT GAG CGG GAC CTC TGG AAC ATC GCC TTC TA CTT CGG AAT CTT TTT CCG GAA CTC CGG GAT CGA TCC CCT CCA CAA CCT GCC ATC ACG AGA TTT CGA AGC CGG ATT CGA AAC CCC CTG CTG CCT GAG GTT TTT CCG GGA CCT CGG GAT GAT CCT CCA CAA CCT GCC ATC ACG AGA CAC CCC CGG GTT CGA ACC CCT CCA CAA CCT GCA GAG AAC CAG GAG CCT CGC GGA GTT CCA CCC CGG GGA GTG GGG AGG CAC CAT CCG GGA CTA TCC GGG CAT CCA CCC CGG CTG CTG GAG GCT GGA CGA CCC CGC GTT TTG CGG GAA AAC CTT GGC GGA GTG GGG AGG CAC CAT CCG GGA CTA TCC GGG CAA CCA GCC GGC TTT GGG GGG AGG CAC CAA CGG CGG CTA TCC GGG CAA CCA GCC GGC TTT GGG GGG AGA GCT GGA CAA CCG GGG GGA AAA CCT CCA CGC GGG CCC AACG CCA ACG CCA GCA GCG GCA ACA GCC GGA TAT AAC ATG GGG CCC GAA TTG CCG CAA CCG CCA CAC CCA CAA CGC GGA TAT AAC ATG GGG AGG CCG CAA TTG CCG CAA CCA GCA GCA CAA CAA GCC GAA CTA ACA GCC GCA TTG CCA CTA C</td> <td>GIG GAT TIGC CAG CTG GGG CGC CCT CTG GTA AGG TTG GGA AGC CCT ACT GGA TGG CTT TCT TGC CGC CGC TATG GTA AGG TTG GGA AGC CCT ACT GGA TGG CTT TCT TGC CGC CGC TTTC GCA TGC TTG AAC AAG ATG CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC TAT TCG GCT ATG ACT AGA CAA TCG GCT GCT CTG ATG CCG CCG TGT TCC GGC TGT CAG CGC CGG TTC TTT TTG TCA AGA CCG ACC TGT CCG GTG CCC TGA ATG ACC AGG CAG CGC GGC TAT CGT GGC TGG CCA CGA CGG GCG TTC CTT GCG GGC AGG TTG TCA CTG AAG CCG ACC TGT CCG GCG GCC TAT TGG GCG GGC AGG ATC TCC TGT CAT CTC ACC TTG CTC CTG CCG AGA AAG TAT CTG ATG CAA TGC GGC GGC TGC ATA CTC ACC TTG CTC CTG CCG AGA AAG TAT CTG ATG CAA TGC GGC GGC TGC ATA CTC ACC TTG CTC CTG CGG AGA AGG TAT CTG ATG CAA TGC GGC GGC TGC ATA CTC AGC CTG ATC CGG GGA TGG AAG ACG GAA ACA ATC GCA TCG AGC GAG CAC GTA CTC GGA TGG AAG CCG ATC AGG ATC TCC TGC CGA ATA CGC TTG ATC CGG CAA GGA ACG ATC AGG ATA CTC GGA CGA AGG AGC GAC GTA CTC GGA TGG AAG CCG CCA GGC TCA AGG CGC GCA TGC CCG ACG GCA AGG ATC TCG GCG CGG ATG CCT GCT GC GGA ATA TCA TGG TGG AAA ATG GCC GCT TT CG ACT GTG GCC GCC TCG GTG TGT GGG GCG AAT GGG CTT ACC GGC CTT ACG GTA TCG CCG CTC CCG ATT CGG GGG ACC GCT ATC AGG ACA TAG CCC GTG ATA TTG CCG AGC ACC TCT GGG GCG AAT GGC CTT CTA TCC ACG AGT TCT TCT GAC GGA CAC TCT GGG GTT CGA AAT GAC CGA CCA CCA CCT GCC ATC ACG AGA ATT CGA TTC CAC CGC CCT CTT ATC ACG AGT TCT TCT CGC GGA CAC CCC CGG GCT GAA ATG GAC CCA CCA CAC CCT GCC ATC ACG AAA CCG CCC CCG GGG CTA CCA TGC CCC CCG CAT GCT GGC ATT CTC CGC CAC CCC CGG GCT CGA AAT GAC CCA CCA CCG GAA TC CA GGA AAC CAG CAG CAC CCC CGG GCA TTC CA TGC CCC CCC GGA AGT CCA GGA AAC CAG CAG CGG CTA TCC CGA CAA TCC CCC CGC GAG CTT CGG CAT CCA GGA AAC CAG CAG CGC CGG CTA CCA ACC CCC CGC GAA AAT CCT GTT TGC TTT CGC CAC CCC CGC GGG CAA ACA ATG CCC CCC GGA AAT CCT GTT TGC TTT CGC CAC CCC CGG GGA AAA CCA GCC CGC GAA ATG CCC GAA AAT CCT GTT TGC TGC CCA CCC CGC GGG GAA AGG CGT TTT GCC TAT CGG TAT ATA ATG CGC CCA CAC CGC GGG CAA ACA GCC GGA CAA CCA CTA CGT CGT ATC CCA CTA CCG AGA CGG GGA CAA ACA CAC CTA GGC GGA TTT CCC CAT TCA CGG C</td> <td>GIG GAC TIG CAG CGA GAC GAC ACC TIG ACT GGA AGC TIG GGA AGC CCT GCA ACC GAA ACC CAA ACT GGA TGG CTT TCT TGC CGC CAA GGA TCT GAT GGC GCA GGG AAC CAA ATC AAA ACAA AGG ACA GAT GAC GGT CGT TTC GCA TGC TG AAC AAA ATG GAT CAA ATG GCT GCT CTG ATG GCA GGA GGC TAT TCC GGC TGT ACA AAG ATG GAC GAC TTT GG GCA GGC TGT TCC CGG CTG TTT CC GGC TGT CAC GGA GGC TAT TTT TTT TTT TTT TTT TTA CAA GA CGA ACC TGT CCG GTG TCC TGA ATG ACC AGA AGG CAA TTT TTT TTT TTT TTA CAA GA CGA ACC TGT CCG GCG TGT TCC CTG ATG ACC AGA GGC TAT TTT TTT TTT TTT TTA CAA GA CGA ACC TGT CCG GCG TGT TAT CGG CGC AGG ACC GTG TCC TTA TGG GCG AAG GGC ACC GTG TCC TTA TGG GCG AAG GGC AGG ATC TCC TGT CAC TGA ACG AAG CGG AAG GGC AGC AGG ATC TCC TGT CAC TGA ATG CGC AGC AGA AGA CAC GTA CTC GGC AGA AAG TAT CCA ACC AGG CAG CAC GTA CTC GGA AAA AG CCC TCG ATC AGG AAA CAC GCA CGA CGC GCA TGC CGA CGC GCA TGC CGA AGA AGG CAC GTA CTC GGA AGA AGG CCG GCG CGA GGC GCA TGC CGA ACG GGA AGG ATC TCG GCG AGA AGA CCG TCG ACG AGG ATG CCG GCT GCT AGG CGG GCG ATG TGG GGA ACA TGA GGA AGA AGG CTT CTG TGG CGA ATA TCA TGG TGG AAA ATG GCC GCT TTT CTG TCG GCG ATG TCG CGC ACC GCT ATC AGG ACA TTA CGC TCC TTT ACC GTG ATA TTG CTG AAG AGC TTG GCG ACC GCT ATC AGG ACA TAG CGT TCC TTT ACG GTA TTT CGA GCA GCT CCC AGC CAT CCC AGC CAC CCC CTT TCT ATC AGC ACC AGC CAC TTC CTG TTT CGG AAA ATG CCG CCT TCT ATC ACC AGC CAC TTC CTG TTT CGG AAA TGC CCT TCT ATC ACC AGC CAC TTC CAC AGC CAC ACC CCC GG ATT CCA CCA CGA CCA AGC CAC TTC CAC AGC CAC ACC CCC CGG CTT CTA ACC AGC CCC CGG AAT CCC CCC CGA ATT CCA GCA ACA AAG CTT CGA GAC ATC CAC GAA AAT CCT CCA CGA CAC AAC CAC ACC CCC CGG CTT CAA CCA GCC CCC CGA AAT CCA GCA CAC ACC CCC CGA AAT CCA GCA CAC ACC CCC CGA AAT CCA GCA CCA ACC CCC CGA AAT CCA GCA CCA ACC CCC CGA AAT CCA GCA CCA ACC AGC CCA ACC A</td>	GIGGGCIIACATGGCGATAGCIACIACCGGAATTGGCTTTCTTGCCGCCAAGGAATCAAGAGACAGGATGACGGTCCTTTCCAGGTTCTCCGGCCGCCTGGGTGCAGAAGACAATCGGCTGCTCTGAGGCCGCCGCGGTTCTTTTTGTCAAGACCGACCTGTAGGCAGCGCGGCTATCGTGGCAGAGGGGGCAGGTTGTCACTGAAGCGGCCATGGAGGCAGTTGTCACTGAAGCGCAGGGGGGGCAGGATCTCCTGTCATCTCACCTTGCTGAAGCCAGCCGCCGCATGCCCAGGCACCAAGCCAGCCGCCGCAACCTGGCCGGCAATCAGGAACATCTGGAAGAGCCCGGCGATGCCAGCCGCCGCAAGCCCGGGGGCGATGCCTGCTGCCGCCGCGGGGGGGGCGATGCCTGCTGCCGCCGCGGGGGCGATGCCTGCTGCGGCGGCGGGGCCGGCAATCTCTGCAAGC </td <td>CIG AAT TGC CAG CTG GGG CGC CCT CTG GTA ACT GGA TGG CTT TCT TGC CGC CAA GGA TCT ATC AAG AGA CAG GAT GAC GGT CGT TTC GCA CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC AGA CAA TCG GCT GCT CTG ATG CCG CCG TGT CGG TTC TTT TTG TCA AGA CCG ACC TGT CCG AGG CAG CGC GGC TAT CGT GGC TGG CCA CGA TCG ACG TTG TCA CTG AAG CGG GAA GGG ACT GGC AGG GTC TCC TGT CAT CTC ACC TTG CTG ATG CAA TGC GGC GGC TGC ATA CGC TTG ACC AAG CGA AAC ATC GAA CGA AGC GAG CAC TCG ATG CAA TGC GGC GGC TGC ATC CTC ACC TTG CCG ATG CAA TGC GGC GGC TGC ATA CGC TTG ACC AAG CGA AAC ATC GCA TCG AGC GAG CAC TCG ATC AGG ATG ATC TGG ACG AAG AGC ATC TCG CAA GGC TCA AGG CGC GCA TGC CCG ACG GCC ATG CCT GCT TGC CGA ATA TCA TGG TGG TCG ACT GTG GCC GCC TGG GTG TGG CGG ACC CCC GTG ATA TCG CGG CTC CCG ATT CGC AGC AGC AAT TCT TCT GAA GAG AGC TTG GCG GCG TTT ACG GTA TCG CCG CTC CCG ATT CGC AGC AGC AGT TCT TCT GAG GGG GAC TTC GGG GTT CAA CCT GCC ATC ACG AGA TTT CGA TCC CAC CTT CGG AAT CGT TTT CGG CGA CCC CGG CTG CTT CGG AAT CGT TTT CGG CGA CCC CGG CTG CTT CGG AAT CGT TTT CCG GGA CGC CGG CTG CTT CGG AAT CGT TTT CGC GGA CGC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTG CTG CCT GAG GCT GGA CGA CGC CGG CTG CAA CCT GCC ATC ACG AGA TTT CGA TCC CAC CTG CTG CCT GAG GCT CGA CAG CGC CGG CTG CAT GCT GGA ATC CTT TC CGC CAA CCC CGG GAA CGG CAT CCA GGA AAC CAG CAG CGC CGC TTT CCA GGA GTG GGG AGG CAC GAT GGC CGC TTT CCA GGA TTA ATG AAT CGG CCC ACT GCC GGC TAT CCC GGA GTG GGG AGG CAC CA CTA CCG GGG GAG CTT CGT CGT ATC CCA CTA CCG AAG CGG CGG TAT CGT CGT ATC CCA CTA CCG AAG CGG GGG TAT CGT CGT ATC CCA CTA CCG AGA TAT CCG TAA TGG CGC GCA TTG CCC CTA TCA GCA AAC CGG CTA AAT CCT GTT TGA TGG CCC AAG CGC CGA TCT TGG GAA CGA TCC CCT CAT TCA GCA AAC CGA CGC TAA CAG CGC CCA CTT CCC GTT CCG CTA TCT GGG AAA CGA TGC CCT CAT TCA GCA AAC CAA GCG CTA ACA CGG CGA TTG CCG CTA CCG CTA TCT GGG TAC CGT CCT CCT CTT CCC GTT CCG CTA TCG ATT TAT GCC AGC CAT CCC GTT CCC GTT CCG CTA TCT</td> <td>GIGGACTIACAGCAGCGGCAGCAGCAGCAGACTGGATGGCTTGTAAGGACTGGATGGCTTTTTTGCCGCCAAGGATGCGATTGCGGATGCGAGTGCGAGTGCGAGTGCGAGTGCGAGGGCTATAGAAGAAGGCGTTTTGGCGGCTGTCCGGCGGGCTATAGAAGAGGCGGCTGTCCGGGCGGCTGTCCGGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGAGCCGGA<!--</td--><td>GIGGACTIAGACGAGCACCCTCTGGAGTTGCACGGATGGCAGCTTTTCTTGCCGCCAAGGATTGGAGGGTTTGCAGGTTCTCCGGCGCCCTGGAGGAGGCTTTCGAGGCTTGCAGGTTCTCCGGCGCCCTGGCGGCTATTCGGGCGGCTATTCGAGACAATCGGCTGCTGCTGGCGGCTATTCGGGCGGCTATTCGAGACAATCGGCTGCTGCTGGCGGCTATTCGGGCGGCGGCTATTCGAGGTTTTTGGCCGCGGCGTGCTCGCCCGGGG</td></td>	CIG AAT TGC CAG CTG GGG CGC CCT CTG GTA ACT GGA TGG CTT TCT TGC CGC CAA GGA TCT ATC AAG AGA CAG GAT GAC GGT CGT TTC GCA CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC AGA CAA TCG GCT GCT CTG ATG CCG CCG TGT CGG TTC TTT TTG TCA AGA CCG ACC TGT CCG AGG CAG CGC GGC TAT CGT GGC TGG CCA CGA TCG ACG TTG TCA CTG AAG CGG GAA GGG ACT GGC AGG GTC TCC TGT CAT CTC ACC TTG CTG ATG CAA TGC GGC GGC TGC ATA CGC TTG ACC AAG CGA AAC ATC GAA CGA AGC GAG CAC TCG ATG CAA TGC GGC GGC TGC ATC CTC ACC TTG CCG ATG CAA TGC GGC GGC TGC ATA CGC TTG ACC AAG CGA AAC ATC GCA TCG AGC GAG CAC TCG ATC AGG ATG ATC TGG ACG AAG AGC ATC TCG CAA GGC TCA AGG CGC GCA TGC CCG ACG GCC ATG CCT GCT TGC CGA ATA TCA TGG TGG TCG ACT GTG GCC GCC TGG GTG TGG CGG ACC CCC GTG ATA TCG CGG CTC CCG ATT CGC AGC AGC AAT TCT TCT GAA GAG AGC TTG GCG GCG TTT ACG GTA TCG CCG CTC CCG ATT CGC AGC AGC AGT TCT TCT GAG GGG GAC TTC GGG GTT CAA CCT GCC ATC ACG AGA TTT CGA TCC CAC CTT CGG AAT CGT TTT CGG CGA CCC CGG CTG CTT CGG AAT CGT TTT CGG CGA CCC CGG CTG CTT CGG AAT CGT TTT CCG GGA CGC CGG CTG CTT CGG AAT CGT TTT CGC GGA CGC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTT CGG AAT CGT TTT CGC GAA CCC CGG CTG CTG CTG CCT GAG GCT GGA CGA CGC CGG CTG CAA CCT GCC ATC ACG AGA TTT CGA TCC CAC CTG CTG CCT GAG GCT CGA CAG CGC CGG CTG CAT GCT GGA ATC CTT TC CGC CAA CCC CGG GAA CGG CAT CCA GGA AAC CAG CAG CGC CGC TTT CCA GGA GTG GGG AGG CAC GAT GGC CGC TTT CCA GGA TTA ATG AAT CGG CCC ACT GCC GGC TAT CCC GGA GTG GGG AGG CAC CA CTA CCG GGG GAG CTT CGT CGT ATC CCA CTA CCG AAG CGG CGG TAT CGT CGT ATC CCA CTA CCG AAG CGG GGG TAT CGT CGT ATC CCA CTA CCG AGA TAT CCG TAA TGG CGC GCA TTG CCC CTA TCA GCA AAC CGG CTA AAT CCT GTT TGA TGG CCC AAG CGC CGA TCT TGG GAA CGA TCC CCT CAT TCA GCA AAC CGA CGC TAA CAG CGC CCA CTT CCC GTT CCG CTA TCT GGG AAA CGA TGC CCT CAT TCA GCA AAC CAA GCG CTA ACA CGG CGA TTG CCG CTA CCG CTA TCT GGG TAC CGT CCT CCT CTT CCC GTT CCG CTA TCG ATT TAT GCC AGC CAT CCC GTT CCC GTT CCG CTA TCT	GIGGACTIACAGCAGCGGCAGCAGCAGCAGACTGGATGGCTTGTAAGGACTGGATGGCTTTTTTGCCGCCAAGGATGCGATTGCGGATGCGAGTGCGAGTGCGAGTGCGAGTGCGAGGGCTATAGAAGAAGGCGTTTTGGCGGCTGTCCGGCGGGCTATAGAAGAGGCGGCTGTCCGGGCGGCTGTCCGGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGCGGAGGAGCCGGA </td <td>GIGGACTIAGACGAGCACCCTCTGGAGTTGCACGGATGGCAGCTTTTCTTGCCGCCAAGGATTGGAGGGTTTGCAGGTTCTCCGGCGCCCTGGAGGAGGCTTTCGAGGCTTGCAGGTTCTCCGGCGCCCTGGCGGCTATTCGGGCGGCTATTCGAGACAATCGGCTGCTGCTGGCGGCTATTCGGGCGGCTATTCGAGACAATCGGCTGCTGCTGGCGGCTATTCGGGCGGCGGCTATTCGAGGTTTTTGGCCGCGGCGTGCTCGCCCGGGG</td>	GIGGACTIAGACGAGCACCCTCTGGAGTTGCACGGATGGCAGCTTTTCTTGCCGCCAAGGATTGGAGGGTTTGCAGGTTCTCCGGCGCCCTGGAGGAGGCTTTCGAGGCTTGCAGGTTCTCCGGCGCCCTGGCGGCTATTCGGGCGGCTATTCGAGACAATCGGCTGCTGCTGGCGGCTATTCGGGCGGCTATTCGAGACAATCGGCTGCTGCTGGCGGCTATTCGGGCGGCGGCTATTCGAGGTTTTTGGCCGCGGCGTGCTCGCCCGGGG	GIG GGA TIA CAI GGC GAG GGC CCT CTG GTA AGG TTG GGA ACT GGA TGG CTT TCT TGC CGC CCA GGA TCT GAT GGC GCA ATC AAG AGA CAG GAT GAC GGT CGT TTC GCA TGC TTG AAC CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC TAT TCG GCT AGA CAA TCG GCT GCT CTG ATG CCG CCG TGT TCC GGC TGT CGG TTC TTT TTG TCA AGA CCG ACC TGT CCC GGC GCT AGG CAG CGC GGC TAT CGT GGC GAC GGA CGG GCG TTC TCG ACG TTG TCA CTG AAG CCG ACC TGT CCC GGC GCG TTC GCG AGG ATC TCC TGT CAT CTC ACC TTG CCC GCG GC TTC ACC AGG CAG CGA AAC ATC GCA TCC ACC TTG CCC GCG CTA ACC AAG CGA AAC ATC GCA TCG AGC GAA GGC ACC GTA CTC GGA TCG ATG CAA TGC GGC GGC TGC ATA CGC TGG ACA GGG CAC GCG ATG CAA GC GGC GCA TGC CCG ACG GCG ACG CCG AGG CTC AAG CCG ACA GCG GCA AGC GCA AGC CCG ATG CAA GC GGC GCA TGC CCG ACG GCG ACG CCG ATG CAA GC GGC CGA TGC CCG ACG GCG AGG CTG CCG ACG GTC AAG CGG GCA TGC CGG ACC GCA ATG GCC TCG ACT GTG GCC GGC TGG GTG TGG CGG ACC GCT ATC AGG CCC GTG ATA TTG CTG AAG AGC TTG GCG ACC GCA ATG GCC TTT ACG GTA TCG CGG CCA TTC CGG ACC GCA ATG GCC TTT ACG GTA TCG CGG CCA TTT CGA GCG GCG AAT GGG CTG TTT ACG GTA TCG CGG CCA CTC GGG GTT CGA AAT GAC CCA CCT GCC ATC CGG GCA TTT CGA GCG GCA ATT GGC CTT CTT CGG AAT CGT TTT CCG GGA CGC CGC CTT CCA CGC CCT TC CTT GGG AAT CGT TTT CCG GGA CGC CGG CTA GAT CCC CAA CCT GCC ATC ACG AGA TTT CAA CGC GCG CTA TCC AGG CCA CT GCC ATC ACG AGA CTT CGG GGT CGA AAT GAC CCAA CCT GCC ATC CGG GAC CGC CGG CTA TCC ACC CCC CGG CAT CAA GGA AAC CAG CAG CGC CGG CTA TCC ACC CGG CAT CAA GGA AAC CAG CAG CCC CGG GAG AGA CAC TCC CAT GCT GGA AGT CTT TCA CAG GCG CGC TTT CCA GCG CAT CCA GGA GTG GGG AGG CAC CAA CAG CGC CGC TTT CCA GCG CAT CCA GGA GTG GGG AGG CAC CAA CGG CGC CTA TCC GGG CAA ACA GCT CCT GGC CCT GAA GAA GTG GCA CGC GCG CTA TCC GCG CAA CAA TTC AAT TGC GTT TCC TTT TA CCA GTG GGG CAA ACA GCT GCA AAT CCT GTT TGA TGG TGG TTA ACG GCG GGA TAT AAC TAT ATG GCT GCC CTT CCC CTT CCG CTA TCG GCC ACA CCA CAA CGC TAA AGA CGA TGC CCT CAT TCA GCA GAC ACC ACG CGG CTA ACA AGC CGC CTT CCC CTT CCG CTA CCG CCA ACG CGA CCA GAT TT ATT TAT GCC AGC CAG CCA CGC CCA ACC AATG CGA CAA	GIG GAT TIG CAG GGG GGG CGC CT CTG GTA AGG TTG GGA AGG ACT GGA TGG CTT TCT TGC CGC CAA GGA TCT GAT GGC GCA GGG ATC AAG AGA CAG GAT GAC GGT CGT TTC GCA TGC TTG AAC AAG CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC TAT TCG GCT ATG AGA CAA TCG GCT GCT CTG ATG CCG CG TGT TCC GGC TGT CAG CGG TTC TTT TTG TCA AGA CCG ACC TGT CCG GTG CCC TGA ATG AGG CAG CGC GGC TAT CGT GAT GGC GGC CAC CGA CGG GGC TTC CTT TCG ACG TTG TCA CTG AAG CGG GAA GGG ACT GGC TGC CTT GGC AGG ATC TCC CTG ATG CTC ACC TTG CTC GGC GAG AAG CTG ATG CAA TGC GGC GGC TGC ATA CGC TTG ATC CGG CTA CCT ACC AAG CGA AAC ATC GCA CGA AGG GAC CAC GTA CTC GGA TGG TCG ATG CAA AGA CCG GCA TGC ATA CGC TTG ATC CGG CTA CCT ACC AAG CGA AAC ATC GCA CGA AGA GAG CAC GTA CTC GGA TGG CGG ATG CCT GCT TGC CGA ATA CGC AGG GGC AGG ATC GG GCG AGG ATC TCG CGC TGG CAC ATA CGC GTA CTC GGA CGG CGG ATG CCT GCT TGC CGA ATA TCA TGG TGG AAA ATG GCC GCT CGG ACT AGG ATG TGG CGG GCG TGC GAT CGG CGC AGG AGG ATC AGG ACC GTG ATA TTG CTG AAG AGC TTG GCG ACC GCA TCC GCC TTC TAC AGG ATA TTG CTG AAG AGC TTG GGG AAT GGC CTT CTT ACG GTA TTG CTG AAG AGC TTG GGG GCG AAT GGC CTT CTT ACG GTA TTG CTG AAG AGC TTC GGG GCG AAT GGC CTT CTT ACG ATT ATG CTG AAG AGC TTC GGG GCG AAT GGC CTT CTT ACG AGT TCT TCT GAG CGG GAC CTC TGG AAC ATC GCC TTC TA CTT CGG AAT CTT TTT CCG GAA CTC CGG GAT CGA TCC CCT CCA CAA CCT GCC ATC ACG AGA TTT CGA AGC CGG ATT CGA AAC CCC CTG CTG CCT GAG GTT TTT CCG GGA CCT CGG GAT GAT CCT CCA CAA CCT GCC ATC ACG AGA CAC CCC CGG GTT CGA ACC CCT CCA CAA CCT GCA GAG AAC CAG GAG CCT CGC GGA GTT CCA CCC CGG GGA GTG GGG AGG CAC CAT CCG GGA CTA TCC GGG CAT CCA CCC CGG CTG CTG GAG GCT GGA CGA CCC CGC GTT TTG CGG GAA AAC CTT GGC GGA GTG GGG AGG CAC CAT CCG GGA CTA TCC GGG CAA CCA GCC GGC TTT GGG GGG AGG CAC CAA CGG CGG CTA TCC GGG CAA CCA GCC GGC TTT GGG GGG AGA GCT GGA CAA CCG GGG GGA AAA CCT CCA CGC GGG CCC AACG CCA ACG CCA GCA GCG GCA ACA GCC GGA TAT AAC ATG GGG CCC GAA TTG CCG CAA CCG CCA CAC CCA CAA CGC GGA TAT AAC ATG GGG AGG CCG CAA TTG CCG CAA CCA GCA GCA CAA CAA GCC GAA CTA ACA GCC GCA TTG CCA CTA C	GIG GAT TIGC CAG CTG GGG CGC CCT CTG GTA AGG TTG GGA AGC CCT ACT GGA TGG CTT TCT TGC CGC CGC TATG GTA AGG TTG GGA AGC CCT ACT GGA TGG CTT TCT TGC CGC CGC TTTC GCA TGC TTG AAC AAG ATG CAG GTT CTC CGG CCG CTT GGG TGG AGA GGC TAT TCG GCT ATG ACT AGA CAA TCG GCT GCT CTG ATG CCG CCG TGT TCC GGC TGT CAG CGC CGG TTC TTT TTG TCA AGA CCG ACC TGT CCG GTG CCC TGA ATG ACC AGG CAG CGC GGC TAT CGT GGC TGG CCA CGA CGG GCG TTC CTT GCG GGC AGG TTG TCA CTG AAG CCG ACC TGT CCG GCG GCC TAT TGG GCG GGC AGG ATC TCC TGT CAT CTC ACC TTG CTC CTG CCG AGA AAG TAT CTG ATG CAA TGC GGC GGC TGC ATA CTC ACC TTG CTC CTG CCG AGA AAG TAT CTG ATG CAA TGC GGC GGC TGC ATA CTC ACC TTG CTC CTG CGG AGA AGG TAT CTG ATG CAA TGC GGC GGC TGC ATA CTC AGC CTG ATC CGG GGA TGG AAG ACG GAA ACA ATC GCA TCG AGC GAG CAC GTA CTC GGA TGG AAG CCG ATC AGG ATC TCC TGC CGA ATA CGC TTG ATC CGG CAA GGA ACG ATC AGG ATA CTC GGA CGA AGG AGC GAC GTA CTC GGA TGG AAG CCG CCA GGC TCA AGG CGC GCA TGC CCG ACG GCA AGG ATC TCG GCG CGG ATG CCT GCT GC GGA ATA TCA TGG TGG AAA ATG GCC GCT TT CG ACT GTG GCC GCC TCG GTG TGT GGG GCG AAT GGG CTT ACC GGC CTT ACG GTA TCG CCG CTC CCG ATT CGG GGG ACC GCT ATC AGG ACA TAG CCC GTG ATA TTG CCG AGC ACC TCT GGG GCG AAT GGC CTT CTA TCC ACG AGT TCT TCT GAC GGA CAC TCT GGG GTT CGA AAT GAC CGA CCA CCA CCT GCC ATC ACG AGA ATT CGA TTC CAC CGC CCT CTT ATC ACG AGT TCT TCT CGC GGA CAC CCC CGG GCT GAA ATG GAC CCA CCA CAC CCT GCC ATC ACG AAA CCG CCC CCG GGG CTA CCA TGC CCC CCG CAT GCT GGC ATT CTC CGC CAC CCC CGG GCT CGA AAT GAC CCA CCA CCG GAA TC CA GGA AAC CAG CAG CAC CCC CGG GCA TTC CA TGC CCC CCC GGA AGT CCA GGA AAC CAG CAG CGG CTA TCC CGA CAA TCC CCC CGC GAG CTT CGG CAT CCA GGA AAC CAG CAG CGC CGG CTA CCA ACC CCC CGC GAA AAT CCT GTT TGC TTT CGC CAC CCC CGC GGG CAA ACA ATG CCC CCC GGA AAT CCT GTT TGC TTT CGC CAC CCC CGG GGA AAA CCA GCC CGC GAA ATG CCC GAA AAT CCT GTT TGC TGC CCA CCC CGC GGG GAA AGG CGT TTT GCC TAT CGG TAT ATA ATG CGC CCA CAC CGC GGG CAA ACA GCC GGA CAA CCA CTA CGT CGT ATC CCA CTA CCG AGA CGG GGA CAA ACA CAC CTA GGC GGA TTT CCC CAT TCA CGG C	GIG GAC TIG CAG CGA GAC GAC ACC TIG ACT GGA AGC TIG GGA AGC CCT GCA ACC GAA ACC CAA ACT GGA TGG CTT TCT TGC CGC CAA GGA TCT GAT GGC GCA GGG AAC CAA ATC AAA ACAA AGG ACA GAT GAC GGT CGT TTC GCA TGC TG AAC AAA ATG GAT CAA ATG GCT GCT CTG ATG GCA GGA GGC TAT TCC GGC TGT ACA AAG ATG GAC GAC TTT GG GCA GGC TGT TCC CGG CTG TTT CC GGC TGT CAC GGA GGC TAT TTT TTT TTT TTT TTT TTA CAA GA CGA ACC TGT CCG GTG TCC TGA ATG ACC AGA AGG CAA TTT TTT TTT TTT TTA CAA GA CGA ACC TGT CCG GCG TGT TCC CTG ATG ACC AGA GGC TAT TTT TTT TTT TTT TTA CAA GA CGA ACC TGT CCG GCG TGT TAT CGG CGC AGG ACC GTG TCC TTA TGG GCG AAG GGC ACC GTG TCC TTA TGG GCG AAG GGC AGG ATC TCC TGT CAC TGA ACG AAG CGG AAG GGC AGC AGG ATC TCC TGT CAC TGA ATG CGC AGC AGA AGA CAC GTA CTC GGC AGA AAG TAT CCA ACC AGG CAG CAC GTA CTC GGA AAA AG CCC TCG ATC AGG AAA CAC GCA CGA CGC GCA TGC CGA CGC GCA TGC CGA AGA AGG CAC GTA CTC GGA AGA AGG CCG GCG CGA GGC GCA TGC CGA ACG GGA AGG ATC TCG GCG AGA AGA CCG TCG ACG AGG ATG CCG GCT GCT AGG CGG GCG ATG TGG GGA ACA TGA GGA AGA AGG CTT CTG TGG CGA ATA TCA TGG TGG AAA ATG GCC GCT TTT CTG TCG GCG ATG TCG CGC ACC GCT ATC AGG ACA TTA CGC TCC TTT ACC GTG ATA TTG CTG AAG AGC TTG GCG ACC GCT ATC AGG ACA TAG CGT TCC TTT ACG GTA TTT CGA GCA GCT CCC AGC CAT CCC AGC CAC CCC CTT TCT ATC AGC ACC AGC CAC TTC CTG TTT CGG AAA ATG CCG CCT TCT ATC ACC AGC CAC TTC CTG TTT CGG AAA TGC CCT TCT ATC ACC AGC CAC TTC CAC AGC CAC ACC CCC GG ATT CCA CCA CGA CCA AGC CAC TTC CAC AGC CAC ACC CCC CGG CTT CTA ACC AGC CCC CGG AAT CCC CCC CGA ATT CCA GCA ACA AAG CTT CGA GAC ATC CAC GAA AAT CCT CCA CGA CAC AAC CAC ACC CCC CGG CTT CAA CCA GCC CCC CGA AAT CCA GCA CAC ACC CCC CGA AAT CCA GCA CAC ACC CCC CGA AAT CCA GCA CCA ACC CCC CGA AAT CCA GCA CCA ACC CCC CGA AAT CCA GCA CCA ACC AGC CCA ACC A

CCG CTG TTA TGG CCG CGT TTG TCT CAT TCC ACG CCT GAC ACT CAG TTC CGG GTA GGC AGT TCG CTC CAA GCT GGA CTG TAT GCA CGA ACC CCC CGT TCA GTC CGA CCG CTG CGC CTT ATC CGG TAA CTA TCG TCT TGA GTC CAA CCC GGA AAG ACA TGC AAA AGC ACC ACT GGC AGC AGC AGC CAC TGG TAA TTG ATT TAG AGG AGT TAG TCT TGA AGT CAT GCG CCG GTT AAG GCT AAA CTG AAA GGA CAA GTT TTG GTG ACT GCG CTC CTC CAA GCC AGT TAC CTC GGT TCA AAG ACA TGC AAA AGC CCT GCA AGG CCG GTT AAG TCT TTT CGT TTT CAG AGC AAG ACA TTC CTC GAA AGC CCT GCA AGG CGG TTT TTT CGT TTT CAG AGC AAG ACA TTA CGC GCA GAC CAA AAC GAT CTC AAG AGG AGG TTT TTT CGT TTT CAG AGC AAG ACA TTA CGC GCA GAC CAA AAC GAT TTA TTA TTC AAG ATT TAT ATC AGA TAA AAT ATT TCT AGA TTT CAG TGC AAT TTA TTC TCA AGT TTG TAA TTC TCA TAT ATC AGA TAA AAT ATT TCT AGA AGT TGT TAA TTC TCA TGT TTG ACA GCT TAT CAT CGA T

Appendix B

Marker Protein Information

210

1. GFP_{UV} (Clontech) (without a N-terminal hexa-histidine tag)

DNA sequence

ATGAGTAAAGGAGAAGAACTTTTCACTGGAGTTGTCCCAATTCTTGTTGAATTAGATGGTGATGTTAATGGGCACAAATTTTCTGTCAGTGGAGAGGGTGAAGGTGAAGGTGATGCAACATACGGAAAACTTACCCTTAAATTTATTTGCACTACTGGAAAACTTCCTGTTCCATGGGAAACACTTGTCACATTTAAACTCTCTTTCACTACTGGTGAAAAACTTCCTGTTCCATGGGAAACACTTGTCACATTTTTCACTTTTTTCACTACTACTACACTTTTCCCTGTTCCATGGGAAACACTTGTCACATTTTTCAAAGGTACTTTCTTCAAAGATGGTGAAACCATTTCCGTTTATCCGGAAACGACAACTACAACTTTCTTCTATGATGATGACAAAACCAAAACCAAAACGAAAACGACAACCAAAACGACCAAAACGAAAACGAAAACGAAAACGAAAACAACAAAAACAACAAAAACAAAAACAAAA

Amino acid sequence

MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTTGKLPVPWPTLVTTFSYGVQCF SRYPDHMKRHDFFKSAMPEGYVQERTISFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLE YNYNSHNVYITADKQKNGIKANFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSALSKDPNE KRDHMVLLEFVTAAGITHGMDELYKLQPSLIS

2. GFP_{UV}

DNA sequence

ATG AGA GGA TCG CAT CAC CAT CAC CAT CAC GGA TCC GTC GAC CTG CAG CCC CGG GTA CCG GTA GAA AAA ATG AGT AAA GGA GAA GAA CTT TTC ACT GGA GTT GTC CCA ATT CTT GTT GAA TTA GAT GGT GAT GTT AAT GGG CAC AAA TTT TCT GTC AGT GGA GAG GGT GAA GGT GAT GCA ACA TAC GGA AAA CTT ACC CTT AAA TTT ATT TGC ACT ACT GGA AAA CTA CCT GTT CCA TGG CCA ACA CTT GTC ACT ACT TTC TCT TAT GGT GTT CAA TGC TTT TCC CGT TAT CCG GAT CAT ATG AAA CGG CAT GAC TTT TTC AAG AGT GCC ATG CCC GAA GGT TAT GTA CAG GAA CGC ACT ATA TCT TTC AAA GAT GAC GGG AAC TAC AAG ACG CGT GCT GAA GTC AAG TTT GAA GGT GAT ACC CTT GTT AAT CGT ATC GAG TTA AAA GGT ATT GAT TTT AAA GAA GAT GGA AAC ATT CTC GGA CAC AAA CTC GAG TAC AAC TAT AAC TCA CAC AAT GTA TAC ATC ACG GCA GAC AAA CAA AAG AAT GGA ATC AAA GCT AAC TTC AAA ATT ATG AGA GGA TCG CAT CAC CAT CAC CAT CAC GGA TCC GTC GAC CTG CAG CCC CGG GTA CCG GTA GAA AAA ATG AGT AAA GGA GAA GAA CTT TTC ACT GGA GTT GTC CCA ATT CTT GTT GAA TTA GAT GGT GAT GTT AAT GGG CAC AAA TTT TCT GTC AGT GGA GAG GGT GAA GGT GAT GCA ACA TAC GGA AAA CTT ACC CTT AAA TTT ATT TGC ACT ACT GGA AAA CTA CCT GTT CCA TGG CCA ACA CTT GTC ACT ACT TTC TCT TAT GGT GTT CAA TGC TTT TCC CGT TAT CCG GAT CAT ATG AAA CGG CAT GAC TTT TTC AAG AGT GCC ATG CCC GAA GGT TAT GTA CAG GAA CGC ACT ATA TCT TTC AAA GAT GAC GGG AAC TAC AAG ACG CGT GCT GAA GTC AAG TTT GAA GGT GAT ACC CTT GTT AAT CGT ATC GAG TTA AAA GGT ATT GAT TTT AAA GAA GAT GGA AAC ATT CTC GGA CAC AAA CTC GAG TAC AAC TAT AAC TCA CAC AAT GTA TAC ATC ACG GCA GAC AAA CAA AAG AAT GGA ATC AAA GCT AAC TTC AAA ATT CGC CAC AAC ATT GAA GAT GGA TCC GTT CAA CTA GCA GAC CAT TAT CAA CAA AAT ACT CCA ATT GGC GAT GGC CCT GTC CTT TTA CCA GAC AAC CAT TAC CTG TCG ACA CAA TCT GCC CTT TCG AAA GAT CCC AAC GAA AAG CGT GAC CAC ATG GTC CTT CTT GAG TTT GTA ACT GCT GCT GGG ATT ACA CAT GGC ATG GAT GAG CTC TAC AAA CTG CAG CCA AGC TTA ATT AGC TGA

Amino acid sequence

MRGSHHHHHHGSVDLQPRVPVEKMSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFIC TTGKLPVPWPTLVTTFSYGVQCFSRYPDHMKRHDFFKSAMPEGYVQERTISFKDDGNYKTRAEVKFEGDTL VNRIELKGIDFKEDGNILGHKLEYNYNSHNVYITADKQKNGIKANFKIRHNIEDGSVQLADHYQQNTPIGD GPVLLPDNHYLSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDELYKLQPSLIS

3. GFP3

F64L, S65T, S99F, and T153M mutations in GFP_{UV}

DNA sequence

ATGAGAGGATCGCATCACCATCACCATCACGGATCCGTCGACCTGCAGCCGCGGGTACCGGTAAAAAAAATGAGTAAAGGAGAACATTTCACTGAAGTTGTCCCAATTCTTGTTGAATTAGATGGTGATGATGATAAAGGACTTTTCAAATTTTTTACTGCAAGTAGTGAGGGTGAAGGTGATGCTGATGGTGAACTTACTACATTTAAACTTACATTTACTACAACACTTACTACAACTACAACACTTACTACATTTAAACTTACAACTACACTTACAACACTTACTACAACTACACTTACAACACTTACTACAACTACACTTACAACACTTACAACTACACTTAAACTTAAATTTAATTCTACTACTACAACTACAACT<td

Amino acid sequence

MRGSHHHHHHGSVDLQPRVPVEKMSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFIC TTGKLPVPWPTLVTTLTYGVQCFSRYPDHMKRHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTL VNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKANFKIRHNIEDGSVQLADHYQQNTPIGD GPVLLPDNHYLSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDELYKLQPSLIS

4. GFP6

L64F mutation and all Phe codons are changed to UUU in GFP3

DNA sequence

ATG AGA GGA TCG CAT CAC CAT CAC CAT CAC GGA TCC GTC GAC CTG CAG CCC CGG GTA CCG GTA GAA AAA ATG AGT AAA GGA GAA GAA CTT TTT ACT GGA GTT GTC CCA ATT CTT GTT GAA TTA GAT GGT GAT GTT AAT GGG CAC AAA TTT TCT GTC AGT GGA GAG GGT GAA GGT GAT GCA ACA TAC GGA AAA CTT ACC CTT AAA TTT ATT TGC ACT

ACTGGAAAACTACCTGTTCCATGGCCAACACTTGTCACTTTTACCTATGGTGTTCAATGCTTTTCCCGTTATCCGGATCATATGAAACGGCATGACTTTTTTAAGAGTGCCATGCCCGAAGGTTATGTACAGGAACGCACTATATTTTTTAAAGATGACGGGAACTACAAGACGCGTGCTGAAGTCAAGGGTATATTTAAAGATGACGGGAACTACAAGACGCGTGATGATGAAGGTGAAGATGAAATTCACCATAATCGTATCGAGTAAAAGGGTATTGATGAAGATGAAGATAACCACAAACTGGAGTAAAACTATGATGATGAAGATGAAAATGAAAACATTCTCGGAAACCACAAACTGGAAGATAACTATAAAATTCGCAACATTGAAAAACAACAAAAACTGGAAGATAACTATAAAATTCGCAACATTGAAAAACAAAAAGAAGAAAATACTCCAAACAATACTGCAAATGATGACAAACAA</

Amino acid sequence

MRGSHHHHHHGSVDLQPRVPVEKMSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFIC TTGKLPVPWPTLVTTFTYGVQCFSRYPDHMKRHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTL VNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKANFKIRHNIEDGSVQLADHYQQNTPIGD GPVLLPDNHYLSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDELYKLQPSLIS

5. GFP_WC

All Phe codons are changed to UUC in GFP3

DNA sequence

ATG AGA GGA TCG CAT CAC CAT CAC CAT CAC GGA TCC GTC GAC CTG CAG CCC CGG GTA CCG GTA GAA AAA ATG AGT AAA GGA GAA GAA CTT TTC ACT GGA GTT GTC CCA ATT CTT GTT GAA TTA GAT GGT GAT GTT AAT GGG CAC AAA TTC TCT GTC AGT GGA GAG GGT GAA GGT GAT GCA ACA TAC GGA AAA CTT ACC CTT AAA TTC ATT TGC ACT ACT GGA AAA CTA CCT GTT CCA TGG CCA ACA CTT GTC ACT ACT CTG ACC TAT GGT GTT CAA TGC TTC TCC CGT TAT CCG GAT CAT ATG AAA CGG CAT GAC TTC TTC AAG AGT GCC ATG CCC GAA GGT TAT GTA CAG GAA CGC ACT ATA TTC TTC AAA GAT GAC GGG AAC TAC AAG ACG CGT GCT GAA GTC AAG TTC GAA GGT GAT ACC CTT GTT AAT CGT ATC GAG TTA AAA GGT ATT GAT TTC AAA GAA GAT GGA AAC ATT CTC GGA CAC AAA CTC GAG TAC AAC TAT AAC TCA CAC AAT GTA TAC ATC ATG GCA GAC AAA CAA AAG AAT GGA ATC AAA GCT AAC TTC AAA ATT CGC CAC AAC ATT GAA GAT GGA TCC GTT CAA CTA GCA GAC CAT TAT CAA CAA AAT ACT CCA ATT GGC GAT GGC CCT GTC CTT TTA CCA GAC AAC CAT TAC CTG TCG ACA CAA TCT GCC CTT TCG AAA GAT CCC AAC GAA AAG CGT GAC CAC ATG GTC CTT CTT GAG TTC GTA ACT GCT GCT GGG ATT ACA CAT GGC ATG GAT GAG CTC TAC AAA CTG CAG CCA AGC TTA ATT AGC TGA

Amino acid sequence

MRGSHHHHHHGSVDLQPRVPVEKMSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFIC TTGKLPVPWPTLVTTLTYGVQCFSRYPDHMKRHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTL VNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKANFKIRHNIEDGSVQLADHYQQNTPIGD GPVLLPDNHYLSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDELYKLQPSLIS

6. GFP6 (2UUU)

AGA (Arg) to UUU (Phe) mutation in position 2 of GFP6

213

7. GFP6 (2UUC)

AGA (Arg) to UUU (Phe) mutation in position 2 of GFP6

8. GFP6 (158Am)

AAA (Lys) to UAG (an amber codon) mutation in position 158 of GFP6

9. GFP3C

GFP3 with a C-terminal hexa-histidine tag

DNA sequence

ATG CGA GGA TCC CTG CAG CCC CGG GTA CCG GTA GAA AAA ATG AGT AAA GGA GAA GAA CTT TTC ACT GGA GTT GTC CCA ATT CTT GTT GAA TTA GAT GGT GAT GTT AAT GGG CAC AAA TTT TCT GTC AGT GGA GAG GGT GAA GGT GAT GCA ACA TAC GGA AAA CTT ACC CTT AAA TTT ATT TGC ACT ACT GGA AAA CTA CCT GTT CCA TGG CCA ACA CTT GTC ACT ACT CTG ACC TAT GGT GTT CAA TGC TTC TCC CGT TAT CCG GAT CAT ATG AAA CGG CAT GAC TTT TTC AAG AGT GCC ATG CCC GAA GGT TAT GTA CAG GAA CGC ACT ATA TTC TTC AAA GAT GAC GGG AAC TAC AAG ACG CGT GCT GAA GTC AAG TTT GAA GGT GAT ACC CTT GTT AAT CGT ATC GAG TTA AAA GGT ATT GAT TTT AAA GAA GAT GGA AAC ATT CTC GGA CAC AAA CTC GAG TAC AAC TAT AAC TCA CAC AAT GTA TAC ATC ATG GCA GAC AAA CAA AAG AAT GGA ATC AAA GCT AAC TTC AAA ATT CGC CAC AAC ATT GAA GAT GGA TCC GTT CAA CTA GCA GAC CAT TAT CAA CAA AAT ACT CCA ATT GGC GAT GGC CCT GTC CTT TTA CCA GAC AAC CAT TAC CTG TCG ACA CAA TCT GCC CTT TCG AAA GAT CCC AAC GAA AAG CGT GAC CAC ATG GTC CTT CTT GAG TTT GTA ACT GCT GCT GGG ATT ACA CAT GGC ATG GAT GAG CTC TAC AAA CTG CAG CAT CAC CAT CAC CAT CAC TAA

Amino acid sequence

MRGSLQPRVPVEKMSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTTGKLPVPWP TLVTTLTYGVQCFSRYPDHMKRHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGID FKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKANFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHY LSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDELYKLQHHHHHH

10. GFP3C (158UUG)

AAA (Lys) to UUG (Leu) mutation in position 158 of GFP3C

11. mDHFR (Qiagen)

C-terminal hexa-histidine tag

DNA sequence

ATG AGA GGA TCC GGC ATC ATG GTT CGA CCA TTG AAC TCG ATC GTC GCC GTG TCC CAA AAT ATG GGG ATT GGC AAG AAC GGA GAC CTA CCC TGG CCT CCG CTC AGG AAC GAG TTC AAG TAC TTC CAA AGA ATG ACC ACA ACC TCT TCA GTG GAA GGT AAA CAG AAT CTG GTG GTG ATT ATG GGT AGG AAA ACC TGG TTC TCC ATT CCT GAG AAG AAT CGA CCT TTA AAG GAC AGA ATT AAT ATA GTT CTC AGT GAT GAC ACA CCT ATA AGA CCA CCA CCA CGA GGA GCT CAT TTT CTT GCC AAA AGT AGT AGA ATG GAT GAT GAT GAC CTC AAG GAA GCC AGT TCT GGA AGG AAG AAT CGA CCG GAA TTG GCA AGT AAA GTA GAC ATG GTT TGG ATA GTC GGA GGC AGT TCT GTT TAC CAG GAA GCC ATG AAT CAA CCA GGC CAC CTT AGA CTC TTT GTG ACA AGG AAT AAT ATA CAA CCA GGC CAC CTT AGA CTC TTT GTG ACA AGG AAT ATA AAA CTT CAA CAA CCA GGC CAC CTT AGA CTC TTT GTG ACA AGG AAT AAT AAA CTA AAA CTA CAA CCA GGC CAC CTT AGA ATT GAT TTG GGG AAA AAT GAC ATG GTT TAC CAA GAA CTC CCA GAA ATT GAA AGT AAA CAA CCA GGC ATC ATT AAA CTT CTC CCA GAA ATT GAT TTG GGG AAA GGC ATC ATT CTC CCA GAA CTT CTC CAA GAA CTT CTT GGG AAA AGT GAC ATG GTT TTT CTC CCA GAA ATT GAT TTT GAG AAA GGC ATC TTT CTC CCA GAA ATT GAT TTT GAA AAA GTT CAA CCA GGC ATC TTT TTT CTC CAA GAA CTC CAG GAA AAA GGC ATC ATC

Amino acid sequence

MRGSGIMVRPLNSIVAVSQNMGIGKNGDLPWPPLRNEFKYFQRMTTTSSVEGKQNLVIMGRKTWFSIPEKN RPLKDRINIVLSRELKEPPRGAHFLAKSLDDALRLIEQPELASKVDMVWIVGGSSVYQEAMNQPGHLRLFV TRIMQEFESDTFFPEIDLGKYKLLPEYPGVLSEVQEEKGI

12. mDHFR2

UUC (Phe) in position 38 and UUU (Phe) in position 95 to UUG (Leu) mutations in mDHFR

DNA sequence

ATGAGAGGATCCGGCATCATGGTTCGACCATTGAACTCGATCGTCGCCGTGTCCCAAAATATGGGGATTGGCAAGAACGGAGACCTACCCTGGCCTCCGCTCAGGAACGAGTTGAAGTTCCAAAGAATGACCACCTCTTCAGTGGAAGGTAAACAGAATCTGGTGATTATGGGTAGGAAAACCTGGTCTTCAGTGGAAGGTAAACAGCATTTGGTGATTATGAGGAAAACCTGGTCTTCAGAGAAACAGCAGGGAGCTCATTTGGCTAGAAATAATATTCTCAGAAAACAGCAGGGAGCTCATTTGGCTAGAAGTAAAACCTCGAAAAGACCACCACAGGGAGCTCATTTGGCTAAAAGTAAAAGTTTGAGAGAACTCAAAAGACCACCACAGGGAGCTCATTTGGCTAAAAGTAAAAGTTTGGATGATGCTAAAAGAAGTAGAAGAAGTAGAAGTAGAAGTAGAAGTAGAAGTAGAAGTAGAAGTAGAAG

Amino acid sequence

MRGSGIMVRPLNSIVAVSQNMGIGKNGDLPWPPLRNELKYFQRMTTTSSVEGKQNLVIMGRKTWFSIPEKN RPLKDRINIVLSRELKEPPRGAHLLAKSLDDALRLIEQPELASKVDMVWIVGGSSVYQEAMNQPGHLRLFV TRIMQEFESDTFFPEIDLGKYKLLPEYPGVLSEVQEEKGIKYKFEVYEKKGSRSHHHHHH

13. mDHFR2 (100UUA)

UUG to UUA mutation in position 100 of mDHFR2

Appendix C

Names and Sequences of PCR Primers

Primer name	Primer sequence
GFPI	5'- AC CTGCAG CCC CGG GTA CCG GTA GAA - 3'
GFP2	5'- AC CTGCAG TTT GTA GAG CTC ATC CAT - 3'
F64LS65T f	5'- CTT GTC ACT ACT CTG ACC TAT GGT GTT CAA
10125001_1	TGC TTC TCC CGT - 3'
F64I \$65T r	5'- ACG GGA GAA GCA TTG AAC ACC ATA GGT CAG
	AGT AGT GAC AAG - 3'
SOOF f	5'- GTA CAG GAA CGC ACT ATA TTC TTC AAA GAT
3991°_1	GAC GGG AAC - 3'
S00E *	5'- GTT CCC GTC ATC TTT GAA GAA TAT AGT GCG
399F_f	TTC CTG TAC - 3'
T152N f	5'- CAC AAT GTA TAC ATC ATG GCA GAC AAA CAA
1153M_f	AAG AAT GGA - 3'
T15214	5'- TCC ATT CTT TTG TTT GTC TGC CAT GAT GTA
1153M_r	TAC ATT GTG - 3'
	5'- GTG CCA CCT GAC GTC TAA GAA ACC ATT ATT
Primer 1	ATC ATG ACA TTA ACC - 3'
	5' - GAG TAA AGG AGA AGA ACT TTT TAC TGG AGT
Primer 2	TGT CCC AAT TC - 3'
	5'- GAA TTG GGA CAA CTC CAG TAA AAA GTT CTT
Primer 3	CTC CTT TAC TC - 3'
	5'- GGC CAA CAC TTG TCA CTA CTT TTA CCT ATG
Primer 4	GTG TTC AAT GCT T - 3'
	$5'_{-}$ A G C AT TGA ACA CCA TAG GTA AAA GTA GTG
Primer 5	$\Delta C \Delta \Delta G T G T T G G C C = 3'$
	5'- CAT ATG AAA CGG CAT GAC TTT TTT AAG AGT
Primer 6	$GCC \Delta TG CCC G \Delta \Delta G = 3'$
	5' CTT CGG GCA TGG CAC TCT TAA AAA AGT CAT
Primer 7	GCC GTT TCA TAT G 3'
	$\frac{1}{5} CTT ATC TAC ACC CCA CTA TAT TTT TCA$
Primer 8	3 - 011 AIG TAC AGG AAC GCA CTA TAT TIT TCA
	AAU ATU ACU UUA ACT ACA A $- 3$
Primer 9	5 - TIG TAG TIC CCG ICA ICI TIG AAA AAT ATA
	GIG CGI ICC IGI ACA TAA C - 3
Primer 10	5'- ACA AAA GAA IGG AAI CAA AGC IAA CII IAA
	AAT TCG CCA CAA CAT TGA AGA TG - 3'
Primer 11	5'- CAT CTT CAA TGT TGT GGC GAA TTT TAA AGT
	TAG CIT IGA TIC CAT ICI TII GI - 3'
Primer 12	5'- CGC CAA GCT AGC TTG GAT TCT CAC CAA TAA
	AAA ACG CCC - 3'
Primer 13	5'- TAC CTA TGG TGT TCA ATG CTT TTC CCG TTA
	TCC GGA TCA TAT G - 3'
Primer 14	5'- CAT ATG ATC CGG ATA ACG GGA AAA GCA TTG
	AAC ACC ATA GGT A - 3'

Primer 15	5'- GTT ATG TAC AGG AAC GCA CTA TAT TTT TTA AAG ATG ACG GGA ACT ACA AG - 3'
Primer 16	5'- CTT GTA GTT CCC GTC ATC TTT AAA AAA TAT AGT GCG TTC CTG TAC ATA AC - 3'
Primer 17	5'- GT GAT GTT AAT GGG CAC AAA TTC TCT GTC
Primer 18	5'- ACC CTC TCC ACT GAC AGA GAA TTT GTG CCC
Primer 19	5'- CA TAC GGA AAA CTT ACC CTT AAA TTC ATT TGC ACT ACT GGA AAA CTA CC- 3'
Primer 20	5'- GGT AGT TTT CCA GTA GTG CAA ATG AAT TTA AGG GTA AGT TTT CCG TAT G- 3'
Primer 21	5'- AT CAT ATG AAA CGG CAT GAC TTC TTC AAG AGT GCC ATG CCC- 3'
Primer 22	5'- GGG CAT GGC ACT CTT GAA GAA GTC ATG CCG TTT CAT ATG AT- 3'
Primer 23	5'- CG CGT GCT GAA GTC AAG TTC GAA GGT GAT ACC CTT GT- 3'
Primer 24	5'- AC AAG GGT ATC ACC TTC GAA CTT GAC TTC AGC ACG CG- 3'
Primer 25	5'- CGT ATC GAG TTA AAA GGT ATT GAT TTC AAA GAA GAT GGA AAC ATT CTC GGA- 3'
Primer 26	5'- TCC GAG AAT GTT TCC ATC TTC TTT GAA ATC AAT ACC TTT TAA CTC GAT ACG - 3'
lacI_AflII_f	5'- CAT ATT CTT AAG TCA TGC GCA CCC GTG GCC
<i>lac</i> I_SacII_r	5'- CAA CTC CCG CGG TTA CGT GCA GAT GAT AAG
pQE_AflII_f	5'-CAA GTA CTT AAG CGG TGT GAA ATA CCG CAC AGA TGC GTA AGG -3'
pQE_SacII_r	5'- GAT CAT CCG CGG TTT TCC CAG TCA CGA CGT TGT AAA ACG - 3'
437_f	5' - GTC GAA ATC GGT AAC NNK GGT ATG TTC AGA CCA GAA ATG CTC G -3'
437_r	5' - CGA GCA TTT CTG GTC TGA ACA TAC CMN NGT TAC CGA TTT CGA C - 3'
412_415_418_f	5' - C AAG CCT ACC TAC NNK CCT TAC NNK GAG CCA NNK ATG GAA ATC TTT T - 3'
412_415_418_r	5'- A AAA GAT TTC CAT MNN TGG CTC MNN GTA AGG MN N GTA GG T AGG CTT G - 3'
2F_UUU_f	5' - C ATT AAA GAG GAG AAA TTA ACT ATG TTT GGA TCG CAT CAC - 3'
2F_UUU_r	5' - GTG ATG CGA TCC AAA CAT AGT TAA TTT CTC CTC TTT AAT G - 3'
2F_UUC_f	5' - C ATT AAA GAG GAG AAA TTA ACT ATG TTC GGA TCG CAT CAC - 3'

2F_UUC_r	5' - GTG ATG CGA TCC GAA CAT AGT TAA TTT CTC CTC TTT AAT G - 3'
R_lacP_f	GCA CCC CAG TCT TTA CAC TAT GCC TAC CGT CTC CGT GAA C
R_lacP_r	GTT CAC GGA GAC GGT AGG CAT AGT GTA AAG ACT GGG GTG C
Ins_SD_A_Del_lacO_f	CGT ATA ATG GAT CC C GAT TTT CAC ATA AGG AGC AGA CC ATG
Ins_SD_A_Del_lacO_r	CAT GGT CTG CTC CTT ATG TGA AAA TCG GGA TCC ATT ATA CG
ytRNA-GAA_f	5' - CAC AGG ACC TCC AGA TTT TCA GTC TGG CGC TCT CC - 3'
ytRNA-GAA_r	5' - GGA GAG CGC CAG ACT GAA AAT CTG GAG GTC CTG TG - 3'
ytRNA-CAA_f	5' - CAC AGG ACC TCC AGA TTT TGA GTC TGG CGC TCT CC - 3'
ytRNA-CAA_r	5' - GGA GAG CGC CAG ACT CAA AAT CTG GAG GTC CTG TG - 3'
L100_UUA_f	5' - CTT GCC AAA AGT TTA GAT GAT GCC TTA AGA C - 3'
L100_UUA_r	5' - GTC TTA AGG CAT CAT CTA AAC TTT TGG CAA G - 3'
PstI_70_f	5' - GCA TGC GAG GAT CCC TGC AGC ATC ACC ATC ACC - 3'
PstI_70_r	5'- GGT GAT GGT GAT GCT GCA GGG ATC CTC GCA TGC - 3'
K158_UUG_f	5' - ATG GCA GAC AAA CAA TTG AAC GGA ATC AAA GCT AAC TTC -3'
K158_UUG_r	5' - GAA GTT AGC TTT GAT TCC GTT CAA TTG TTT GTC TGC CAT - 3'
K158_UAG_f	5' - CAT CAT GGC AGA CAA ACA ATA GAA TGG AAT CAA AGC TAA C - 3'
K158_UAG_r	5' - GTT AGC TTT GAT TCC ATT CTA TTG TTT GTC TGC CAT GAT G- 3'