

## Bibliography

- Aegerter-Wilmsen T., Aegerter C.M., Bisseling T. (2005). Model for the robust establishment of precise proportions in the early Drosophila embryo. *J Theor Biol.* 234: 13-19.
- Ahn S., Joyner A. L. (2004). Dynamic changes in the response of cells to positive hedgehog signaling during mouse limb patterning. *Cell.* 118: 505-516.
- Akam M. (1987). The molecular basis for metamerization in the Drosophila embryo. *Development.* 101: 1-22.
- Alon U., Surette M.G., Barkai N., Leibler S. (1999). Robustness in bacterial chemotaxis. *Nature.* 397: 168-171.
- Apionishev S., Katanayeva N.M., Marks S.A., Kalderon D., Tomlinson A. (2005). Drosophila Smoothened phosphorylation sites essential for Hedgehog signal transduction. *Nat Cell Biol.* 7: 86-92.
- Ashe H. L., Briscoe J. (2006). The interpretation of morphogen gradients. *Development.* 133: 385-394.
- Barkai N., Liebler S. (1997). Robustness in simple biochemical networks. *Nature.* 387: 913-917.
- Belvin M.P., Jin Y., Anderson K.V. (1995). Cactus protein degradation mediates Drosophila dorsal-ventral signaling. *Genes Dev.* 9: 783-793.
- Ben-Zvi D., Barkai N. (2010). Scaling of morphogen gradients by an expansion-

repression integral feedback control. Proc Natl Acad Sci U S A. 107: 6924-6929.

Ben-Zvi D., Shilo B.Z., Fainsod A., Barkai N. (2008). Scaling of the BMP activation gradient in *Xenopus* embryos. Nature. 453: 1205-1211.

Bergmann A., Stein D., Geisler R., Hagenmaier S., Schmid B., et al. (1996). A gradient of cytoplasmic Cactus degradation establishes the nuclear localization gradient of the dorsal morphogen in *Drosophila*. Mech Dev. 60: 109-123.

Bergmann S., Sandler O., Sberro H., Shnider S., Schejter E., et al. (2007). Pre-steady-state decoding of the Bicoid morphogen gradient. PLoS Biol. 5: e46. doi:10.1371/journal.pbio.0050046.

Biehs B., Francois V., Bier E. (1996). The *Drosophila* short gastrulation gene prevents Dpp from autoactivating and suppressing neurogenesis in the neuroectoderm. Genes Dev 10: 2922–2934.

Bier E., Jan L.Y., Jan Y.N. (1990). Rhomboid, a gene required for dorsoventral axis establishment and peripheral nervous system development in *Drosophila melanogaster*. Genes Dev. 4: 190-203.

Bijlsma M.F., Spek C.A., Zivkovic D., van de Water S., Rezaee F., Peppelenbosch M. P. (2006). Repression of smoothened by patched-dependent (pro-)vitamin D3 secretion. PLoS Biol. 4:e232.

Blair S.S. (1992). Engrailed expression in the anterior lineage compartment of the developing wing blade of *Drosophila*. Development. 115:21–33.

- Bollenbach T., Pantazis P., Kicheva A., Bökel C., González-Gaitán M., Jülicher F. (2008). Precision of the Dpp gradient. *Development*. 135: 1137-1146.
- Bolouri H. (2008). Embryonic pattern formation without morphogens. *Bioessays*. 30: 412-417.
- Bourillot P.Y., Garret N., Gurdon J.B. (2002). A changing morphogen gradient is interpreted by continuous transduction flow. *Development*. 129: 2167-2180.
- Brand A.H., Perrimon N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development*. 118: 401-415.
- Bryant P.J. (1969). Mosaicism. *Bioscience*. 19: 1126.
- Burke R., Nellen D., Bellotto M., Hafen E., Senti K.A., et al. (1999). Dispatched, a novel sterol-sensing domain protein dedicated to the release of cholesterol-modified hedgehog from signaling cells. *Cell*. 99: 803–815.
- Capdevila J., Estrada M.P., Sanchez-Herrero E., Guerrero I. (1994). The Drosophila segment polarity gene patched interacts with decapentaplegic in wing development. *EMBO J*. 13: 71–82.
- Carroll S.B., Grenier J.K., Weatherbee S.D. (2005). From DNA to diversity: molecular genetics and the evolution of animal design. 2<sup>nd</sup> Edition. Blackwell Publishing. Malden, MA.
- Casali A., Struhl G. (2004). Reading the Hedgehog morphogen gradient by measuring the ratio of bound to unbound Patched protein. *Nature*. 431: 76-80.

Chamoun Z., Mann R.K., Nellen D., von Kessler D.P., Bellotto M., et al. (2001). Skinny hedgehog, an acyltransferase required for palmitoylation and activity of the hedgehog signal. *Science*. 293: 2080–2084.

Chen Y., Struhl G. (1996). Dual roles for patched in sequestering and transducing Hedgehog. *Cell*. 87: 553–563.

Child C. M. (1941). Patterns and Problems of Development. Chicago University Press. Chicago, IL.

Chuang P.T., McMahon A.P. (1999) Vertebrate Hedgehog signalling modulated by induction of a Hedgehog-binding protein. *Nature*. 397: 617–621.

Cohen J.E. (2004). Mathematics is biology's next microscope, only better; biology is mathematics' next physics, only better. *PLoS Biol*. 2: e439.

Cooke J. (1995). Morphogens in vertebrate development: how do they work? *Bioessays*. 17: 93–96.

Cowden J., Levine M. (2003). Ventral dominance governs sequential patterns of gene expression across the dorsal-ventral axis of the neuroectoderm in the *Drosophila* embryo. *Dev Biol*. 262: 335-349.

Dahmann C., Basler K. (2000). Opposing transcriptional outputs of Hedgehog signalling and engrailed control compartmental cell sorting at the *Drosophila* A/P boundary. *Cell*. 100: 411-422.

Dalcq A.M. (1938). Form and causality in early development. Cambridge University Press. Cambridge, UK.

- Dalcq A., Pasteels J.J. (1937). Une conception nouvelle des bases physiologiques de la morphogénèse. *Arch Biol.* 48:669–710.
- Davidson E.H. (2006). *The Regulatory Genome: Gene Regulatory Networks in Development and Evolution*. Academic Press. Burlington, MA.
- de Celis J.F., Ruiz-Gomez M. (1995) groucho and hedgehog regulate engrailed expression in the anterior compartment of the *Drosophila* wing. *Development*. 121: 3467–3476.
- de Lachapelle A.M., Bergmann S. (2010). Precision and scaling in morphogen gradient read-out. *Mol Syst Biol.* 6: 351.
- De Robertis E.M. (2006). Spemann's organizer and self-regulation in amphibian embryos. *Nat Rev Mol Cell Biol.* 7:296-302.
- Denef N., Neubuser D., Perez L., Cohen S.M. (2000). Hedgehog induces opposite changes in turnover and subcellular localization of patched and smoothened. *Cell*. 102: 521–531.
- Dessaud E., Ribes V., Balaskas N., Yang L.L., Pierani A., et al. (2010). Dynamic assignment and maintenance of positional identity in the ventral neural tube by the morphogen sonic hedgehog. *PLoS Biol.* 8: 1000382.
- Dessaud E., Yang L.L., Hill K., Cox B., Ulloa F., et al. (2007). Interpretation of the sonic hedgehog morphogen gradient by a temporal adaptation mechanism. *Nature*. 450: 717–720.
- Drier E.A., Huang L.H., Steward R. (1999). Nuclear import of the *Drosophila* Rel protein

Dorsal is regulated by phosphorylation. *Genes Dev.* 13: 556-568.

Driever W., Nüsslein-Volhard C. (1988a). A gradient of bicoid protein in Drosophila embryos. *Cell.* 54: 83-93

Driever W., Nüsslein-Volhard C. (1988b). The bicoid protein determines position in the Drosophila embryo in a concentration-dependent manner. *Cell.* 54: 95-104.

Drossopoulou G., Lewis K.E., Sanz-Ezquerro J.J., Nikbakht N., McMahon A.P., et al. (2000) A model for anteroposterior patterning of the vertebrate limb based on sequential long- and short-range Shh signalling and Bmp signalling. *Development.* 127: 1337–1348.

Dyson S., Gurdon J.B. (1998). The interpretation of position in a morphogen gradient as revealed by occupancy of activin receptors. *Cell.* 93: 557–568.

Edwards D.N., Towb P., Wasserman S.A. (1997) An activity-dependent network of interactions links the Rel protein Dorsal with its cytoplasmic regulators. *Development.* 124: 3855-3864.

Eldar A., Rosin D., Shilo B.Z., Barkai N. (2003). Self-enhanced ligand degradation underlies robustness of morphogen gradients. *Dev Cell.* 5: 635-646.

Ferguson E.L., Anderson K.V. (1992). Decapentaplegic acts as a morphogen to organize dorsal-ventral pattern in the Drosophila embryo. *Cell.* 71: 451-461.

Fischer J.A., Giniger E., Maniatis T., Ptashne M. (1988). GAL4 activates transcription in Drosophila. *Nature.* 332: 853-856.

Francois V., Solloway M., O'Neill J.W., Emery J., Bier E. (1994). Dorsal-ventral

patterning of the *Drosophila* embryo depends on a putative negative growth factor encoded by the short gastrulation gene. *Genes Dev.* 8: 2602-2616.

García-Bellido A. (1975). Genetic control of wing disc development in *Drosophila*. *Ciba Found Symp.* 0: 161-182.

Gierer A., Meinhardt H. (1972) A theory of biological pattern formation. *Kybernetik.* 12: 30-39.

Golembio M., Schweitzer R., Freeman M., Shilo B.Z. (1996). Argos transcription is induced by the *Drosophila* EGF receptor pathway to form an inhibitory feedback loop. *Development.* 122: 223–230.

Gonzalez A., Chaouiya C., Thieffry D. (2008). Logical modelling of the role of the Hh pathway in the patterning of the *Drosophila* wing disc. *Bioinformatics.* 24: 234–240.

González-Gaitán M., Capdevila M.P., García-Bellido A. (1994). Cell proliferation patterns in the wing disc of *Drosophila*. *Mech Dev.* 46: 183-200.

Green J. (2002). Morphogen gradients, positional information and *Xenopus*: interplay of theory and experiment. *Dev Dyn.* 225: 392-408.

Green J., Smith J.C. (1990). Graded changes in dose of a *Xenopus* activin elicit stepwise transitions in embryonic cell fate. *Nature.* 347: 391–394.

Gregor T., Bialek W., de Ruyter van Steveninck R.R., Tank D.W., Wieschaus E.F. (2005). Diffusion and scaling during early embryonic pattern formation. *Proc Natl Acad Sci U S A.* 102: 18403-18407.

Gregor T., McGregor A.P., Wieschaus E.F. (2008). Shape and function of the Bicoid

morphogen gradient in dipteran species with different sized embryos. *Dev Biol.* 316:350-358.

Gregor T., Tank D.W., Wieschaus E.F., Bialek W. (2007). Probing the limits to positional information. *Cell.* 130: 153-164.

Gregor T., Wieschaus E.F., McGregor A.P., Tank D.W. (2007). Stability and nuclear dynamics of the bicoid morphogen gradient. *Cell* 130: 141-152.

Grier A., Meinhardt H. (1972). A theory of biological pattern formation. *Kybernetik* 12: 30-39.

Gunbin K.V., Omelyanchuk L.V., Kogai V.V., Fadeev S.I., Kolchanov N.A. (2007). Model of the reception of hedgehog morphogen concentration gradient: comparison with an extended range of experimental data. *J Bioinform Comput Biol.* 5: 491-506.

Gurdon J.B., Bourillot P.Y. (2001). Morphogen gradient interpretation. *Nature.* 413: 797-803.

Gurdon J.B., Dyson S., St Johnston D. (1998). Cells' perception of position in a concentration gradient. *Cell.* 95: 159-162.

Gurdon J.B., Harger P., Mitchell A., Lemaire P. (1994). Activin signaling and response to a morphogen gradient. *Nature.* 371: 487-492.

Gurdon J.B., Mitchell A., Mahony D. (1995). Direct and continuous assessment by cells of their position in a morphogen gradient. *Nature.* 376: 520–521.

Gurdon J.B., Standley H., Dyson S., Butler K., Langon T., et al. (1999). Single cells can sense their position in a morphogen gradient. *Development.* 126: 5309-5317.

Gustafson T. (1965). In: The Biochemistry of Development. Weber E. (ed.). Academic Press. New York, NY. p.139.

Han C., Belenkaya T.Y., Wang B., Lin X. (2004). Drosophila glypicans control the cell-to-cell movement of Hedgehog by a dynamin-independent process. *Development*. 131: 601-611.

Harfe B.D., Scherz P.J., Nissim S., Tian H., McMahon A.P., et al. (2004). Evidence for an expansion-based temporal Shh gradient in specifying vertebrate digit identities. *Cell*. 118: 517-528.

Harvey S.A., Smith J.C. (2009). Visualisation and quantification of morphogen gradient formation in the Zebrafish. *PLoS Biol*. 7: e1000101.

He F., Wen Y., Cheung D., Deng J., Lu L.J., et al. (2010). Distance measurements via the morphogen gradient of Bicoid in Drosophila embryos. *BMC Dev Biol*. 10: 80.

He F., Wen Y., Deng J., Lin X., Lu L.J., et al. (2008). Probing intrinsic properties of a robust morphogen gradient in Drosophila. *Dev Cell*. 15: 558-567.

Hepker J., Blackman R.K., Holmgren R. (1999). Cubitus interruptus is necessary but not sufficient for direct activation of a wing-specific decapentaplegic enhancer. *Development*. 126: 3669-3677.

Ho K.S., Suyama K., Fish M., Scott M.P. (2005). Differential regulation of Hedgehog target gene transcription by Costal2 and Suppressor of Fused. *Development*. 132: 1401-1412.

- Holloway D.M, Harrison L.G, Kosman D, Vanario-Alonso C.E, Spirov A.V. (2006). Analysis of pattern precision shows that *Drosophila* segmentation develops substantial independence from gradients of maternal gene products. *Dev Dyn.* 235: 2949-2960.
- Horstadius S. (1939). The mechanics of sea urchin development, studied by operative method. *Biol Rev.* 14: 132–179.
- Houchmandzadeh B., Wieschaus E., Liebler S. (2002). Establishment of developmental precision and proportions in the early *Drosophila* embryo. *Nature.* 415: 798-802.
- Hsiung F., Ramirez-Weber F.A., Iwaki D.D., Kornberg T.B. (2005). Dependence of *Drosophila* wing imaginal disc cytonemes on Decapentaplegic. *Nature.* 437: 560–563.
- Huang A.M., Rusch J., Levine M. (1997). An anteroposterior Dorsal gradient in the *Drosophila* embryo. *Genes Dev.* 11: 1963-1973.
- Huangfu D., Liu A., Rakeman A.S. , Murcia N.S., Niswander L., Anderson K.V. (2003). Hedgehog signalling in the mouse requires intraflagellar transport proteins. *Nature.* 426: 83-87.
- Huxley J.S., de Beer G.R. (1934). *The Elements of Experimental Embryology.* Cambridge University Press. Cambridge, UK.
- Ibañes M., Izpisúa-Belmonte J.C. (2008). Theoretical and experimental approaches to understand morphogen gradients. *Mol Syst Biol.* 4: 176.
- Imler J.L., Zheng L. (2004). Biology of Toll receptors: lessons from insects and mammals. *J Leukoc Biol.* 75: 18-26.

Ingham P.W., McMahon A.P. (2001). Hedgehog signaling in animal development: paradigms and principles. *Genes Dev.* 15: 3059-3087.

Ip Y.T., Park R.E., Kosman D., Bier E., Levine M. (1992). The dorsal gradient morphogen regulates stripes of rhomboid expression in the presumptive neuroectoderm of the *Drosophila* embryo. *Genes Dev.* 6: 1728-1739.

Jacob F., Monod J. (1961). Genetic regulatory mechanisms in the synthesis of proteins. *J Mol Biol.* 3: 318-356.

Jaeger J., Reinitz J. (2006). On the dynamic nature of positional information. *Bioessays*. 28: 1102-11.

Jaeger J., Surkova S., Blagov M., Janssens H., Kosman D., et al. (2004) Dynamic control of positional information in the early *Drosophila* embryo. *Nature*. 430: 368–371.

Jazwinska A., Rushlow C., Roth S. (1999). The role of brinker in mediating the graded response to Dpp in early *Drosophila* embryos. *Development*. 126: 3323–3334.

Jaźwińska A., Rushlow C., Roth S. (1999). The role of brinker in mediating the graded response to Dpp in early *Drosophila* embryos. *Development*. 126: 3323-3334.

Jeong J., McMahon A.P. (2005). Growth and pattern of the mammalian neural tube are governed by partially overlapping feedback activities of the hedgehog antagonists Patched1 and Hhip1. *Development*. 132: 143-154.

Jia J., Tong C., Wang B., Luo L., Jiang J. (2004). Hedgehog signalling activity of Smoothened requires phosphorylation by protein kinase A and casein kinase I. *Nature* 432: 1045-1050.

Jiang J., Hui C.C. (2008). Hedgehog signaling in development and cancer. *Dev Cell.* 15: 801-812.

Jiang J., Kosman D., Ip Y.T., Levine M. (1991). The dorsal morphogen gradient regulates the mesoderm determinant twist in early *Drosophila* embryos. *Genes Dev.* 5: 1881-1891.

Johnson R.L., Grenier J.K., Scott M.P. (1995). patched overexpression alters wing disc size and pattern: transcriptional and post-transcriptional effects on hedgehog targets. *Development* 121: 4161-4170.

Kanodia J.S., Rikhy R., Kim Y., Lund V.K., DeLotto R., et al. (2009). Dynamics of the Dorsal morphogen gradient. *Proc Natl Acad Sci USA.* 106: 21707-21712.

Kent D., Bush E.W., Hooper J.E. (2006). Roadkill attenuates Hedgehog responses through degradation of Cubitus interruptus. *Development.* 133: 2001-2010.

Kicheva A., González-Gaitán M. (2008). The Decapentaplegic morphogen gradient: a precise definition. *Curr Opin Cell Biol.* 20: 137-143.

Kicheva A., Pantazis P., Bollenbach T., Kalaidzidis Y., Bittig T., Jülicher F., González-Gaitán M. (2007). Kinetics of morphogen gradient formation. *Science.* 315: 521-525.

Kitano H. (2002). Systems biology: a brief overview. *Science.* 295: 1662-1664.

Kondo S., Asai R. (1995). A reaction-diffusion wave on the skin of the marine angelfish *Pomacanthus*. *Nature.* 376: 765-768.

Kornberg T.B., Guha A. (2007). Understanding morphogen gradients: a problem of dispersion and containment. *Curr Opin Genet Dev.* 17: 264-271.

- Kreutz C., Timmer J. (2009). Systems biology: experimental design. *FEBS J.* 276: 923-942.
- Kruger U., Zhang J., Xie L. (2007). Developments and applications of nonlinear Principal Component Analysis: A review. In: *Principal manifolds for data visualization and dimension-reduction*, Gorban A., et al. (eds.) pp. 1-44. Berlin-Heidelberg: Springer.
- Kutejova E., Briscoe E., Kicheva A. (2009). Temporal dynamics of patterning by morphogen gradients. *Curr Opin Genet Dev.* 19: 315-322.
- Lai K., Robertson M.J., Schaffer D.V. (2004). The sonic hedgehog signaling system as a bistable genetic switch. *Biophys J.* 86: 2748–2757.
- Lander A.D., Nie Q., Wan F.Y. (2002). Do morphogen gradients arise from diffusion? *Dev Cell.* 2: 785-796.
- Lander A.D. (2007). Morpheus unbound: reimagining the morphogen gradient. *Cell.* 128: 245-256.
- Lawrence P.A. (1966). Development and determination of hairs and bristles in the milkweed bug, *Oncopeltus fasciatus* (Lygaeidae, Hemiptera). *J Cell Sci.* 1: 475-498.
- Lawrence P.A. (1992). *The Making of a Fly*. Blackwell Publishing. Oxford, UK.
- Lawrence P.A. (2001). Morphogens: how big is the big picture? *Nat Cell Biol.* 3: E151-154.
- Lawrence P.A., Struhl G. (1996). Morphogens, compartments, and pattern: lessons from *Drosophila*? *Cell.* 85: 951–961.

Levine M., Davidson E.H. (2005). Gene regulatory networks for development. Proc Natl Acad Sci U S A. 102: 4936-4942.

Lewis J., Slack J.M., Wolpert L. (1977). Thresholds in development. J Theor Biol. 65: 579–590.

Lberman L.M., Reeves G.T., Stathopoulos A. (2009). Quantitative imaging of the Dorsal nuclear gradient reveals limitations to threshold-dependent patterning in *Drosophila*. Proc Natl Acad Sci USA. 106: 22317-22322.

Lott S.E., Kreitman M., Palsson A., Alekseeva E., Ludwig M.Z. (2007). Canalization of segmentation and its evolution in *Drosophila*. Proc Natl Acad Sci USA. 104: 10926-10931.

Ma C., Zhou Y., Beachy P.A., Moses K. (1993). The segment polarity gene hedgehog is required for progression of the morphogenetic furrow in the developing *Drosophila* eye. Cell. 75: 927-938.

McGuire S.E., Roman G., Davis R.L. (2004). Gene expression systems in *Drosophila*: a synthesis of time and space. Trends Genet. 20: 384-391.

McHale P., Rappel W.J., Levine H. (2006). Embryonic pattern scaling achieved by oppositely directed morphogen gradients. Phys Biol. 3: 107-120.

Meinhardt H. (1978). Space-dependent cell determination under the control of a morphogen gradient. J Theor Biol. 74: 307-321.

Meinhardt, H. (1983). Cell determination boundaries as organizing regions for secondary

embryonic fields. *Dev Biol.* 96: 375-385.

Meinhardt H. (2004). Different strategies for midline formation in bilaterians. *Nat Rev Neurosci.* 5: 502-510.

Methot N., Basler K. (1999). Hedgehog controls limb development by regulating the activities of distinct transcriptional activator and repressor forms of Cubitus interruptus. *Cell.* 96: 819-831.

Methot N., Basler K. (2001). An absolute requirement for Cubitus interruptus in Hedgehog signaling. *Development.* 128: 733-742.

Morata G., Lawrence P.A. (1975). Control of compartment development by the engrailed gene in Drosophila. *Nature.* 255: 614-617.

Morgan T.H. (1934). *Embryology and Genetics.* Columbia University Press. New York, NY.

Morgan T.H. (1897). Regeneration in oligochaete worms. *Science.* 6: 692-693.

Morisato D. (2001). Spätsle regulates the shape of the Dorsal gradient in the Drosophila embryo. *Development.* 128: 2309-2319.

Moussian B., Roth S. (2005). Dorsoventral axis formation in the Drosophila embryo--shaping and transducing a morphogen gradient. *Curr Biol.* 15: 887-899.

Müller B., Basler K. (2000). The repressor and activator forms of Cubitus interruptus control Hedgehog target genes through common generic gli-binding sites. *Development.* 127: 2999-3007.

- Mullor J.L., Calleja M., Capdevila J., Guerrero I. (1997). Hedgehog activity, independent of decapentaplegic, participates in wing disc patterning. *Development*. 124: 1227-1237.
- Mullor J.L., Guerrero I. (2000). A gain-of-function mutant of patched dissects different responses to the hedgehog gradient. *Dev Biol*. 228: 211-224.
- Nahmad M., Stathopoulos A. (2009). Dynamic interpretation of Hedgehog signaling in the Drosophila wing disc. *PLoS Biol*. 7: 1000202; DOI:10.1371/journal.pbio.1000202.
- Nakagoshi H., Shirai T., Nabeshima Y., Matsuzaki F. (2002). Refinement of wingless expression by a wingless- and notch-responsive homeodomain protein, defective proventriculus. *Dev Biol*. 249: 44-56.
- Nedham J. (1942). Biochemistry and Morphogenesis. Cambridge University Press. Cambridge, UK.
- Nellen D., Burke R., Struhl G., Basler K. (1996). Direct and long-range action of a DPP morphogen gradient. *Cell*. 85: 357-368.
- Nilson L.A., Schüpbach T. (1998). Localized requirements for windbeutel and pipe reveal a dorsoventral prepattern within the follicular epithelium of the Drosophila ovary. *Cell*. 93: 253-262.
- Nusslein-Volhard C., Wieschaus E. (1980). Mutations affecting segment number and polarity in Drosophila. *Nature*. 287: 795–801.
- Oates A.C., Gorfinkiel N., González-Gaitán M., Heisenberg C.P. (2009). Quantitative approaches in developmental biology. *Nat Rev Genet*. 10: 517-530.
- Pan D.J., Huang J.D., Courey A.J. (1991). Functional analysis of the Drosophila twist

promoter reveals a dorsal-binding ventral activator region. *Genes Dev.* 5: 1892-1901.

Panakova D., Sprong H., Marois E., Thiele C., Eaton S. (2005). Lipoprotein particles are required for Hedgehog and Wingless signalling. *Nature*. 435: 58–65.

Peri F., Technau M., Roth S. (2002). Mechanisms of Gurken-dependent pipe regulation and the robustness of dorsoventral patterning in *Drosophila*. *Development*. 129: 2965-2975.

Price J.V., Clifford R.J., Schüpbach T. (1989) The maternal ventralizing locus torpedo is allelic to faint little ball, an embryonic lethal, and encodes the *Drosophila* EGF receptor homolog. *Cell*. 56: 1085-1092.

Reach M., Galindo R.L., Towb P., Allen J.L., Karin M., Wasserman S.A. (1996). A gradient of cactus protein degradation establishes dorsoventral polarity in the *Drosophila* embryo. *Dev Biol*. 180: 353-364.

Reeves G.T., Stathopoulos A. (2009). Graded dorsal and differential gene regulation in the *Drosophila* embryo. *Cold Spring Harb Perspect Biol*. 1: a000836.

Reversade B., De Robertis E.M. (2005). Regulation of ADMP and BMP2/4/7 at opposite embryonic poles generates a self-regulating morphogenetic field. *Cell*. 123: 1147-1160.

Riddle R.D., Johnson R.L., Laufer E., Tabin C. (1993). Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell*. 75: 1401–1416.

Roberts P. (1964). Mosaics Involving Aristapedia, a Homeotic Mutant of *Drosophila melanogaster*. *Genetics*. 49: 593-598.

Robertson F.W. (1963). The ecological genetics of growth in *Drosophila*. VI. The genetic correlation between the duration of the larval period and body size in relation to larval diet. *Gene Res Camb.* 4: 74-92.

Roelink H., Porter J.A., Chiang C., Tanabe Y., Chang D.T., et al. (1995). Floor plate and motor neuron induction by different concentrations of the amino-terminal cleavage product of sonic hedgehog autoproteolysis. *Cell.* 81: 445-455.

Roth S., Stein D., Nüsslein-Volhard C. (1989). A gradient of nuclear localization of the dorsal protein determines dorsoventral pattern in the *Drosophila* embryo. *Cell.* 59: 1189-1202.

Rushlow C.A., Han K., Manley J.L., Levine M. (1989). The graded distribution of the dorsal morphogen is initiated by selective nuclear transport in *Drosophila*. *Cell.* 59: 1165-1177.

Rushlow C., Frasch M., Doyle H., Levine M. (1987). Maternal regulation of *zerknüllt*: a homoeobox gene controlling differentiation of dorsal tissues in *Drosophila*. *Nature.* 330: 583-586.

Saha K., Schaffer D.V. (2006). Signal dynamics in Sonic Hedgehog tissue patterning. *Development.* 133: 889-900.

Sander K. (1994). Of gradients and genes: Developmental concepts of Theodor Boveri and his students. *Dev Genes Evol.* 203: 295-297.

Schejter E.D., Shilo B.Z. (1989). The *Drosophila* EGF receptor homolog (DER) gene is allelic to faint little ball, a locus essential for embryonic development. *Cell.* 56: 1093-

Scherz P.J., McGlinn E., Nissim S., Tabin C.J. (2007). Extended exposure to Sonic hedgehog is required for patterning the posterior digits of the vertebrate limb. *Dev Biol.* 308: 343-354.

Sen J., Goltz J.S., Stevens L., Stein D. (1998). Spatially restricted expression of pipe in the *Drosophila* egg chamber defines embryonic dorsal-ventral polarity. *Cell.* 95: 471-81.

Shimizu K., Gurdon J.B. (1999). A quantitative analysis of signal transduction from activin receptor to nucleus and its relevance to morphogen gradient interpretation. *Proc Natl Acad Sci U S A.* 96: 6791-6796.

Shinar G., Feinberg M. (2010). Structural sources of robustness in biochemical reaction networks. *Science.* 327: 1389-1391.

Smith J., Thoedoris C., Davidson E.H. (2007). A gene regulatory network subcircuit drives pattern of gene expression. *Science.* 318: 794-797.

Sondhi K.C. (1963). The biological foundations of animal patterns. *Q Rev Biol.* 38: 289-327.

Spemann H. (1938). *Embryonic Development and Induction*. Yale University Press. New Haven, CT.

Spemann H, Mangold H. (1924). Induction of embryonic primordia by implantation of organizers from a different species. *Roux's Arch Entw Mech.* 100: 599-638.

Stathopoulos A., Levine M. (2002). Linear signaling in the Toll-Dorsal pathway of

Drosophila: activated Pelle kinase specifies all threshold outputs of gene expression while the bHLH protein Twist specifies a subset. Development. 129: 3411-3419.

Stathopoulos A., Levine M. (2005). Genomic regulatory networks and animal development. Dev Cell. 9:449-462.

Stern C. (1954). Genes and developmental patterns. Proc. 9<sup>th</sup> Internat Cong Genetics. Part 1. pp. 355-369.

Steward R. (1989). Relocalization of the dorsal protein from the cytoplasm to the nucleus correlates with its function. Cell. 59: 1179-1188.

Strigini M., Cohen S.M. (1997). A Hedgehog activity gradient contributes to AP axial patterning of the Drosophila wing. Development. 124: 4697-4705.

Strutt H., Thomas C., Nakano Y., Stark D., Neave B., et al. (2001). Mutations in the sterol-sensing domain of Patched suggest a role for vesicular trafficking in Smoothened regulation. Curr Biol. 11: 608-613.

Stumpf H.F. (1966). Mechanism by which cells estimate their location within the body. Nature. 212: 430-431.

Stumpf H.F. (1967). On positional determination of cuticle zones within a segment of *Galleria mellonella*. Dev Biol. 16: 144-167.

Sturmfels B. (2005). Can biology lead to new theorems? Clay Mathematics Institute Annual Report .

Su V.F., Jones K.A., Brodsky M., The I. (2007). Quantitative analysis of Hedgehog gradient formation using an inducible expression system. *BMC Dev Biol.* 7: 43.

Tabata T., Takei Y. (2004). Morphogens, their identification and regulation. *Development.* 131: 703-712.

Taipale J., Cooper M.K., Maiti T., Beachy P.A. (2002). Patched acts catalytically to suppress the activity of Smoothened. *Nature.* 418: 892-897.

Tanimoto H., Itoh S., ten Dijke P., Tabata T. (2000). Hedgehog creates a gradient of DPP activity in Drosophila wing imaginal discs. *Mol Cell.* 5: 59-71.

Thompson C.R., Reichelt S., Kay R.R. (2004). A demonstration of pattern formation without positional information in Dictyostelium. *Dev Growth Differ.* 46: 363-369.

Tokunaga C. (1978). Genetic mosaic studies in *Drosophila melanogaster*, with special reference to the prepattern hypothesis. In: *Genetic Mosaics and Cell Differentiation.* Gehring W.J. (ed.). Springer-Verlag. Berlin, Germany. pp. 157-204.

Tomlin C.J, Axelrod J.D. (2007). Biology by numbers: mathematical modelling in developmental biology. *Nat Rev Genet.* 8: 331-340.

Torroja C., Gorfinkel N., Guerrero I. (2004). Patched controls the Hedgehog gradient by endocytosis in a dynamin-dependent manner, but this internalization does not play a major role in signal transduction. *Development.* 131: 2395-2408.

Towb P., Galindo R.L., Wasserman S.A. (1998). Recruitment of Tube and Pelle to signaling sites at the surface of the *Drosophila* embryo. *Development.* 125: 2443-2550.

Towers M., Mahood R., Yin Y., Tickle C. (2008). Integration of growth and specification in chick wing digit-patterning. *Nature*. 452: 882-886.

Turing A.M. (1952). The chemical basis of morphogenesis. *Phil Trans R Soc Lond B Biol Sci*. 237: 37-72.

Varjosalo M., Taipale J. (2008). Hedgehog: functions and mechanisms. *Genes Dev*. 22: 2454-2472.

Vervoort M. (2000). Hedgehog and wing development in Drosophila: a morphogen at work? *Bioessays*. 22: 460-468.

Vervoort M., Crozatier M., Valle D., Vincent A. (1999). The COE transcription factor Collier is a mediator of short-range Hedgehog-induced patterning of the Drosophila wing. *Curr Biol*. 9: 632-639.

von Dassow G., Meir E., Munro E.M., Odell G.M. (2000). The segment polarity network is a robust developmental module. *Nature*. 406: 188-192.

von Ohlen T., Doe C.Q. (2000). Convergence of dorsal, dpp, and egfr signaling pathways subdivides the drosophila neuroectoderm into three dorsal-ventral columns. *Dev Biol*. 224: 362-372.

Waddington C.H. (1940). Organisers and Genes. Cambridge University Press. Cambridge, UK.

Waddington C.H. (1956). Principles of Embryology. George Allen & Unwin. London, UK.

Witkowski J. (1985). The hunting of the organizer: an episode in biochemical embryology. *TIBS*. 10: 379–381.

Wolpert L. (1968). The French Flag problem: a contribution to the discussion on pattern development and regulation. In: Waddington C.H. (ed.). *Towards a theoretical biology*. Edinburgh University Press. Edinburgh, UK. pp. 125–133.

Wolpert L. (1969). Positional information and the spatial pattern of cellular differentiation. *J Theor Biol*. 25: 1-47.

Wolpert L. (1971). Positional information and pattern formation. *Curr Top Dev Biol*. 6: 183-224.

Yi T. M., Huang Y., Simon M. I., Doyle J. (2000). Robust perfect adaptation in bacterial chemotaxis through integral feedback control. *Proc Natl Acad Sci USA*. 97: 4649-4653.

Yu S.R., Burkhardt M., Nowak M., Ries J., Petrásek Z., et al. (2009). Fgf8 morphogen gradient forms by a source-sink mechanism with freely diffusion molecules. *Nature*. 461: 533-536.

Zhang Q., Zhang L., Wang B., Ou C.Y., Chien C.T., Jiang J. (2006). A hedgehog-induced BTB protein modulates hedgehog signaling by degrading Ci/Gli transcription factor. *Dev Cell*. 10: 719-729.

Zhao Y., Tong C., Jiang J. (2007). Hedgehog regulates smoothened activity by inducing a conformational switch. *Nature*. 450: 252-258.

Zhu A.J., Scott M.P. (2004). Incredible journey: how do developmental signals travel through tissue? *Genes Dev*. 18: 2985-2997.