

## Chapter 3

### *Synthesis and Biological Activity of Cyclic Pyrrole-Imidazole Polyamide Libraries*

*The text of this chapter was taken in part from a manuscript co-authored with Benjamin C. Li, James W. Puckett, and Peter B. Dervan (California Institute of Technology)*

(Li, B. C.; Montgomery, D. C.; Puckett, J. W.; Dervan, P. B. "Synthesis of Cyclic Py-Im Polyamide Libraries," *J. Org. Chem.* **2013**, 78, 124-133)

## Abstract

Cyclic Py-Im polyamides containing two GABA turn units exhibit enhanced DNA binding affinity, but extensive studies of their biological properties have been hindered due to synthetic inaccessibility. A facile modular approach towards cyclic polyamides has been developed via microwave-assisted solid-phase synthesis of hairpin amino acid oligomer intermediates followed by macrocyclization. A focused library of cyclic polyamides **1-7** targeted to the androgen response element (ARE) and the estrogen response element (ERE) were synthesized in 12-17% overall yield. The Fmoc protection strategy also allows for selective modifications on the GABA turn units that have been shown to improve cellular uptake properties. The DNA binding affinities of a library of cyclic polyamides were measured by DNA thermal denaturation assays and compared to the corresponding hairpin polyamides. Fluorescein-labeled cyclic polyamides have been synthesized and imaged via confocal microscopy in A549 and T47D cell lines. The IC<sub>50</sub> values of compounds **1-7** and **9-11** were determined, revealing remarkably varying levels of cytotoxicity.


## Introduction

The selective modulation of eukaryotic gene expression by small molecules may have important implications in the field of chemical biology and human medicine. Pyrrole-imidazole polyamides are a class of synthetic ligands that can be programmed to bind the minor groove of specific DNA sequences.<sup>1</sup> Antiparallel, side-by-side N-methylpyrrole (Py) and N-methylimidazole (Im) carboxamides (Im/Py) pairs distinguish G·C from C·G base pairs, N-methyl-3-hydroxypyrrole (Hp)/Py shows specificity for T·A over A·T, whereas Py/Py pairs are specific for both T·A and A·T.<sup>2-5</sup> By linking two strands of these heterocyclic oligomers via a  $\gamma$ -amino butyric acid (GABA) turn unit, hairpin Py-Im polyamides can be programmed to bind a large library of DNA sequences with affinities comparable to natural DNA-binding proteins.<sup>6-8</sup> Hairpin polyamides have been shown to localize to the nuclei of living cells, and regulate endogenous gene expression by disrupting protein/DNA interfaces.<sup>9-17</sup> Cyclic polyamides containing a second GABA turn unit exhibit further enhanced DNA binding properties.<sup>18-21</sup> We have recently demonstrated their gene regulatory effects on AR-activated gene expression in prostate cancer models.<sup>22</sup>

This discovery has opened a new area of research towards transcriptional regulation with small molecules, but the relative synthetic inaccessibility of cyclic polyamides has remained a bottleneck for examining libraries of structural variants that would modulate affinity, cell uptake and biological activity. Initial solid-phase methods were low yielding, and required substantial pre-modifications of the PAM resin.<sup>18-20</sup> While the solution-phase synthesis of cyclic polyamides remains useful in large-scale target-oriented synthesis, it has limited practicality towards libraries for screening

biological activities.<sup>22</sup> The recent report by Morinaga et al. offers a modular approach to achieve cyclic polyamides by intramolecular coupling of a cysteine and a chloroacetyl residue, but the modification of the optimal three-carbon GABA turn into a sulfur-containing four-atom linker compromises its DNA binding affinity and may alter its biological properties.<sup>23</sup>

We report here a solid-phase polyamide synthesis of a key hairpin amino acid oligomer intermediate, which followed by intramolecular cyclization, affords cyclic polyamides **1-8** in good yields. The polyamides were synthesized step-wise on 2-chlorotrityl resin. The modular approach led to rapid access of a focused library of cyclic polyamides **1-7** with various core sequences and turn unit modifications. The utilization of Fmoc chemistry allowed for differentially protected turn units, which were modified selectively to complement existing cellular imaging and cell uptake enhancement technologies.<sup>24</sup> We examined the DNA binding properties and cytotoxicity profiles of compounds **1-11**, and the cellular localization of cyclic polyamides **12-14** by fluorescence microscopy.



**Figure 3.1:** The basic strategy for synthesis of differentially protected cyclic Py-Im polyamides.

## Results and Discussion

### *Microwave-Assisted Solid-Phase Synthesis*


Due to previously observed decomposition of the conjugated C-terminal free carboxylic acid in polyamide intermediates, 2-chlorotrityl-chloride (2-Cl-Trt-Cl) resin was chosen for its mild synthesis and cleavage conditions. Polyamide synthesis on this resin has been previously reported by Aldrich-Wright and co-workers, but a resin-bound  $\beta$ -alanine linker was used in both instances, and a new loading procedure was therefore needed.<sup>25,26</sup> 2-Cl-Trt-Cl resin was first loaded with the Fmoc-protected Py monomer in N,N-dimethylformamide (DMF) and capped with methanol. Resin substitution levels were determined by the Fmoc test and confirmed by weighing the dry mass of the loaded resin. Fmoc deprotection was achieved using a 50% piperidine in DMF solution. In light of the recent improvements in both efficiency and yield, the couplings were performed under microwave-assisted conditions using the desired PyBOP-activated monomers.<sup>27</sup>

Resin-Bound Nucleophile	Deprotection Times <sup>a</sup>	Coupling Times <sup>b</sup> (min)		
		Py	Im	GABA/ $\beta$ -Ala
Py	3 x 10min	20	20	20
Im	3 x 10min	IR <sup>c</sup>	30	30
GABA/ $\beta$ -Ala	2 x 5min	20	20	20

**Table 3.1** Standard Fmoc deprotection and microwave-assisted coupling times for solid-phase polyamide synthesis. <sup>a</sup>All deprotections were performed in 50% piperidine in DMF. <sup>b</sup>All coupling reactions were conducted under microwave-assisted conditions at 50 °C with a 0.3M solution of the activated monomers (3 eq. monomer acid, 3 eq. PyBOP, 8 eq. DIEA, DMF). <sup>c</sup>FmocPyOH coupling onto resin with N-terminal Im was incomplete even at 60°C for up to 1h. Synthesis of polyamide sequences that require this linkage should use the FmocPyImOH dimer instead, demonstrated later in the synthesis of **8**.

Initial syntheses performed at 60°C led to premature cleavage of intermediates off the 2-Cl-Trt resin, and 50°C couplings were therefore preferable. The challenging Im to Py coupling required an FmocPyImOH dimer, which was obtained via an optimized procedure

















by Weltzer and Wemmer.<sup>28</sup> The deprotection and coupling conditions are detailed in Table 3.1. This 2-step deprotection-coupling procedure was repeated until the desired polyamide sequence was achieved. To build the small library of polyamides in a modular fashion, the resin was split into different batches at corresponding steps for further derivatization. Upon completion, the N-terminal Fmoc-protected polyamide oligomer was cleaved from the resin with 30% hexafluoroisopropanol (HFIP) in dichloromethane (DCM), concentrated *in vacuo*, and the resulting residue was subjected to a 20% piperidine solution to remove the Fmoc group. Direct cleavage of the free-amine polyamide oligomer was attempted, but found to be ineffective due to poor solubility of the zwitterion intermediate in the cleavage solution. After purification by high-performance liquid chromatography (HPLC), the desired polyamide intermediates **18-24** were obtained in 31-40% yields.



**Scheme 3.1:** Microwave-assisted synthesis of cyclic polyamides **1-7**. <sup>1</sup>All PyBOP-mediated coupling conditions were performed under microwave-assisted conditions (see Table 3.1). <sup>2</sup>Reagents and conditions: (i) 50% piperidine, DMF; (ii) FmocPyOH, PyBOP, DIEA, DMF; (iii) 50% piperidine, DMF; (iv) FmocPyOH, PyBOP, DIEA, DMF; (v) 50% piperidine, DMF; (vi) FmocImOH, PyBOP, DIEA, DMF; (vii) 50% piperidine, DMF; (viii) Z-β-Dab(Fmoc)-OH (for **1-6**) or Boc-β-Dab(Fmoc)-OH (for **7**), PyBOP, DIEA, DMF; (ix) 50% piperidine, DMF; (x) FmocPyOH, PyBOP, DIEA, DMF; (xi) 50% piperidine, DMF; (xii) FmocPyOH, PyBOP, DIEA, DMF; (xiii) 50% piperidine, DMF; (xiv) FmocPyOH (for **1-3**) or FmocImOH (for **4-7**), PyBOP, DIEA, DMF; (xv) 50% piperidine, DMF; (xvi) FmocImOH, PyBOP, DIEA, DMF; (xvii) 50% piperidine, DMF; (xviii) Z-β-Dab(Fmoc)-OH (for **1 & 4 & 7**) or Fmoc-GABA-OH (for **2 & 5**) or Boc-β-Dab(Fmoc)-OH (for **3 & 6**), PyBOP, DIEA, DMF; (xix) 30% HFIP, DCM; (xx) 20% piperidine, DMF; (xxi) DPPA, DIEA, DMF; (xxii) 10% TFMSA, TFA (for **1, 2, 4 & 5**) or TFA (for **3, 6 & 7**).

### DPPA-Mediated Macrocyclization

The polyamide macrocyclization step was achieved by a DPPA-mediated ring-closing reaction between the N-terminal amino group and the C-terminal carboxylic acid. This method was first employed by Cho et al. in the synthesis of cyclic polyamides.<sup>18,29,30</sup> In order to obtain a general work-up procedure applicable to polyamides of various lipophilicities, diisopropylethylamine (DIEA) was used as the base in place of sodium bicarbonate ( $\text{NaHCO}_3$ ) in the original conditions.<sup>31</sup> Deprotection of the turn units with trifluoromethanesulfonic acid (TFMSA) or trifluoroacetic acid (TFA), followed by HPLC purification, afforded polyamides **1-7** in **37-43%** yields over 2 steps.


Polyamide Precursor	$[\text{M}+\text{H}]^+$ (expected)	m/z (observed)	Solid-Phase Yield (%)	Cyclic Polyamide	$[\text{M}+\text{H}]^+$ (expected)	m/z (observed)	Cyclization Yield (%)	Overall Yield (%)
 18	1465.4	1465.9	33	 1	1179.5	1179.9	35	12
 19	1316.5	1316.9	40	 2	1164.5	1164.9	42	17
 20	1431.6	1453.9 [M+Na] <sup>+</sup>	34	 3	1313.6	1314.0	47	16
 21	1466.5	1466.9	31	 4	1180.5	1180.9	39	12
 22	1317.5	1317.2	34	 5	1165.5	1165.5	48	16
 23	1432.6	1455.0 [M+Na] <sup>+</sup>	32	 6	1314.5	1314.7	37	12
 24	1432.6	1454.9 [M+Na] <sup>+</sup>	33	 7	1314.5	1314.8	38	13
 25	1414.6	1372.0 [M-CO <sub>2</sub> ] <sup>+</sup>	34	 8	1129.5	1130.0	39	13

**Table 3.2:** Summary Table of MALDI-TOF Data and Synthetic Yields for Cyclic Polyamides **1-8** and Intermediates **18-25**.

### Selective Derivatization of Cyclic Polyamide Turn Units


By taking advantage of the Fmoc protection scheme, the two GABA  $\beta$ -amino groups in **3** were differentially protected. This is further highlighted in **6** and **7**, which

share the same asymmetric polyamide core targeted to the 5'-WGGWCW-3' sequence, and allowed for the selective conjugation of a benzoic acid moiety on a single turn unit in polyamides **9-11** that has been recently developed to enhance the cellular localization properties of hairpin polyamides.<sup>24</sup>



**Scheme 3.2:** Preparation of cyclic polyamides **9-11**. Reagents and conditions: (i) BzOH, PyBOP, DIEA, DMF; (ii) 10% TFMSA, TFA.

Mono-substituted benzyl carbamate (Cbz) polyamides **3**, **6** and **7** were chosen as targets based on unpublished results indicating Cbz-functionalized polyamides are biologically active (Supp. Fig. S16). Cyclic polyamides **12-14** with a fluorescein dye were synthesized in a similar fashion, and imaged in live cells via fluorescence microscopy. Furthermore, both the solubility and the pharmacokinetic profiles of cyclic polyamides can be highly dependent on subtle structural modifications, and this method allows for the modular synthesis of these structural variants in an efficient manner.<sup>31-33</sup>




**Scheme 3.3:** Preparation of FITC-labeled cyclic polyamides **12-14**. Reagents and conditions: (i) FITC, DIEA, DMF; (ii) 10% TFMSA, TFA.

### *Synthesis of Cyclic and Hairpin Polyamides with C-terminal Imidazole Units*

Previously established solid-phase polyamide synthesis methods have been generally limited to sequences beginning with a pyrrole monomer.<sup>34,35</sup> Solid-phase synthesis of polyamides starting with imidazole units on the commonly used Kaiser oxime resin have been low yielding, mainly attributed to the sensitivity of the oxime-imidazole linkage that lead to premature cleavage of resin-bound intermediates. The addition of an aliphatic linker (e.g. Boc- $\beta$ -Ala-PAM resin) circumvents this issue, but previous studies on hairpin polyamides with C-terminal  $\beta$ -alanine motifs have shown reduced cellular uptake properties and thus diminished gene regulatory effects of these products.<sup>9</sup> Using the microwave-assisted conditions reported above, cyclic polyamide **8** targeted to the 5'-

WCGWGW-3' sequence found in E-Box binding sites have been synthesized in 13% yield overall.




**Scheme 3.4:** Preparation of cyclic polyamide **8**. All PyBOP-mediated coupling conditions were performed under microwave-assisted conditions (see Table 3.1). Reagents and conditions: (i) 50% piperidine, DMF; (ii) FmocPyOH, PyBOP, DIEA, DMF; (iii) 50% piperidine, DMF; (iv) FmocPyOH, PyBOP, DIEA, DMF; (v) 50% piperidine, DMF; (vi) Z- $\beta$ -Dab(Fmoc)-OH, PyBOP, DIEA, DMF; (vii) 50% piperidine, DMF; (viii) FmocImOH, PyBOP, DIEA, DMF; (ix) 50% piperidine, DMF; (x) Fmoc- $\beta$ -Ala-OH, PyBOP, DIEA, DMF; (xi) 50% piperidine, DMF; (xii) FmocPyImOH, PyBOP, DIEA, DMF; (xiii) 50% piperidine, DMF; (xiv) Z- $\beta$ -Dab(Fmoc)-OH, PyBOP, DIEA, DMF; (xv) 30% HFIP, DCM; (xvi) 20% piperidine, DMF; (xvii) DPPA, DIEA, DMF; (xviii) 10% TFMSA, TFA.

Initial attempts starting with FmocImOH-loaded resin resulted in undesired cleavage during subsequent steps, and so syntheses began with loading of the FmocPyImOH dimer onto 2-chlorotrityl resin. After deprotection-coupling of the corresponding monomer units, followed by resin cleavage and Fmoc removal, polyamide

intermediate **25** was isolated by HPLC purification in 34% yield. DPPA-mediated cyclization of **25**, followed by Cbz deprotection afforded cyclic polyamide **8** in 39% yield over 2 steps.

Hairpin polyamide **17** was synthesized in a similar stepwise manner to afford the Im-capped, C-terminal acid intermediate, which was then coupled to a 3,3'-diamino-N-methyldipropylamine linker, followed by isophthalic acid conjugation, Boc deprotection, and isolation in 24% yield over 16 steps (Scheme 3.5). This is a step forward which allows for the synthesis of both cyclic and non- $\beta$ -alanine-linked hairpin polyamides with sequences beginning with an imidazole unit, further expanding the scope of targetable DNA sequences and inhibition of transcription factor-mediated gene expression by Py-Im polyamides.




**Scheme 3.5:** Preparation of hairpin polyamide **17**. All PyBOP-mediated coupling conditions were performed under microwave-assisted conditions (see Table 3.1). Reagents and conditions: (i) 50% piperidine, DMF; (ii) FmocPyOH, PyBOP, DIEA, DMF; (iii) 50% piperidine, DMF; (iv) FmocPyOH, PyBOP, DIEA, DMF; (v) 50% piperidine, DMF; (vi) Boc-b-Dab(Fmoc)-OH, PyBOP, DIEA, DMF; (vii) 50% piperidine, DMF; (viii) FmocImOH, PyBOP, DIEA, DMF; (ix) 50% piperidine, DMF; (x) Fmoc- $\beta$ -Ala-OH, PyBOP, DIEA, DMF; (xi) 50% piperidine, DMF; (xii) PyImOH, PyBOP, DIEA, DMF; (xiii) 30% HFIP, DCM; (xiv) 3,3'-diamino-N-methyldipropylamine, PyBOP, DMF; (xv) Isophthalic acid, PyBOP, DIEA, DMF; (xvi) TFA.

### *Thermal Stabilization of DNA Duplexes by Polyamides*

Py-Im polyamide-DNA binding affinities and specificities have historically been measured by quantitative DNase I footprinting assays.<sup>36</sup> As previously reported, however, cyclic polyamides have exceptionally high DNA binding affinities that exceed the detection limit of this experiment (i.e.  $K_a \geq 2 \times 10^{10} \text{ M}^{-1}$ ).<sup>22,37,38</sup> The DNA binding affinities of cyclic polyamides **1-14** have been rank ordered by magnitude of DNA thermal stabilization ( $\Delta T_m$ ), and compared to the corresponding hairpin polyamides **14-16**. Spectroscopic measurements were performed on 12-mer DNA duplexes with sequences 5'-CGATGTTCAAGC-3', 5'-CGATGGTCAAGC-3', and 5'-CGATCGTGAAGC-3', each containing a match binding site for the corresponding polyamides.


















Consistent with the findings of Chenoweth et al, the  $\Delta T_m$  value for bis-amino cyclic polyamide **1** ( $\Delta T_m = 26.1^\circ\text{C}$ ) was calculated to be significantly higher than that of hairpin polyamide **15** ( $\Delta T_m = 22.0^\circ\text{C}$ ).<sup>22</sup> So while **15** has an established binding affinity to the match androgen response element (ARE) half-site (5'-WGWWCW-3') in the subnanomolar range, cyclic polyamide **1** provides even greater stabilization to such DNA duplexes. Mono-unsubstituted cyclic polyamide **2** provides less DNA stabilization compared to **1** ( $\Delta T_m = 20.4^\circ\text{C}$ ), which is likely due to the loss of a positive charge and thus the loss of favorable electrostatic interactions with the negatively-charged DNA backbone. Perhaps more surprising is the high  $\Delta T_m$  values retained by mono-protected cyclic polyamides **3** ( $\Delta T_m = 27.3^\circ\text{C}$ ) and **9** ( $\Delta T_m = 28.0^\circ\text{C}$ ), each containing a lone free amino group and net +1 charge. As shown in Figure 3.2, the benzoyl (Bz) group in **9** projects straight down the minor groove, avoiding unfavorable steric interactions with the groove

wall, and may offer insight into the high degree of DNA stabilization by **3** and **9** comparable to the bis-amino cycle **1**.



**Figure 3.2** (Left) Stick model and (right) space-filling model of benzoyl-substituted turn along the DNA minor groove from three different angles (A-C). Based on published crystal structure by Chenoweth et al (PDB ID: 3OMJ).

Fluorescein-conjugate **12** ( $\Delta T_m = 21.8^\circ\text{C}$ ) has a DNA binding affinity lower than **3** and **9**, perhaps due to increased steric clashes from the larger substitution group and unfavorable electrostatic interactions from the negatively-charged fluorescein group, but still binds DNA at a similar level to **15**.

Polyamides		5'- CGA <b>TGGTCA</b> AGC -3'	
		$T_m / ^\circ\text{C}$	$\Delta T_m / ^\circ\text{C}$
—		53.1 ( $\pm 0.2$ )	—
	(1)	79.2 ( $\pm 0.2$ )	26.1
	(2)	73.5 ( $\pm 0.2$ )	20.4
	(3)	80.4 ( $\pm 0.2$ )	27.3
	(9)	81.1 ( $\pm 0.2$ )	28.0
	(12)	74.9 ( $\pm 0.1$ )	21.8
	(15)	75.1 ( $\pm 0.1$ )	22.0
Polyamides		5'- CGA <b>TGGTCA</b> AGC -3'	
		$T_m / ^\circ\text{C}$	$\Delta T_m / ^\circ\text{C}$
—		56.2 ( $\pm 0.3$ )	—
	(4)	79.4 ( $\pm 0.3$ )	23.2
	(5)	74.8 ( $\pm 0.2$ )	18.6
	(6)	79.8 ( $\pm 0.2$ )	23.6
	(7)	80.5 ( $\pm 0.3$ )	24.3
	(10)	82.2 ( $\pm 0.1$ )	26.0
	(11)	81.5 ( $\pm 0.1$ )	25.1
	(13)	78.4 ( $\pm 0.6$ )	22.2
	(14)	77.2 ( $\pm 0.1$ )	21.0
	(16)	72.9 ( $\pm 0.2$ )	16.7
Polyamides		5'- CGA <b>TCGTGA</b> AGC -3'	
		$T_m / ^\circ\text{C}$	$\Delta T_m / ^\circ\text{C}$
—		56.7 ( $\pm 0.2$ )	—
	(8)	70.1 ( $\pm 0.1$ )	13.4
	(17)	63.3 ( $\pm 0.3$ )	6.6

**Table 3.3:**  $T_m$  Values for Polyamide Library.






The magnitude of stabilization provided by cyclic polyamides **4-7**, **10-11**, and **13-14** targeted to estrogen response element (ERE) half-sites (5'-WGGWCW-3') follows a similar pattern to the aforementioned ARE-targeting series. Mono-unsubstituted cycle **5** ( $\Delta T_m = 18.6^\circ\text{C}$ ) stabilizes the duplex at a comparable level to hairpin polyamide **16** ( $\Delta T_m = 16.7^\circ\text{C}$ ), whereas bis-amino compound **4** ( $\Delta T_m = 23.2^\circ\text{C}$ ) has a higher  $\Delta T_m$  value. The mono-Cbz cycles **6** ( $\Delta T_m = 23.6^\circ\text{C}$ ) and **7** ( $\Delta T_m = 24.3^\circ\text{C}$ ), mono-Bz substituted **10** ( $\Delta T_m = 26.0^\circ\text{C}$ ) and **11** ( $\Delta T_m = 25.1^\circ\text{C}$ ), and mono-fluorescein conjugates **13** ( $\Delta T_m = 22.2^\circ\text{C}$ ) and **14** ( $\Delta T_m = 21.0^\circ\text{C}$ ) each bind DNA similar to **4**.

Cyclic polyamide **8** ( $\Delta T_m = 13.4^\circ\text{C}$ ) binds the match 5'-WCGWGW-3' sequence at an elevated level compared to hairpin **17** ( $\Delta T_m = 6.6^\circ\text{C}$ ), which may prove important towards targeting oncogenic transcription factors such as c-Myc that act through binding canonical E-Box (5'-CACGTG-3') sequences.

#### *Sulforhodamine B Cytotoxicity Assay for Compounds 1-11*








The cytotoxicities of compounds **1-11** were assessed in A549 human lung carcinoma and T47D human breast cancer cell lines (Tables 3.4 & 3.5). The cyclic polyamides targeted to the 5'-WGWWCW-3' sequence generally exhibit a higher level of cytotoxicity than the 5'-WGGWCW-3' series, which is consistent with the trends observed in the DNA thermal denaturation analysis and confocal microscopy studies. Detailed inspection of the  $\text{IC}_{50}$  values within the series of compounds and across the cell lines, however, offers some unanticipated insights into the different biological properties of these minor structural variants.

Bis-amino cycle **1**, which has previously been shown to be biologically active, did not display any significant level of cytotoxicity in either cell line ( $IC_{50} > 30 \mu M$ ). Mono-unsubstituted compound **2** ( $IC_{50} = 4.9 \mu M$ ) and mono-Bz **9** ( $IC_{50} = 1.0 \mu M$ ) were comparably cytotoxic to hairpin polyamide **15** ( $IC_{50} = 3.1 \mu M$ ) in A549 cells, but an order of magnitude more cytotoxic ( $IC_{50} = 74 \text{ nM}$  &  $79 \text{ nM}$ , respectively) than **15** ( $IC_{50} = 710 \text{ nM}$ ) in T47D cells. The mono-Cbz cycle **3** was consistently the most cytotoxic compound in both A549 ( $IC_{50} = 160 \text{ nM}$ ) and T47D ( $IC_{50} = 25 \text{ nM}$ ) cell lines.

Cyclic Polyamide	A549	T47D
	$IC_{50} (\mu M)$	$IC_{50} (\mu M)$
 <b>1</b>	>30	>30
 <b>2</b>	$4.9 \pm 1.8$	$0.074 \pm 0.011$
 <b>3</b>	$0.16 \pm 0.05$	$0.025 \pm 0.005$
 <b>9</b>	$1.0 \pm 0.2$	$0.079 \pm 0.003$
 <b>15</b>	$3.1 \pm 0.6$	$0.71 \pm 0.10$

**Table 3.4:** SRB cytotoxicity data on compounds **1-3**, **9**, and **15**, in A549 and T47D cells, 72 hr incubation.

For the 5'-WGGWCW-3' targeting polyamides, the only compound that exhibited an  $IC_{50}$  value lower than  $30 \mu M$  in A549 cells was the mono-Cbz compound **6** ( $IC_{50} = 1.9 \mu M$ ). In T47D cells, consistent with the 5'-WGWWCW-3' series, bis-amino cycle **4** ( $IC_{50} > 30 \mu M$ ) was found to be not significantly cytotoxic and **6** ( $IC_{50} = 460 \text{ nM}$ ) was the most cytotoxic compound. Mono-unsubstituted cycle **5** ( $IC_{50} = 0.82 \mu M$ ) again shares a comparable level of cytotoxicity with the reference hairpin **16** ( $IC_{50} = 1.1 \mu M$ ), whereas mono-Bz compounds **10** ( $IC_{50} = 13.3 \mu M$ ) and **11** ( $IC_{50} = 7.5 \mu M$ ) are both an order of magnitude less cytotoxic.

Cyclic Polyamide	A549	T47D
	IC <sub>50</sub> (μM)	IC <sub>50</sub> (μM)
 4	>30	>30
 5	>30	0.82 ± 0.21
 6	1.9 ± 0.2	0.46 ± 0.27
 7	>30	>30
 10	>30	13.3 ± 0.6
 11	>30	7.5 ± 0.7
 16	>30	1.1 ± 0.1

**Table 3.5:** SRB cytotoxicity data on compounds **4-7**, **10-11**, and **16**, in A549 and T47D cells, 72 hr incubation.

Perhaps most interestingly, mono-Cbz compound **7** did not exhibit observable levels of cytotoxicity ( $IC_{50} > 30 \mu M$ ) in either A549 or T47D cells. Considering that **7** is a regioisomer of **6**, where the Cbz group is simply swapped onto the other turn, and that **7** only differs from **3** by a single  $-CH$  to  $-N$  substitution, it is rather surprising that **7** is more than 15- to 65- fold less cytotoxic than **6** and at least 180- to 1200- fold less cytotoxic than **3** in the two examined cell lines. Given the comparable DNA stabilization properties between **6** and **7**, and their common core sequence, we would not have predicted this vast discrepancy in cytotoxicity.

This study has demonstrated the large and somewhat unpredictable effects in biological activity induced by small structural variations of cyclic polyamides. Based on our preliminary work, the aggregation and pharmacokinetic properties of polyamides also vary greatly depending on structural modifications.<sup>31-33</sup> All this combines to highlight the

importance of a fast and reliable method to generate focused libraries of cyclic polyamides for future research.


#### *Confocal Microscopy of Cyclic Polyamide-Fluorescein Conjugates 12-14*

To directly examine the cellular localization of cyclic polyamides, fluorescein conjugates **12-14** were synthesized and visualized in living cells via confocal microscopy (Figs. 2 & 3). The selective conjugation of a single fluorescein molecule not only helped retain a free amino group for solubility purposes, but also allowed for the qualitative comparison of the two fluorescein conjugates **13** and **14** that share the same asymmetric polyamide core.


In each of the cases examined, cyclic polyamides **12-14** appear to permeate through the cellular membrane and localize in the cell nucleus, which was confirmed by the co-localization with Hoechst 33258 DNA stain. For ease of qualitatively assessing compound uptake, all A549 images were taken at a 660 fluorescence gain level, and all T47D images were taken at 600 fluorescence gain. Comparing Figures 3.3 and 3.4, the fluorescence levels of compounds **12-14** in T47D cells are all significantly higher than in A549, and this is only further amplified by this difference in gain levels.

Compound **12** matched to the 5'-WGWWCW-3' sequence exhibits the highest level of nuclear localization in both cell lines. Among the two 5'-WGGWCW-3' targeting cycles, polyamide **14** qualitatively appears to have a relatively higher fluorescence signal in the cell nuclei in both A549 and T47D cells. This may help explain the cytotoxicity data reported above, where compounds **6** and **11** with Cbz and Bz substitutions on the same side

as the fluorescein in **14** consistently display larger biological effects than **7** and **10**, which are more structurally similar to cyclic polyamide **13**.



**Figure 3.3:** Confocal microscopy of cyclic polyamide-fluorescein conjugates **12** (top), **13** (middle), and **14** (bottom) in A549 cells. In order to confirm nuclear localization, the fluorescence panel (left) was compared with Hoechst 33258 staining (middle) and overlay (right).



**Figure 3.4:** Confocal microscopy of cyclic polyamide-fluorescein conjugates **12** (top), **13** (middle), and **14** (bottom) in T47D cells. In order to confirm nuclear localization, the fluorescence panel (left) was compared with Hoechst 33258 staining (middle) and overlay (right).

## Conclusions

We have described a modular solid-phase synthesis method, which, when combined with an established DPPA-mediated macrocyclization step, afforded cyclic polyamides in a high-yielding and time-efficient manner. Using this method, we have overcome previous limitations and synthesized both cyclic and hairpin polyamides that start with an imidazole unit. The binding affinities of all synthesized cycles have been assessed by DNA thermal denaturation assays and compare favorably to hairpin polyamides that bind their match DNA sequences at subnanomolar concentrations. Furthermore, the protection strategy of our method allows for selective modification of the GABA turn units, which we have used to rapidly generate a focused library of compounds. The cytotoxicity and uptake analysis of the cyclic polyamides revealed unexpected properties that further highlight the need for an efficient method to synthesize structural variants of cyclic polyamides for future studies.

## Materials and Methods

### *Reagents and Equipment*

2-Chlorotrityl chloride (2-Cl-Trt-Cl) resin was purchased from Bachem. FmocPyOH and FmocImOH monomers were purchased from Wako. PyBOP was purchased from NovaBioChem. Boc- $\beta$ -Dab(Fmoc)-OH was purchased from Peptides International. All DNA oligomers were purchased HPLC purified from Integrated DNA Technologies. Cell culture media was purchased from Gibco. Fetal bovine serum was purchased from Omega Scientific. Microwave-assisted coupling reactions were conducted on a Biotage Initiator Eight synthesizer.

### *Polyamide Quantification*

Polyamide concentrations were measured in 20% MeCN in 0.1% (v/v) aqueous TFA using an approximated extinction coefficient of  $69\,200\text{ M}^{-1}\text{ cm}^{-1}$  at  $\lambda_{\text{max}}$  near 310 nm, unless otherwise specified.<sup>32, 36</sup>

### *Monomer Loading onto 2-Cl-Trt Resin*

Prior to manual microwave-assisted synthesis, 2-Cl-Trt-Cl resin (1.0 g, 1.59 mmol/g) was first loaded by mixing with 576 mg (1.59 mmol, 1 eq.) of FmocPyOH monomer, followed by addition of 6 mL dimethylformamide (DMF) and 1.38 mL diisopropylethylamine (DIEA) (7.59 mmol, 5 eq.). The suspended mixture was stirred for 18 hr, then capped by addition of 1 mL methanol (MeOH) and stirred for 1 hr. The orange-colored, loaded resin was then collected on a fritted peptide synthesis vessel, washed with DMF (2x), MeOH (2x), DMF (2x), MeOH (2x), and diethyl ether (Et<sub>2</sub>O). (Owing to the

sensitivity of the 2-Cl-Trt resin towards hydrolysis/methanolysis, this final Et<sub>2</sub>O wash was found to be essential and all loaded resin was sealed and stored at -20°C.) The loading efficiency was quantitated via the Fmoc test, and confirmed by measuring the mass of the dried resin. Typical monomer loading was calculated to be 0.4-0.8 mmol/g.

#### *Microwave-Assisted Solid-Phase Synthesis (18-24)*

All solid-phase polyamide coupling reactions were performed manually on a Biotage Initiator Eight microwave synthesizer on a 200-500 mg scale of loaded resin. Prior to each monomer coupling reaction, the N-terminal Fmoc group was first removed in a piperidine solution. The Fmoc deprotections were performed in a fritted peptide synthesis vessel at room temperature, and the specific conditions for each N-terminal monomer are as follows:

N-Fmoc-Pyrrole / Imidazole: (a) swell resin in DCM; (b) wash with DMF; (c) add 50% piperidine in DMF; (d) shake suspension for 10 min; (e) wash with DMF; (f) repeat steps a-e twice.

N-Fmoc-GABA /  $\beta$ -Alanine: (a) swell resin in DCM; (b) wash with DMF; (c) add 50% piperidine in DMF; (d) shake suspension for 5 min; (e) wash with DMF; (f) repeat steps a-e once.

Following Fmoc removal, the resin was de-swelled in MeOH, washed with Et<sub>2</sub>O, dried *in vacuo*, and transferred to a microwave synthesis vessel as a dry powder. The corresponding monomer acid (3 eq.) was activated with PyBOP (3 eq.) and DIEA (6 eq.) in DMF (0.3M concentration of monomer), and added to the resin. The coupling reactions were then set up in the microwave reactor at 50°C for the time durations described in **Table**

**3.1.** After the listed microwave-assisted coupling times, the reaction mixture was filtered into a peptide synthesis vessel, and the collected resin was washed with DMF (3x), MeOH (3x), Et<sub>2</sub>O, and dried *in vacuo*. To ensure completion of each deprotection and coupling step, analytical HPLC spectra were taken by cleaving a small resin sample in 30% hexafluoroisopropanol (HFIP) in DCM.

The polyamide core was synthesized on 2-Cl-Trt resin in an iterative manner by repeating the deprotection-coupling procedures described above using the corresponding monomeric units. Upon completion of the sequence, 100-200 mg of the resin was suspended in 1 mL 30% HFIP in DCM and stirred for 1 hr to yield the crude N-terminal Fmoc-protected polyamide intermediate. The reaction mixture was then run through a cotton filter to remove the resin, and the filtrate was concentrated *in vacuo*. The residual oil was resuspended in 5 mL of a 1:1 MeOH:DCM mixture, and reconcentrated *in vacuo* to give an off-white/beige solid. To remove the N-terminal Fmoc group, the solid was redissolved in 800  $\mu$ L DMF, followed by addition of 200  $\mu$ L piperidine, and the solution was stirred for 30 min. Upon confirmation of complete deprotection by analytical HPLC, the solution was added to 4 mL 30% MeCN in 0.1% aqueous TFA. The precipitated 9-methylenefluorene side product was then removed by centrifugation, and washed twice with 2 mL 30% MeCN in 0.1% aqueous TFA. The combined aqueous solution was purified by reverse-phase HPLC and lyophilized to dryness to yield pre-cyclic polyamide intermediates **18-24**. All dried samples of **18-24** were stored at -20°C prior to DPPA-mediated macrocyclization.

Synthetic yields and MALDI-TOF Characterization Data for **18-24**:

**(18)**: 12.6  $\mu\text{mol}$  recovered (38.0  $\mu\text{mol}$  theoretical, 33% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{70}\text{H}_{77}\text{N}_{22}\text{O}_{15}^+ = 1465.4$ , observed = 1465.9.

**(19)**: 15.2  $\mu\text{mol}$  recovered (38.0  $\mu\text{mol}$  theoretical, 40% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{62}\text{H}_{70}\text{N}_{21}\text{O}_{13}^+ = 1316.5$ , observed = 1316.9.

**(20)**: 12.8  $\mu\text{mol}$  recovered (38  $\mu\text{mol}$  theoretical, 34% yield). MALDI-TOF  $[\text{M} + \text{Na}]^+$  calc'd for  $\text{C}_{67}\text{H}_{79}\text{N}_{22}\text{NaO}_{15}^+ = 1453.6$ , observed = 1453.9.

**(21)**: 15.1  $\mu\text{mol}$  recovered (49  $\mu\text{mol}$  theoretical, 31% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{69}\text{H}_{76}\text{N}_{23}\text{O}_{15}^+ = 1466.5$ , observed = 1466.9.

**(22)**: 17.2  $\mu\text{mol}$  recovered (51  $\mu\text{mol}$  theoretical, 34% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{61}\text{H}_{69}\text{N}_{22}\text{O}_{13}^+ = 1317.5$ , observed = 1317.2.

**(23)**: 16.2  $\mu\text{mol}$  recovered (51  $\mu\text{mol}$  theoretical, 32% yield). MALDI-TOF  $[\text{M} + \text{Na}]^+$  calc'd for  $\text{C}_{66}\text{H}_{77}\text{N}_{23}\text{NaO}_{15}^+ = 1454.5$ , observed = 1455.0.

**(24)**: 8.2  $\mu\text{mol}$  recovered (25  $\mu\text{mol}$  theoretical, 33% yield). MALDI-TOF  $[\text{M} + \text{Na}]^+$  calc'd for  $\text{C}_{66}\text{H}_{78}\text{N}_{23}\text{NaO}_{15}^+ = 1454.5$ , observed = 1454.9.

#### *DPPA-mediated Macrocyclization (1-7)*

The macrocyclization reactions were run on a 2  $\mu\text{mol}$  to 16  $\mu\text{mol}$  scale. Intermediates **18-24** were first dissolved in DMF (0.25 mM) in a round-bottom flask equipped with a magnetic stir bar, followed by addition of DIEA (200 eq.), and purged with argon for 15 min. Diphenylphosphoryl azide (DPPA) (50 eq.) was then added to the reaction mixture in a drop wise manner, while rapidly stirring. Upon full addition of the DPPA, the solution was allowed to react and stirred at room temperature for 16-20 hr. After

confirmation of reaction completion by analytical HPLC, the reaction mixture was concentrated *in vacuo*, and the resulting oil residue was dissolved in 3 mL MeCN and transferred to a 15 mL Falcon Tube. The MeCN was then removed with air flow and 3 mL 0.1% aqueous TFA was added to the remaining oil layer to yield an off-white suspension, which was isolated via centrifugation and lyophilized to dryness.

For reactions starting with **18**, **19**, **21**, and **22**, the lyophilized residue was submitted to 1 mL 10% trifluoromethanesulfonic acid (TFMSA) in TFA, stirred for 5 min, frozen in LN2, and thawed by layering 1 mL DMF. For reactions starting with **20**, **23**, and **24**, the lyophilized residue was submitted to 1 mL neat TFA, stirred for 15 min, frozen in LN2, and thawed by layering 1 mL DMF. All the thawed solutions were then diluted with 6 mL 0.1% aqueous TFA, purified by reverse-phase HPLC and lyophilized to dryness to yield cyclic polyamides **1-7**.

Synthetic yields and MALDI-TOF characterization data for **1-7**:

**(1)**: 4.5  $\mu\text{mol}$  recovered (12.9  $\mu\text{mol}$  theoretical, 35% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{54}\text{H}_{63}\text{N}_{22}\text{O}_{10}^+ = 1179.5$ , observed = 1179.9.

**(2)**: 0.84  $\mu\text{mol}$  recovered (2.0  $\mu\text{mol}$  theoretical, 42% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{54}\text{H}_{62}\text{N}_{21}\text{O}_{10}^+ = 1164.5$ , observed = 1164.6.

**(3)**: 3.2  $\mu\text{mol}$  recovered (6.7  $\mu\text{mol}$  theoretical, 47% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{62}\text{H}_{69}\text{N}_{22}\text{O}_{12}^+ = 1313.6$ , observed = 1314.0.

**(4)**: 3.1  $\mu\text{mol}$  recovered (8.0  $\mu\text{mol}$  theoretical, 39% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{53}\text{H}_{62}\text{N}_{23}\text{O}_{10}^+ = 1180.5$ , observed = 1180.9.

**(5):** 0.96  $\mu\text{mol}$  recovered (2.0  $\mu\text{mol}$  theoretical, 48% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{53}\text{H}_{61}\text{N}_{22}\text{O}_{10}^+ = 1165.5$ , observed = 1165.5.

**(6):** 3.7  $\mu\text{mol}$  recovered (10.0  $\mu\text{mol}$  theoretical, 37% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{61}\text{H}_{68}\text{N}_{23}\text{O}_{12}^+ = 1314.5$ , observed = 1314.5.

**(7):** 2.2  $\mu\text{mol}$  recovered (5.7  $\mu\text{mol}$  theoretical, 38% yield). MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{61}\text{H}_{68}\text{N}_{23}\text{O}_{12}^+ = 1314.5$ , observed = 1314.8.

### *Selective Conjugation of Benzoic Acid Derivatives (9-11)*

A solution of benzoic acid (3.0 mg, 0.025 mmol, 25 eq.) and PyBOP (13 mg, 0.025 mmol, 25 eq.) in DMF (0.5 mL) and DIEA (44  $\mu\text{L}$ , 0.25 mmol, 250 eq.) was stirred at room temperature for 10 min. The activated solution was then added to **3** (1.0  $\mu\text{mol}$ ) and stirred for 3 hr. After confirmation of complete reaction by analytical HPLC, 12 mL cold  $\text{Et}_2\text{O}$  was added to the reaction mixture and cooled at  $-20^\circ\text{C}$  for 16 hr. The precipitate was then isolated by centrifugation and allowed to air dry. The resulting residue was submitted to 1 mL 10% trifluoromethanesulfonic acid (TFMSA) in TFA, stirred for 5 min, frozen in LN<sub>2</sub>, and thawed by layering 1 mL DMF. The thawed solution was then diluted with 6 mL 0.1% aqueous TFA, purified by reverse-phase HPLC and lyophilized to dryness to yield cyclic polyamides **9** (684 nmol, 68% yield). Using the same procedure described above, starting with **6** (1.60  $\mu\text{mol}$ ) and **7** (750 nmol), yielded mono-benzoyl substituted cyclic polyamides **10** (1.05  $\mu\text{mol}$ , 67% yield) and **11** (367 nmol, 49% yield), respectively.

**(9):** MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{61}\text{H}_{67}\text{N}_{22}\text{O}_{11}^+ = 1283.5$ , observed = 1284.1.

**(10):** MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{60}\text{H}_{66}\text{N}_{23}\text{O}_{11}^+ = 1284.5$ , observed = 1284.5.

**(11):** MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{60}\text{H}_{66}\text{N}_{23}\text{O}_{11}^+ = 1284.5$ , observed = 1284.9.

*Synthesis of Cyclic Polyamide-Fluorescein Conjugates (12-14)*

A solution of fluorescein isothiocyanate (FITC) (2.7 mg, 7.0  $\mu$ mol, 25 eq.) in DMF (0.2 mL) and DIEA (12  $\mu$ L, 0.07 mmol, 250 eq.) was added to **3** (0.28  $\mu$ mol) and stirred for 2 hr. After confirmation of complete reaction by analytical HPLC, 12 mL cold Et<sub>2</sub>O was added to the reaction mixture and cooled at -20°C for 16 hr. The precipitate was then isolated by centrifugation and allowed to air dry. The resulting residue was submitted to 1 mL 10% trifluoromethanesulfonic acid (TFMSA) in TFA, stirred for 5 min, frozen in LN<sub>2</sub>, and thawed by layering 1 mL DMF. The thawed solution was then diluted with 6 mL 0.1% aqueous TFA, purified by reverse-phase HPLC and lyophilized to dryness to yield cyclic polyamide **12** (65 nmol, 23% yield). Using the same procedure described above, starting with **6** (0.40  $\mu$ mol) and **7** (0.40  $\mu$ mol), yielded cyclic polyamide-fluorescein conjugated **13** (345 nmol, 86% yield) and **14** (118 nmol, 29% yield), respectively.

**(12):** ESI-MS  $[M + H]^+$  calc'd for C<sub>75</sub>H<sub>74</sub>N<sub>23</sub>O<sub>15</sub>S<sup>+</sup> = 1568.6, observed = 1568.3.

**(13):** ESI-MS  $[M + H]^+$  calc'd for C<sub>74</sub>H<sub>73</sub>N<sub>24</sub>O<sub>15</sub>S<sup>+</sup> = 1569.5, observed = 1569.2.

**(14):** ESI-MS  $[M + H]^+$  calc'd for C<sub>74</sub>H<sub>73</sub>N<sub>24</sub>O<sub>15</sub>S<sup>+</sup> = 1569.5, observed = 1569.3.

*Synthesis of Cyclic Polyamide Targeted to 5'-WCGWG-3' Sequence (8)*

2-Cl-Trt-Cl resin (200 mg, 1.59 mmol/g) was first loaded with FmocPyImOH dimer (96 mg, 0.20 mmol), which was obtained from published procedures.<sup>13</sup> Experimental details were analogous to the monomer loading protocol reported above. The obtained Fmoc-Py-Im-(2-Cl-Trt) resin (265 mg, 0.59 mmol/g) was subjected to the previously described microwave-assisted solid-phase synthesis conditions to build the corresponding

polyamide sequence. *A quarter* of the resin (0.15  $\mu\text{mol}$  theoretical) was then cleaved and purified by reverse-phase HPLC to yield pre-cyclic polyamide intermediate **25** as an off-white powder (13.3  $\mu\text{mol}$ , 34% yield). The isolated **25** (2.0  $\mu\text{mol}$ ) was subjected to DPPA-mediated macrocyclization conditions analogous to that for compounds **1-7**, with the Cbz groups removed with 10% TFMSA in TFA, and purified by reverse-phase HPLC to afford cyclic polyamide **8** (773 nmol, 39% yield).

**(25):** MALDI-TOF  $[\text{M} - \text{CO}_2 + \text{H}]^+$  calc'd for  $\text{C}_{65}\text{H}_{675}\text{N}_{22}\text{O}_{13}^+ = 1371.6$ , observed = 1371.7.

**(8):** MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{61}\text{H}_{67}\text{N}_{22}\text{O}_{11}^+ = 1129.5$ , observed = 1130.0

#### *Synthesis of Hairpin Polyamide Targeted to 5'-WCGWGW-3' Sequence (17)*

Fmoc-Py-Im-(2-Cl-Trt) resin, obtained via the same procedure as for **8**, was subjected to the previously described microwave-assisted solid-phase synthesis conditions to build the corresponding polyamide sequence with an N-terminal PyImOH cap. The resin (70 mg, 0.039 mmol) was then suspended in 2 mL 30% HFIP in DCM, stirred for 1 hr, filtered, washed, and concentrated to afford the crude C-terminal free acid. The residue was then dissolved in DMF (1.8 mL) and added drop wise to a pre-activated solution of 3,3'-diamino-N-methyldipropylamine (252  $\mu\text{L}$ , 1.56 mmol, 40 eq.) and PyBOP (40 mg, 0.078 mmol, 2 eq.) in DMF (6.0 mL). After stirring for 1 hr and confirmation of complete conjugation by HPLC, the crude product was isolated by  $\text{Et}_2\text{O}$  precipitation, air dried, redissolved in 8 mL 15% MeCN: 0.1% aqueous TFA, and purified by reverse-phase HPLC to yield polyamide intermediate **26** as an off-white powder (11.8  $\mu\text{mol}$ , 30% yield).

The isolated **26** (3.1  $\mu\text{mol}$ ) was dissolved in DMF (200  $\mu\text{L}$ ) and added drop wise to a pre-activated solution of isophthalic acid (12 mg, 0.072 mmol) and PyBOP (5 mg, 9  $\mu\text{mol}$ ) in DMF (800  $\mu\text{L}$ ) and DIEA (13  $\mu\text{L}$ ). After stirring for 1 hr and confirmation of complete conjugation by HPLC, the reaction mixture was then precipitated in  $\text{Et}_2\text{O}$ , isolated by centrifugation, and air-dried. The remaining residue was then subjected to 500  $\mu\text{L}$  neat TFA, stirred for 15 min, frozen in  $\text{LN}_2$ , thawed by addition of 1 mL DMF, diluted with 6 mL 0.1% aqueous TFA, and purified by reverse-phase HPLC to afford cyclic polyamide **8** (3.1 nmol, 86% yield). (Note: Extinction coefficient for compounds **26** and **17** were established as  $40,000 \text{ M}^{-1} \text{ cm}^{-1}$  at  $\lambda_{\text{max}} = 302\text{nm}$ .)

**(26):** MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{58}\text{H}_{79}\text{N}_{22}\text{O}_{11}^+ = 1259.6$ , observed = 1260.1.

**(17):** MALDI-TOF  $[\text{M} + \text{H}]^+$  calc'd for  $\text{C}_{61}\text{H}_{75}\text{N}_{22}\text{O}_{12}^+ = 1307.6$ , observed = 1307.8.

### *Thermal Denaturation Analysis*

Melting temperature analysis was performed on a Varian Cary 100 spectrophotometer equipped with a thermo-controlled cell holder possessing a cell path length of 1 cm. A degassed aqueous solution of 10 mM sodium cacodylate, 10 mM KCl, 10 mM  $\text{MgCl}_2$ , and 5 mM  $\text{CaCl}_2$  at pH 7.0 was used as analysis buffer. DNA duplexes and polyamides were mixed in 1:1 stoichiometry to a final concentration of 2  $\mu\text{M}$  for each experiment. Prior to analysis, samples were heated to  $95^\circ\text{C}$  and cooled to a starting temperature of  $25^\circ\text{C}$  with a heating rate of  $5^\circ\text{C}/\text{min}$  for each ramp. Denaturation profiles were recorded at  $\lambda = 260 \text{ nm}$  from 25 to  $95^\circ\text{C}$  with a heating rate of  $0.5^\circ\text{C}/\text{min}$ . The reported melting temperatures were defined as the maximum of the first derivative of the denaturation profile.

### *Cell Culture*

Cell lines were cultured at 37°C under 5% CO<sub>2</sub> using standard cell culture and sterile techniques. Cell medium was supplemented with 10% fetal bovine serum. Ham's F-12K (Kaighn's) medium was used for A549 cells, and RPMI 1640 was used for T47D cells.

### *Confocal Microscopy*

For each experiment, cells were plated in 200 µL of the proper medium onto glass-bottom cell culture plates at a density of  $1 \times 10^5$  (A549) cells/mL, or  $1.5 \times 10^5$  cells/mL (T47D). Cells were grown for 24 hr, and media was replaced with fresh media containing polyamide to give a final DMSO concentration of 0.1%. Next, cells were incubated for 16 hr, followed by removal of media, washing, and addition of fresh media. Hoechst 33258 was added 2 hr prior to imaging. Imaging was performed at the Caltech Beckman Imaging Center using a Zeiss LSM 510 Meta NLO 2-photon inverted laser scanning microscope equipped with a 40x oil-immersion objective lens. Polyamide–fluorescein conjugates **12-14** were imaged in multi-track mode using 488 nm laser excitation at 15% output with a pinhole of 375 µm and a standard fluorescein filter set. Hoechst was imaged using 800 nm two-photon excitation with an HFT KP680 dichroic and a 390- to 465-nm bandpass filter with a fully open pinhole. All images were analyzed using Zeiss LSM Zen software.


### *Sulforhodamine B Cytotoxicity Assay*

For cytotoxicity assays, cell lines were plated in 96-well cell culture plates in 100 µL media at a density of  $1 \times 10^4$  cells/mL (A549), or  $5 \times 10^4$  cells/mL (T47D). IC<sub>50</sub> values were determined using the sulforhodamine B (SRB) colorimetric assay as previously


described.<sup>39</sup> Cells were grown for 24 hr, before polyamides in 100  $\mu$ L media were added in serial dilution, in quadruplicate for each concentration. After incubation for 72 hr, cell media was replaced with 100  $\mu$ L fresh media, and cells were allowed to recover for an additional 24 hr. Cells were then fixed by adding 100  $\mu$ L of 10% trichloroacetic acid directly to each well and stored at 4°C for 1 hr, before being washed, dried, stained with 100  $\mu$ L 0.057% SRB solution per well for 30 min, and washed and dried again as described. After solubilizing the bound dye with 200  $\mu$ L of 10 mM Tris (pH 10.5) per well, absorbance at 490nm was measured on a PerkinElmer Victor microplate reader. The data are charted as a percentage of untreated controls, corrected for background absorbance. IC<sub>50</sub> is defined as the concentration that inhibits 50% of control cell growth. These values were determined by non-linear least-squares regression fit to  $Y = A + (B - A)/(1 + 10^{((\text{Log EC}_{50} - X) \times H))}$ , where  $A = \text{max.}$ ,  $B = \text{min.}$  and  $H = \text{Hill Slope}$ . All calculations were performed using Prism 4 (GraphPad) software. Stated IC<sub>50</sub> values represent the mean and standard deviation of three independent biological replicates.

## Acknowledgements


This work was supported by the National Institutes of Health (GM27681). BCL is grateful to the California Tobacco-Related Disease Research Program (18DT-0015) for a dissertation award. DCM is thankful to the National Institutes of Health for a predoctoral research training grant (5T32GM007616). The authors would like to thank Dr. Amanda E. Hargrove for helpful discussions, and Dr. Jordan L. Meier for assistance with molecular modeling.



**Figure S3.1:** Confocal microscopy analysis of 12-14 in A549 cells with included bright field images.



**Figure S3.2:** Confocal microscopy analysis of 12-14 in T47D cells with included bright field images.



S1

**Figure S3.3:** In previous unpublished results, the cytotoxicity of compound **S1** in A549 cells was examined using the same Sulforhodamine B assay described in the text. The IC<sub>50</sub> was determined to be 0.003μM ± 0.002μM.

## References

1. Dervan, P. B., Molecular recognition of DNA by small molecules. *Bioorg. Med. Chem.* **2001**, 9, 2215–2235.
2. Trauger, J. W.; Baird, E. E.; Dervan, P. B., Recognition of DNA by designed ligands at subnanomolar concentrations. *Nature* **1996**, 382, 559–561.
3. White, S.; Szewczyk, J. W.; Turner, J. M.; Baird, E. E.; Dervan, P. B., Recognition of the four Watson-Crick base pairs in the DNA minor groove by synthetic ligands. *Nature* **1998**, 391, 468–470.
4. Kielkopf, C. L.; Baird, E. E.; Dervan, P. B.; Rees, D. C., Structural basis for G·C recognition in the DNA minor groove. *Nat. Struct. Biol.* **1998**, 5, 104–109.
5. Kielkopf, C. L.; White, S.; Szewczyk, J. W.; Turner, J. M.; Baird, E. E.; Dervan, P. B.; Rees, D. C., A structural basis for recognition of A·T and T·A base pairs in the minor groove of B-DNA. *Science* **1998**, 282, 111–115.
6. Mrksich, M.; Parks, M. E.; Dervan, P. B., Hairpin peptide motif. A new class of oligopeptides for sequence-specific recognition in the minor groove of double-helical DNA. *J. Am. Chem. Soc.* **1994**, 116, 7983–7988.
7. Herman, D. M.; Baird, E. E.; Dervan, P. B., Stereochemical control of the DNA binding affinity, sequence specificity, and orientation preference of chiral hairpin polyamides in the minor groove. *J. Am. Chem. Soc.* **1998**, 120, 1382–1391.
8. Hsu, C. F.; Phillips, J. W.; Trauger, J. W.; Farkas, M. E.; Belitsky, J. M.; Heckel, A.; Olenyuk, B. Z.; Puckett, J. W.; Wang, C. C. C.; Dervan, P. B., Completion of a programmable DNA-binding small molecule library. *Tetrahedron* **2007**, 63, 6146–6151.
9. Best, T. B.; Edelson, B. S.; Nickols, N. G.; Dervan, P. B., Nuclear localization of pyrrole-imidazole polyamide-fluorescein conjugates in cell culture. *Proc. Natl. Acad. Sci. USA* **2003**, 100, 12063–12068.
10. Dervan, P. B.; Edelson, B. S., Recognition of the DNA minor groove by pyrrole-imidazole polyamides. *Curr. Opin. Struct. Biol.* **2003**, 13, 284–299.
11. Hsu, C. F.; Dervan, P. B., Quantitating the concentration of Py-Im polyamide-fluorescein conjugates in live cells. *Bioorg. Med. Chem. Lett.* **2008**, 18, 5851–5855.
12. Olenyuk, B. Z.; Zhang, G. J.; Klco, J. M.; Nickols, N. G.; Kaelin Jr., W. G.; Dervan, P. B., Inhibition of vascular endothelial growth factor with a sequence-specific hypoxia response element antagonist. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, 101, 16768–16773.
13. Kageyama, Y.; Sugiyama, H.; Ayame, H.; Iwai, A.; Fujii, Y.; Huang, L. E.; Kizaka-Kondoh, S.; Hiraoka, M.; Kihara, K., Suppression of VEGF transcription in renal cell carcinoma cells by pyrrole-imidazole hairpin polyamides targeting the hypoxia responsive element. *Acta Oncol.* **2006**, 45, 317–324.

14. Nickols, N. G.; Jacobs, C. S.; Farkas, M. E.; Dervan, P. B., Modulating hypoxia-inducible transcription by disrupting the HIF-1-DNA interface. *ACS Chem. Biol.* **2007**, 2, 561–571.
15. Nickols, N. G.; Dervan, P. B., Suppression of androgen receptor-mediated gene expression by a sequence-specific DNA-binding polyamide. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, 104, 10418–10423.
16. Matsuda, H.; Fukuda, N.; Ueno, T.; Tahira, Y.; Ayame, H.; Zhang, W.; Bando, T.; Sugiyama, H.; Saito, S.; Matsumoto, K.; Mugishima H.; Serie, K., Development of gene silencing pyrrole-imidazole polyamide targeting the TGF-beta1 promoter for treatment of progressive renal diseases. *J. Am. Soc. Nephrol.* **2006**, 17, 422-432.
17. Raskatov, J. A.; Meier, J. L.; Puckett, J. W.; Yang, F.; Ramakrishnan, P.; Dervan, P. B., Modulation of NF-kappaB-dependent gene transcription using programmable DNA minor groove binders. *Proc. Natl. Acad. Sci. USA* **2012**, 109, 1023–1028.
18. Cho, J.; Parks, M. E.; Dervan, P. B., Cyclic polyamides for recognition in the minor groove of DNA. *Proc. Natl. Acad. Sci. USA* **1995**, 92, 10389-10392.
19. Herman, D. M.; Turner, J. M.; Baird, E. E.; Dervan, P. B., Cycle polyamide motif for recognition of the minor groove of DNA. *J. Am. Chem. Soc.* **1999**, 121, 1121-1129.
20. Melander, C.; Herman, D. M.; Dervan, P. B., Discrimination of A/T sequences in the minor groove of DNA within a cyclic polyamide motif. *Chem. Eur. J.* **2000**, 6, 4487-4497.
21. Zhang, Q.; Dwyer, T. J.; Tsui, V.; Case, D. A.; Cho, J.; Dervan, P. B.; Wemmer, D. E., NMR structure of a cyclic polyamide-DNA complex. *J. Am. Chem. Soc.* **2004**, 126, 7958-7966.
22. Chenoweth, D. M.; Harki, D. A.; Phillips, J. W.; Dose, C.; Dervan, P. B., Cyclic pyrrole-imidazole polyamides targeted to the androgen response element. *J. Am. Chem. Soc.* **2009**, 131, 7182-7188.
23. Morinaga, H.; Bando, T.; Takagaki, T.; Yamamoto, M.; Hashiya, K.; Sugiyama, H., Cysteine cyclic pyrrole-imidazole polyamide for sequence-specific recognition in the DNA minor groove. *J. Am. Chem. Soc.* **2011**, 133, 18924-18930.
24. Meier, J. L.; Montgomery, D. C.; Dervan, P. B., Enhancing the cellular uptake of Py-Im polyamides through next-generation aryl turns. *Nucleic Acids Res.* **2012**, 40, 2345-2356.
25. Taleb, R. I.; Jaramillo, D.; Wheate, N. J.; Aldrich-Wright, J. R., Synthesis of DNA-sequence-selective hairpin polyamide platinum complexes. *Chem. Eur. J.* **2007**, 13, 3177-3186.
26. Van Holst, M.; Le Pevelen, D.; Aldrich-Wright, J. R., Terpyridineplatinum(II) incorporation in N-methylpyrrole-based polyamides by solid phase techniques. *Eur. J. Inorg. Chem.* **2008**, 29, 4608-4615.

27. Puckett, J. W.; Green, J. T.; Dervan, P. B., Microwave assisted synthesis of Py-Im polyamides. *Org. Lett.* **2012**, 14, 2774-2777.
28. Weltzer, M.; Wemmer, D. E., Facile dimer synthesis for DNA-binding polyamide ligands. *Org. Lett.* **2010**, 12, 3488-3490.
29. Brady, S. F.; Varga, S. L.; Freidinger, R. M.; Schwenk, D. A.; Mendlowski, M.; Holly, F. W.; Veber, D. F., Practical synthesis of cyclic peptides, with an example of dependence of cyclization yield upon linear sequence. *J. Org. Chem.* **1979**, 44, 3101-3105.
30. Boger, D. L.; Yohannes, D., Studies on the total synthesis of bouvardin and deoxybouvardin: cyclic hexapeptide cyclization studies and preparation of key partial structures. *J. Org. Chem.* **1988**, 53, 487-499.
31. Raskatov, J. A.; Hargrove, A. E.; So, A. Y. Dervan, P. B., Pharmacokinetics of Py-Im polyamides depend on architecture: cyclic versus linear. *J. Am. Chem. Soc.* **2012**, 134, 7995-7999.
32. Hargrove, A. E.; Raskatov, J. A.; Meier, J. L.; Montgomery, D. C.; Dervan, P. B., Characterization and solubilization of pyrrole-imidazole polyamide aggregates. *J. Med. Chem.* **2012**, 55, 5425-5432.
33. Synold, T. W.; Xi, B.; Wu, J.; Yen, Y.; Li, B. C.; Yang, F.; Phillips, J. W.; Nickols, N. G.; Dervan, P. B., Single-dose pharmacokinetic and toxicity analysis of pyrrole-imidazole polyamides in mice. *Cancer Chemoth. Pharm.* **2012**, 70, 617-625.
34. Baird, E. E.; Dervan, P. B., Solid phase synthesis of polyamides containing imidazole and pyrrole amino acids. *J. Am. Chem. Soc.* **1996**, 118, 6141-6146.
35. Wurtz, N. R.; Turner, J. M.; Baird, E. E.; Dervan, P. B., Fmoc solid phase synthesis of polyamides containing pyrrole and imidazole amino acids. *Org. Lett.* **2001**, 3, 1201-1203.
36. Trauger, J. W.; Dervan, P. B., Footprinting methods for analysis of pyrrole-imidazole polyamide/DNA complexes. *Methods Enzymol.* **2001**, 340, 450-466.
37. Dose, C.; Farkas, M. E.; Chenoweth, D. M.; Dervan, P. B., Next generation hairpin polyamides with (R)-3,4-diaminobutyric acid turn unit. *J. Am. Chem. Soc.* **2009**, 130, 6859-6866.
38. Farkas, M. E.; Li, B. C.; Dose, C.; Dervan, P. B., DNA sequence selectivity of hairpin polyamide turn units. *Bioorg. Med. Chem. Lett.* **2009**, 19, 3919-3923.
39. Vichai, V.; Kirtikara, K., Sulforhodamine B colorimetric assay for cytotoxicity screening *Nat. Protoc.* **2006**, 1, 1112-1116.