

Table S3

Primer sequences for quantitative real-time RT-PCR

Gene	Forward	Reverse
<i>Ski</i>	ACTACACGGGCAAGGAGGAG	TGGACTGCTTGTCTTCTCA
<i>Smad3</i>	TGTCCCAGCACACAATAAC	GTAAGTTCCACGGCTGCATT
<i>TGFBR1</i>	AGAAGAGCGTTCATGGTTCC	AATCTGACACCAACCACAGC
<i>IL7Ra</i>	TGGCTCTGGGTAGAGCTTTC	GTGGCACCAGAAGGAGTGAT
<i>Hck</i>	AGTCCAGTTTCTCCGAGAT	GCTGTTGCTGTTGTTTGGTC
<i>Bambi</i>	CGTCTCTCTCCTTCCAAGA	GGAAGTCAGCTCCTGCATCT
<i>Limd1</i>	CTCCACTGCGGAATTGAAGT	AAGCACGCATCGTGGTAGAG
<i>Psen1</i>	ACTTCCAGAATGCCAGATG	TGTTCCACCACCTGTCTTGA
<i>Psen2</i>	GGAGGATGGAGAGAGCACTG	TGCCCGTTCTTCTCAGTGTA
<i>Zfp710</i>	CACTGAGTGTGGCATGGAGT	TCTTGAGGGTTTGTCTTCTGC
<i>Fes</i>	GCAAAGTGGAGCGTGCTGAT	TACCAGGGTCCGGTACTGAG

Other primer sequences used were previously described ([David-Fung et al., 2009](#); [Li et al., 2010](#); [Yui et al., 2010](#)).