# Hydrogen Bonding Constrains Free Radical Reaction Dynamics at Serine and Threonine Residues in Peptides

(Reproduced in part with permission from Thomas, D. A.; Sohn, C. H.; Gao, J.; Beauchamp, J. L. *J. Phys. Chem. A* **2014**, *118*, 8380-8392. http://dx.doi.org/10.1021/jp501367w Copyright 2014 American Chemical Society.)

## 5.1 Abstract

Free radical-initiated peptide sequencing (FRIPS) mass spectrometry derives advantage from the introduction of highly selective low-energy dissociation pathways in target peptides. An acetyl radical, formed at the peptide N-terminus via collisional activation and subsequent dissociation of a covalently attached radical precursor, abstracts a hydrogen atom from diverse sites on the peptide, yielding sequence information through backbone cleavage as well as side-chain loss. Unique free radical-initiated dissociation pathways observed at serine and threonine residues lead to cleavage of the neighboring N-terminal  $C_{\alpha}$ -C or N- $C_{\alpha}$  bond rather than the typical  $C_{\alpha}$ -C bond cleavage observed with other amino acids. These reactions were investigated by FRIPS of model peptides of the form AARAAAXAA, where X is the amino acid of interest. In combination with density functional theory (DFT) calculations, the experiments indicate the strong influence of hydrogen bonding at serine or threonine on the observed free radical chemistry. Hydrogen bonding of the side chain hydroxyl group with a backbone carbonyl oxygen aligns the singly occupied π orbital on the β-carbon and the N- $C_{\alpha}$  bond, leading to low-barrier βcleavage of the N–C<sub>a</sub> bond. Interaction with the N-terminal carbonyl favors a hydrogenatom transfer process to yield stable *c* and *z*<sup>•</sup> ions, whereas C-terminal interaction leads to effective cleavage of the C<sub>a</sub>–C bond through rapid loss of isocyanic acid. Dissociation of the C<sub>a</sub>–C bond may also occur via water loss followed by  $\beta$ -cleavage from a nitrogencentered radical. These competitive dissociation pathways from a single residue illustrate the sensitivity of gas-phase free radical chemistry to subtle factors such as hydrogen bonding that affect the potential energy surface for these low-barrier processes.

## **5.2 Introduction**

Over the last two decades, the use of mass spectrometry experiments for the sequencing of biological molecules has been a major field of research, especially in the area of protein analysis, where high resolution and unparalleled sensitivity make tandem mass spectrometry (MS/MS) experiments the method of choice.<sup>237-241</sup> Successful dissociation of peptides or proteins in the gas phase is essential for obtaining peptide sequence information, and collision-induced dissociation (CID) and infrared multiphoton dissociation (IRMPD) remain the most widely utilized approaches for inducing backbone cleavage even after several decades of research.<sup>242-245</sup> In these techniques, input energy is statistically redistributed throughout the degrees of freedom of the peptide, resulting predominantly in the cleavage of the amide linkage and generally yielding significant sequence coverage.<sup>246</sup> However, shortcomings of these methods become apparent in the examination of peptides containing aspartic and glutamic acid or proline residues, where cleavage is highly selective, as well as in the analysis of post-translational modifications (PTMs), which are typically lost as neutral fragments in low-energy pathways.<sup>245,247</sup>

Partially in response to these limitations, the past decade has seen the development of new techniques that rely on radical-induced dissociation to provide sequence information. These methods can generally be grouped by their utilization of either hydrogen-abundant or hydrogen-deficient radicals.<sup>248</sup> Hydrogen-abundant radical techniques include electron capture and electron transfer dissociation (ECD and ETD, respectively), which mainly effect cleavage of backbone N– $C_{\alpha}$  bonds through the capture of low-energy electrons, yielding N-terminal c and C-terminal z' product ions.<sup>244,249-251</sup> The unexpected cleavage of the comparatively strong N– $C_{\alpha}$  bond has led to a large body of work investigating the mechanism of ETD and ECD,<sup>249,252-258</sup> with the Utah-Washington (UW) mechanism proposed independently by Simons and co-workers<sup>259-267</sup> and Tureček and coworkers<sup>257,268-274</sup> being the most prevalent explanation. This mechanism posits that  $N-C_{\alpha}$ bond dissociation occurs by electron attachment to an amide  $\pi^*$  orbital, either by direct electron capture or by intramolecular transfer from a high n-Rydberg state, yielding an amide radical "superbase" anion, which rapidly undergoes N–C<sub> $\alpha$ </sub> bond cleavage.<sup>255,269</sup> Though significant progress has been achieved in elucidating the mechanism of ECD and ETD, many questions about this intriguing process remain under debate, such as the initial site of electron capture or transfer and the order of N-C<sub> $\alpha$ </sub> bond cleavage and proton transfer.<sup>255,275-277</sup> The success of these techniques, however, is not disputed; they provide extensive sequence coverage through predominantly nonselective backbone cleavage with retention of PTMs.<sup>245,278-280</sup> These attributes, along with the ability to selectively cleave disulfide bonds, make ECD and ETD excellent methods for a wide range of proteomic analyses, from identification of glycosylation sites to the analysis of disulfide linkages.<sup>240,245,252,253,281,282</sup>

**Scheme 5.1.** Generation of Peptide Free Radicals by N-Terminal Derivatization and Gas-Phase Collisional Activation



Parallel to the development and implementation of the hypervalent radical techniques, many research groups investigated methods for the formation of hydrogen-deficient radical peptides. Initially, these species were generated by UV photoionization of aromatic residues or CID of copper coordination complexes to generate copper-bound peptide radicals.<sup>283</sup> More recent work has established simple techniques for the generation of hydrogen-deficient peptide radicals, including CID of an array of peptide-metal complexes,<sup>284-290</sup> peptide irradiation with 157 nm light,<sup>291-293</sup> and photodissociation of carbon-iodine bonds.<sup>294-296</sup> To achieve regioselective free radical generation via collisional

activation, Porter and co-workers converted the lysine side chain to a peroxycarbamate moiety,<sup>297,298</sup> whereas O'Hair and coworkers utilized nitrosylation and nitrate ester formation at cysteine and serine residues, respectively.<sup>299-303</sup> In addition, Hodyss and coworkers developed the technique of free-radical-initiated peptide sequencing (FRIPS), in which a well-defined radical site is produced at the N-terminus of a peptide or protein by dissociation of a covalently bound free radical precursor through a low-energy pathway.<sup>304</sup> The initial free radical precursor developed for this experiment was based on the Vazo-68 free radical initiator, which was coupled to the N-terminus via an acetyl group. As shown on the left side of Scheme 5.1, a two-step gas-phase collisional activation resulted in the generation of an acetyl radical at the N-terminus of the peptide. Later work by Lee and coworkers as well as Sohn and co-workers employed processes involving use of the stable TEMPO (2,2,6,6-tetramethylpiperidine-1-oxyl) free radical,<sup>305-308</sup> which provides the advantage of requiring only a single stage of collisional activation for radical generation, as demonstrated on the right side of Scheme 5.1. The latter methodology is employed in the current investigation.

Scheme 5.2. Mechanism of Peptide Dissociation by Hydrogen-Deficient Free Radical Chemistry



Unlike ECD and ETD, FRIPS and other hydrogen-deficient radical techniques produce primarily cleavage of  $C_{\alpha}$ -C bonds and side-chain losses.<sup>304,309,310</sup> These products are formed by the abstraction of a hydrogen atom by the initial radical center from the backbone  $C_{\alpha}$  or the side-chain  $C_{\beta}$  or  $C_{\gamma}$  position of a given amino acid residue followed by  $\beta$ -cleavage. As outlined in Scheme 5.2, the site of hydrogen atom abstraction generally determines the types of product ions formed. Hydrogen abstraction from  $C_{\alpha}$  results in the loss of an odd-electron side-chain fragment through  $\beta$ -cleavage of the  $C_{\beta}$ - $C_{\gamma}$  bond (Scheme

5.2a). The loss of an even-electron side-chain fragment results from  $C_{\gamma}$ -H abstraction through a similar process, giving a radical peptide ion (Scheme 5.2c). Backbone cleavage occurs following hydrogen abstraction from  $C_{\beta}$ , except at methionine residues, where the loss of a sulfhydryl radical is favored (Scheme 5.2b). This process is highly sensitive to amino acid side-chain chemistry, owing to subtle differences in dissociation energetics that modulate which product is favored. A study by Julian and co-workers using a noncovalently associated 18-crown-6 free radical precursor concluded that the amino acids generally fall into three main categories: (1) nonreactive residues (Ala, Gly, Pro), (2) backbone  $C_{\alpha}$ -C bond cleavage residues (Asp, Asn, His, Tyr, Phe, Val, Trp), and (3) sidechain loss residues (Lys, Glu, Gln, Arg, Met, Cys, Ile, Leu).<sup>311</sup> The amino acid residues that primarily undergo side-chain loss or are largely unreactive tend to possess high bond dissociation energies (BDEs) for the  $C_{\beta}$ -H bond, whereas those that yield backbone  $C_{\alpha}$ -C bond cleavage have comparable BDEs for the  $C_{\alpha}$ -H and  $C_{\beta}$ -H bonds. This process is largely independent of the charge state of the precursor ion, as evidenced by the generation of similar product ions from either cations or anions,<sup>311,312</sup> suggesting that barriers for these processes are typically lower than those facilitated by a mobile proton to yield b- and ytype ions. In addition, there is evidence that  $C_{\beta}$ -H BDEs are not always sufficient to predict the observed product ions, as the initial radical position and the peptide or protein conformation can strongly influence observed dissociation pathways. For example, product ions are often formed at residues in close physical proximity to the site of radical formation in the low-energy peptide or protein conformations.<sup>287,294,296,313,314</sup> Alternately, the radical site may migrate multiple times (i.e., participate in several C-H abstraction processes) before achieving a configuration with a low barrier to bond cleavage.<sup>306,315,316</sup> These studies indicate that the relative importance of competitive dissociation pathways are highly dependent on subtle factors that introduce small differences in activation energies.

The unique properties of hydrogen-deficient free radical chemistry have been shown to be especially useful for a wide range of specialized proteomics experiments, including the elucidation of phosphorylation sites,<sup>317</sup> the analysis of cysteine residues and disulfide linkages, 306,307,318 and the detection of D-amino acids. 319 The discovery of even more innovative applications of this technique is contingent upon a comprehensive knowledge of the underlying free radical reaction dynamics. To this end, the present study investigates the atypical dissociation pathways observed at serine and threonine residues in peptides. As reported by Julian and co-workers and Reilly and co-workers,<sup>311,320</sup> activation of hydrogen-deficient radical ions containing serine or threenine at the  $n^{\text{th}}$  residue from the N-terminus and  $m^{\text{th}}$  residue from the C-terminus results in the formation of  $[a_{n-1}+H]^{\bullet}$  and  $z_m$  ions, as well as  $c_{n-1}$  and  $[z_m-H]^{\bullet}$  ions. However, no compelling rationale has been proposed for the formation of these products in preference to the more commonly observed  $a_n$  and  $x_{m+1}$  ions. This work posits that the hydrogen bonding capabilities of the serie and threonine moieties lead to a number of novel dissociation pathways following hydrogen atom abstraction from either  $C_{\alpha}$  or  $C_{\beta}$ . Specifically, a simple hydrogen bonding interaction alters the relative activation energies for competitive dissociation pathways by stabilizing the transition state, either by optimizing orbital overlap to yield  $\pi$  bond formation via N–C<sub>a</sub> bond cleavage or through accessing a hydrogen atom transfer intermediate. The sensitivity of these low-barrier free radical dissociation pathways to alterations in the free energy landscape via noncovalent interactions is a key distinction between free-radical-initiated biomolecule dissociation and standard collisional activation methods.

## 5.3 Methods

#### 5.3.1 Materials

The model peptides AARAAAAAA, AARAAMAHA, AARAAASAA, AARAAATAA, AARAAASATA, AATAAARAAA, and AARAAAT(OMe)AA (where T(OMe) denotes a methylated threonine residue) were purchased as crude synthesis products (minimum 70% purity) from Biomer Technology (Pleasanton, CA, USA). HPLC grade methanol, acetonitrile, and water were purchased from EMD Merck (Gibbstown, NJ, USA). For desalting, 10  $\mu$ L pipet ZipTips with 0.6  $\mu$ L C<sub>18</sub> resin were purchased from Millipore Corp (Billerica, MA, USA). All other chemicals were purchased from Sigma-Aldrich (St. Louis, MO, USA).

#### 5.3.2 Synthesis of TEMPO-Based FRIPS Reagent

The TEMPO-based FRIPS reagent recently developed by Sohn and co-workers in this group, based upon the procedure outlined by Lee and co-workers,<sup>305</sup> was synthesized and employed for free radical generation.<sup>306</sup> Full synthesis details can be found elswhere.<sup>321</sup> Briefly, the FRIPS reagent was synthesized starting with methyl 2-bromoacetate, to which the TEMPO (2,2,6,6-tetramethylpiperidine-1-oxyl) reagent was coupled to give methyl 2- (2,2,6,6-tetramethylpiperidin-1-yloxy)acetate. This compound was then converted to 2- (2,2,6,6-tetramethylpiperidin-1-yloxy)acetic acid by stirring in 2 M potassium hydroxide in tetrahydrofuran for 24 h. The free acid was then activated by mixing with trifluoroacetic-*N*-hydroxysuccinimide ester in dry *N*,*N*-dimethylformamide for 24 h to yield 2,5-dioxopyrrolidin-1-yl 2-(2,2,6,6-tetramethylpiperidin-1-yloxy)acetate, the desired TEMPO-based FRIPS reagent.

## 5.3.3 Peptide Conjugation

Approximately 20 µg of peptide and 100 µg of the TEMPO-based FRIPS reagent were dissolved in 20 µL of a 50 mM triethylammonium bicarbonate solution (pH 8.5) in 50/50% (v/v) acetonitrile/water. The conjugation of the radical precursor to the peptide was allowed to progress for 2 h before quenching the reaction by the addition of 10 µL 25% (w/w) hydroxylamine hydrochloride in H<sub>2</sub>O to reverse any derivatization of the hydroxyl groups. The solvent was then removed with use of a centrifugal evaporator, and the peptide was resuspended in 0.1% (v/v) trifluoroacetic acid in H<sub>2</sub>O. The solution was desalted using Millipore C<sub>18</sub> ZipTips according to the manufacturer's instructions. The 5 µL elution solvent containing approximately 5 µg of peptide was brought to a final volume of 250 µL in a 50/50/0.1% (v/v) solution of methanol/water/formic acid and was electrosprayed directly into the mass spectrometer. The acquired mass spectrum demonstrated 95% or greater derivatization of the peptide with the FRIPS reagent for all peptides examined.

#### 5.3.4 Mass Spectrometry

Experiments were performed on an LCQ Deca XP quadrupole mass spectrometer (Thermo-Fisher, Waltham, MA, USA). Peptide solutions were infused directly into the electrospray source of the mass spectrometer by a syringe pump at a flow rate of 3  $\mu$ L/min. An electrospray voltage of 3.5 kV, capillary voltage of 41-42 V, capillary temperature of 275 °C, and tube lens voltage of -40 to -60 V were used. Other ion optics parameters were optimized by the autotune function in the LCQ tune program for maximizing the signal intensity. The precursor isolation window for MS<sup>*n*</sup> experiments was set to 3.0 m/z, and the normalized collisional energy in the LCQ tune program was varied from 26% to 30%

according to residual precursor ion intensities. Spectra were recorded for 50 or 100 scans based upon signal intensity.

#### **5.3.5** Computational Methods

Seven structures were chosen as simple model systems for the study of the dissociation pathways of threonine. The molecules 2-acetamido-3-hydroxy-*N*-methylbutanamide, 2acetamido-3-methoxy-*N*-methylbutanamide, and 2-acetamido-3,*N*-methylbutanamide were utilized as models for threonine, *O*-methylthreonine, and valine, respectively. These structures were composed of amino acids with an additional N-terminal acetyl group and C-terminal methylated amine and were employed to investigate the energetics of N–C<sub>a</sub> and C<sub>a</sub>–C bond cleavage with an initial radical center at C<sub>β</sub> (Figure 5.3a, b, d). The threonine and *O*-methylthreonine model structures were also optimized with a radical centered at C<sub>a</sub> to examine loss of water or methanol from the side-chain (Figure 5.3c,e, respectively). Finally, the alanine-based structures 2-acetamidopropanimide and 2-acetamido-*N*-(prop-1en-2-yl)propanamide were utilized for the examination of cleavage of the C<sub>a</sub>–C bond beginning from a nitrogen-centered radical (Figure 5.3f, g).

Initial geometries for each molecule were generated by MC/MM conformer search with OPLS 2005 as the force field using Macromol 8.0 as implemented in Maestro 8.0 (Schrödinger Inc., Portland, OR, USA) under the Linux environment. All initial structures falling within 5 kcal/mol of the lowest-energy structure were recorded and examined. Conformation space of the radical for each system was searched by substitution of the carbon radical center with boron to simulate the trigonal bonding environment. Candidate structures from these simulations were then screened manually to avoid redundancy, and

the boron atom was replaced by a carbon radical site for further optimization utilizing density functional theory (DFT) with Jaguar 7.5 (Schrödinger Inc., Portland, OR, USA) at the B3LYP/6-31+G(d) level of theory. By monitoring the occurrence of imaginary vibrational frequencies, only nontransition state structures (i.e., no imaginary vibrational frequencies) were further optimized using a higher basis set at the B3LYP/6-311++G(d,p) level.

To obtain an initial potential energy surface for each reaction, the energetics of the bond dissociation process being examined were explored by geometry optimization at fixed bond distances (i.e., relaxed coordinate scan) with a step size of 0.1 Å at the B3LYP/6-31+G(d) level of theory. After visualization in Molden,<sup>322</sup> transition state structures were determined by a standard transition state search at the B3LYP/6-311++G(d,p) level of theory from the highest-energy conformer found, and observation of the correct transition state was confirmed by IRC calculations. Generally, only the transition state for cleavage of the bond of interest was found, as well as the energies of the products. Small barriers due to rotations about a bond or noncovalent interactions were not investigated, except in the case where the transition state for bond cleavage was separated from the enthalpy of reaction by a noncovalent complex of the product ions.

The single point energy for each structure was also calculated using the M05-2X and M06-2X density functionals with the 6-311g++G(d,p) basis set. The two new-generation meta-hybrid functionals other than B3LYP were chosen for their ability to reliably predict the energetics of organic radical reactions, as well as their strong performance in assessing hydrogen bond interactions and dipole moments.<sup>323-325</sup> Thermochemical corrections (zero point energy plus thermal internal energy) were calculated with the B3LYP/6-311++G(d,p)

basis set at 298 K and applied to SCF calculations from all utilized density functionals. Charges derived from the electrostatic potential and Mulliken populations were calculated for selected molecules using Jaguar, and natural atomic charges were calculated using NBO 5.0 (Theoretical Chemistry Institute, University of Wisconsin, Madison, WI) at the B3LYP/6-311g++G(d,p) level of theory. All optimizations were performed using the spin unrestricted methodology, and the spin contamination was found to be small (S<sup>2</sup><0.782). All calculations were performed using computational resources kindly provided by the Material and Process Simulation center at the Beckman Institute, Caltech.

## 5.4 Experimental Results and Discussion

#### 5.4.1 Unique Product Ions Observed at Serine and Threonine

Peptide ions derivatized by the TEMPO-based FRIPS reagent were collisionally activated to provide precursors for studies of hydrogen-deficient radical-driven dissociation. The use of the TEMPO-based reagent greatly improves the signal intensity of FRIPS spectra over the previously employed Vazo 68 FRIPS reagent by eliminating an additional stage of MS/MS experiments (Scheme 5.1). In all experiments presented in this work, the singly protonated peptide derivatized with the TEMPO-based reagent was isolated and subjected to CID, producing the acetyl radical peptide ion (right-hand side of Scheme 5.1). Further dissociation of this species by collisional activation (i.e., MS<sup>3</sup>) produced the desired FRIPS spectrum.



**Figure 5.1.** Free radical chemistry of serine and threonine in model peptides. For both serine (a) and threonine (b), the spectra are dominated by the formation of  $[a_6+H]^{\bullet}$  and  $c_6$  ions, instead of the more typical  $a_7$  ion and side-chain losses. Reaction at threonine is more prominent than at serine, as seen in (e). Methylation of the hydroxyl group of threonine (c) prevents  $c_6$  ion formation. The FRIPS spectrum of a model peptide with the threonine and arginine residues transposed to place the charge site on the C-terminal side of threonine (d) yields the  $z_7$ ,  $[z_7-H]^{\bullet}$ , and  $x_7-H_2O$  ions.

Parts a and b of Figure 5.1 show the FRIPS spectra of AARAAASAA and AARAAATAA, respectively. Contrary to the expected dissociation to yield a ions or sidechain losses typically observed in FRIPS spectra, reaction at serine or threonine generates the  $[a_6+H]^{\bullet}$  and  $c_6$  ions as the dominant products, with minor formation of the  $a_7$  ion. These atypical product ions, whose assignment was confirmed by MS<sup>*n*</sup> experiments, cannot be explained by the general mechanisms outlined in Scheme 5.2. Also notable is a small amount of water loss and  $a_7$ –H<sub>2</sub>O ions, which have been observed previously in free radical reactions at serine and threonine.<sup>320</sup> Although the FRIPS spectra for these two model peptides are nearly identical, there is a significant difference in the abundance of CO<sub>2</sub> loss, which can be attributed to a difference in reactivity between serine and threonine. The FRIPS spectrum of AARAASATA shown in Figure 5.1e further highlights this disparity, with a significantly higher abundance of product ions from reaction at threonine than serine. These differences in reactivity were investigated using DFT calculations on model systems, and it was found that the activation energy for the rate-limiting β-cleavage reactions was systematically higher for serine than threonine due to the absence of the stabilizing methyl group on the side chain.<sup>321</sup>

To further investigate the mechanism of formation of these atypical product ions, the transposed sequence of the threonine model peptide, AATAAARAA, was examined. The FRIPS spectrum shown in Figure 5.1d is dominated by the formation of the  $z_7$  ion, with the  $[z_7-H]^{\bullet}$  (inset),  $a_7$ , and  $x_7-H_2O$  ions also of notable abundance. The  $y_6$  ion is likely formed from dissociation of the  $z_7$  species, as this ion is prominently observed upon collisional activation of the  $z_7$  ion. The  $a_7$  ion results from  $\beta$ -cleavage of the  $C_{\alpha}$ -C bond at the arginine residue, a process that gave negligible yields of the  $a_3$  ion in the model peptides possessing an arginine near the N-terminus. This difference can again be attributed to the somewhat regioselective nature of the FRIPS reagent. Similarly, the increased abundance of CO<sub>2</sub> neutral loss can be attributed to the decreased accessibility of the reactive threonine residue to the radical site. However, given that the most prominent ions in the spectrum occur from the threonine residue near the C-terminus, the chemical properties of the various moieties

on the peptide are more influential than amino acid position in determining product ion yields.

In explaining the atypical free radical chemistry of serine and threonine, attention is naturally focused on the hydroxyl group, which not only serves as a key site for modifying protein structure and function through PTMs but also has the ability to affect protein stability directly by hydrogen bonding. To determine the role this functional group plays in observed dissociation patterns, the threonine residue in the model peptide was modified by methylation of the hydroxyl group and examined using FRIPS. As shown in Figure 5.1c, the FRIPS spectrum is dominated by the loss of a methyl radical. This process likely occurs by hydrogen abstraction from  $C_{\beta}$  of the O-methylthreonine residue followed by  $\beta$ -cleavage of the O-CH<sub>3</sub> bond, resulting in the formation of a stable ketone product. Similarly, the neutral loss of the methoxy radical (CH<sub>3</sub>O<sup>•</sup>) can be explained by hydrogen abstraction from  $C_{\alpha}$  of *O*-methylthreonine and subsequent  $\beta$ -cleavage of the C–OCH<sub>3</sub> bond. Interestingly, the  $c_6$  ion is completely absent from the spectrum, implicating the necessity of the hydroxyl group for formation of this ion. In contrast, the  $[a_6+H]^{\bullet}$  ion is still present, although its relative abundance is decreased, possibly due to competition with the more favorable loss of the methyl radical. The a<sub>7</sub> ion is similarly present with decreased relative intensity.



**Figure 5.2**. Collisional activation (MS<sup>4</sup>) of the water neutral loss from model serine and threonine peptides. Each spectrum is dominated by a prominent product ion, either the  $[a_{n-1}+H]^{\bullet}$  ion or the  $x_7$  ion. All product ion identifications include loss of water from serine or threonine, except those marked with \*, and in (e), where <sup>#</sup> explicitly denotes water loss from serine.

The presence of the  $x_7$ -H<sub>2</sub>O ion in such great abundance in the FRIPS spectrum of AATAAARAA (Figure 5.1d) is rather unexpected, indicating that a favorable dissociation pathway exists for C<sub> $\alpha$ </sub>-C bond cleavage coupled with water loss, likely from the threonine side-chain. To further investigate the water loss pathway, MS<sup>4</sup> experiments were performed on the neutral loss of water from each of the model peptides, as shown in Figure 5.2. In each case, the spectrum predominantly consists of a single ion, either the [a<sub>n-1</sub>+H]<sup>•</sup> ion when an N-terminal charge site is present, or the  $x_m$  ion when there is a C-terminal charge

site. Interestingly, the MS<sup>4</sup> spectrum of AARAASATA (Figure 5.2e) shows a much greater abundance of the  $[a_5+H]^{\bullet}$  ion than the  $[a_7+H]^{\bullet}$  ion, suggesting that the neutral loss of water occurs largely from serine rather than threonine in this case. However, whether this trend is due to peptide structural constraints or the inherent differences in the chemistry of the serine and threonine side chains is not clear from these data. Overall, these results reveal multiple reaction pathways to the  $[a_{n-1}+H]^{\bullet}$  ion that generate distinct C-terminal product ions, further highlighting the complex free radical chemistry of serine and threonine.

#### 5.4.2 Proposed Mechanism of Reaction at Serine and Threonine Residues

It is clear from the experimental results that the common processes of  $C_{\alpha}$ –C bond cleavage (from a  $C_{\beta}$ -centered radical) and side-chain loss (from a  $C_{\alpha}$ -centered radical) are replaced by distinct lower-energy dissociation pathways at serine and threonine. Scheme 5.3 outlines the proposed mechanisms by which the novel product ions shown in Figure 5.1 are formed. All of these mechanisms posit that the presence of hydrogen bonds between the side-chain hydroxyl group and backbone amide group enables new dissociation pathways by accessing energetically favorable fragmentation processes. In spite of the relative weakness of the hydrogen bonds (~5 kcal/mol)<sup>326-329</sup> in comparison to the cleaved covalent bonds, these interactions are capable of significantly altering the energetic landscape by guiding access to and stabilizing transition states for otherwise noncompetitive dissociation reactions.





The  $[a_{n-1}+H]^{\bullet}$  ion is proposed to form as a result of hydrogen bonding between the Cterminal carbonyl oxygen and the hydroxyl group, forming a six-membered ring that stabilizes the reaction intermediate, as illustrated in Scheme 5.3a. Due to the favorable interactions between the hydroxyl group and the carbonyl oxygen, the energy required to cleave the N–C<sub>a</sub> bond is lowered, leading to preferential β-cleavage. This process gives the  $z_m$  ion and a  $c_{n-1}^{\bullet}$  ion that rapidly loses isocyanic acid to give the observed  $[a_{n-1}+H]^{\bullet}$  ion. The dominance of the  $[a_{n-1}+H]^{\bullet}$  (Figure 5.1a,b) and  $z_m$  ions (Figure 5.1d) in the FRIPS spectra suggest that this pathway should be the most energetically favored. The hydroxyl group of serine or threonine may alternately form a seven-membered ring through hydrogen bonding with the N-terminal carbonyl oxygen atom, as shown in Scheme 5.3b.

This conformation facilitates hydrogen atom transfer from the hydroxyl group in conjunction with hydrogen abstraction from  $C_{\beta}$ . As the N– $C_{\alpha}$  bond is elongated, the transfer of the hydrogen atom occurs in a concerted manner, reducing the reaction barrier and yielding an enol-imine  $c_{n-1}$  ion and the  $C_{\alpha}$ -centered radical  $[z_m-H]^{\bullet}$  ion. This hydrogen atom transfer process is supported by the experimental observation that methylation of the threonine hydroxyl group eliminates formation of the  $c_{n-1}$  ion. Both of these mechanisms are in agreement with those proposed by Sun et al. for the formation of  $c_{n-1}$  and radical  $a_{n-1}$  ions at serine and threonine residues through hydrogen-deficient radical dissociation from a crown ether-based photolabile radical precursor, but the importance of the hydrogen bonding interactions has not been previously demonstrated.<sup>311</sup>

As evidenced by the formation of  $[a_{n-1}+H]^*$  ions in the absence of a hydrogen bond donor (Figure 5.1c) as well as the formation of  $[a_{n-1}+H]^*$  and  $x_m$ –H<sub>2</sub>O ions following neutral loss of water (Figure 5.2), a third dissociation pathway must be considered at serine and threonine residues. In the mechanism proposed in Scheme 5.3c, the hydroxyl group of serine or threonine acts as hydrogen bond acceptor instead of donor, leading to a strong interaction with the N-terminal amide hydrogen. Following hydrogen abstraction from  $C_{\alpha}$ by the acetyl radical, the new radical center may undergo  $\beta$ -cleavage in concert with hydrogen abstraction from the backbone amide nitrogen, yielding a neutral loss of water and a nitrogen-centered radical. This nitrogen-centered radical may then undergo further  $\beta$ -cleavage of the N-terminal  $C_{\alpha}$ –C bond to yield the observed  $[a_{n-1}+H]^*$  and  $x_m$ –H<sub>2</sub>O ions. A similar mechanism was previously proposed by Reilly and co-workers in their examination of the collisional activation of radical a ions produced by 157 nm photodissociation.<sup>320</sup> Importantly, this mechanism offers an explanation for the formation of the  $[a_{n-1}+H]^{\bullet}$  ion even when the hydroxyl group of threonine is methylated (Figure 5.1c), as the methoxy group may still act as hydrogen bond acceptor and yield the nitrogencentered radical via loss of methanol.

## 5.5 Computational Analysis

## 5.5.1 Models for Serine and Threonine Free Radical Chemistry

Given the strong experimental evidence for the mechanisms proposed in Scheme 5.3, in both this work and previous studies,<sup>311,320</sup> we investigate these reactions utilizing computational techniques to better understand the free radical chemistry of serine and threonine. The C<sub>β</sub>-centered radical structures shown in Figure 5.3a,b,d were employed to examine cleavage of the N–C<sub>a</sub> and C<sub>a</sub>–C bond via β-cleavage, and the C<sub>a</sub>-centered radicals in Figure 5.3c,e were used to explore the neutral loss of water from the side-chain. Finally, the nitrogen-centered radicals shown in Figures 5.3f,g were used to study β-cleavage of the C<sub>a</sub>–C bond. The structures shown beneath the schematic diagram for each molecule represent the low-energy conformer or conformers found for that compound, with the relative enthalpy (B3LYP/6-311++G(d,p) level of theory) indicated for molecules with multiple low-energy conformations of interest.



**Figure 5.3**. Model structures for computational investigation of threonine free radical chemistry. The structures of compounds possessing a  $C_{\beta}$ -centered radical are shown for valine (a), threonine (b), and *O*-methylthreonine (d), whereas those possessing a  $C_{\alpha}$ -centered radical are shown for threonine (c) and *O*-methylthreonine (e). The final two compounds (f and g) were utilized to study  $C_{\alpha}$ -C bond cleavage from a nitrogen-centered radical. Relative enthalpies [B3LYP/6-311++G(d,p)] are indicated for multiple structures. Hydrogen bonding interactions are denoted with dashed lines (see text for further discussion).

For all bond cleavage processes examined, the reaction energetics were calculated using the M05-2X and M06-2X density functionals as well as B3LYP. These functionals give reaction enthalpies differing by as much as 13 kcal/mol for  $\beta$ -cleavage processes, with slightly smaller differences for activation energies. Previous work by Izgorodina et al. demonstrated that B3LYP may underestimate the energetics of  $\beta$ -cleavage reactions by up to 9 kcal/mol.<sup>330</sup> The M05-2X and M06-2X density functionals are expected to be significantly more accurate for these reactions, although they were found to overestimate energetics of  $\beta$ -cleavage by ~2 kcal/mol compared to G3(MP2)-RAD benchmarks.<sup>325</sup> These results suggests that the computational energetics presented here likely represent upper and lower bounds on the actual thermochemistry.

#### 5.5.2 $\beta$ -Cleavage of the N-C<sub> $\alpha$ </sub> vs C<sub> $\alpha$ </sub>-C Bond

To investigate the favored  $\beta$ -cleavage of the N–C<sub>a</sub> bond over the C<sub>a</sub>–C bond from a radical centered at C<sub>β</sub> (Scheme 5.3a,b), the energetics for these processes were calculated for valine, serine, and threonine model compounds (Figure 5.4). The energetics for the valine model compound shown in Figure 5.4a provide reference dissociation energetics in the absence of hydrogen bonding. The low-energy conformer (**1**) has the backbone in the all-trans configuration with the side-chain oriented perpendicular to the backbone. Cleavage of the C<sub>a</sub>–C bond was found to be favored over N–C<sub>a</sub> bond cleavage by 6–10 kcal/mol depending on the density functional. Dissociation of the N–C<sub>a</sub> bond proceeds through a transition state in which a stabilizing hydrogen bond interaction between products is preserved, yielding an energetic minimum between dissociation transition state and overall reaction enthalpy of the separated products (right-hand side of Figure 5.4a).



**Figure 5.4**.  $\beta$ -cleavage of the N–C<sub> $\alpha$ </sub> vs C<sub> $\alpha$ </sub>–C bond from a C<sub> $\beta$ </sub>-centered radical on model amino acids. For the value model compound (a), dissociation of the C<sub> $\alpha$ </sub>–C bond is favored, whereas hydrogen bonding in threonine (b and c) favors N–C<sub> $\alpha$ </sub> bond cleavage via stabilization of the reaction intermediate. Enthalpies are given relative to structure **1** for the value molecule and structure **2** for the threonine molecule.

Two low-energy structures were found for the threonine model compound possessing a radical center at  $C_{\beta}$  (Figure 5.3b), and both conformers were examined to obtain reaction energetics as shown in Figure 5.4b,c. These two structures possess an all-trans backbone conformation, and the side-chain hydroxyl group forms a hydrogen bond with either the Cterminal (2, 0.0 kcal/mol) or N-terminal (3, 0.8 kcal/mol) carbonyl oxygen. Although the energetics of  $C_{\alpha}$ -C bond cleavage are quite similar between value (Figure 5.4a) and the C-terminal hydrogen bonding structure of threonine (Figure 5.4b), the energy required for  $N-C_{\alpha}$  bond cleavage is greatly decreased for threonine. The hydrogen bonding interactions are eliminated at the transition state for  $C_{\alpha}$ -C bond cleavage (20) to enhance the  $\pi$ -orbital interactions of the N-terminal product (22), but the hydrogen bond persists in the transition state for N–C<sub> $\alpha$ </sub> bond dissociation (17), reducing the energy required to form the c<sup>•</sup> (18) and z (19) ion analogues of the products proposed in Scheme 5.3a. Similar to the case for the valine model compound, the loose transition state for N– $C_{\alpha}$  bond cleavage is found to be lower than the product enthalpies due to formation of a stabilizing hydrogen bond following dissociation. Though the B3LYP calculations clearly favor cleavage of the N-C<sub>a</sub> bond over the  $C_{\alpha}$ -C bond, the M05-2X and M06-2X calculations give similar reaction enthalpies for the two processes. Experimentally, product ions resulting from N–C<sub> $\alpha$ </sub> bond cleavage are on the order of 10 times more abundant than those from  $C_{\alpha}$ -C bond cleavage, meaning a significant difference in the reaction rates would be expected on the ion trap time scale. Previous work has demonstrated that even small differences in activation energy can have large impacts on the kinetics and branching ratios of peptide dissociation,<sup>331,332</sup> and dissociation of the N– $C_{\alpha}$  bond need only be slightly enthalpically favored to be the dominant product ion (assuming similar frequency factors for the two processes).

For the N-terminal hydrogen bonding structure (3) shown in Figure 5.4c, the preference for dissociation of the N–C<sub> $\alpha$ </sub> bond over the C<sub> $\alpha$ </sub>–C bond is quite clear. In this conformation, the energy required for cleavage of the N– $C_{\alpha}$  bond is significantly reduced due to the approach of the hydroxyl and carbonyl groups at the transition state (23) and subsequent hydrogen atom transfer to give the enol-imine (24) and carbon-centered radical (25) products (c and [z-H]<sup>•</sup> ion analogues of Scheme 5.3b, respectively). The reaction coordinate for  $C_{\alpha}$ -C bond cleavage shows that the threonine side-chain rotates during dissociation, disrupting hydrogen bonding at the transition state (26) and forming the Nterminal product (27) with the hydroxyl group oriented outward along the axis of the N- $C_{\alpha}$ bond. This N-terminal product (a ion analogue) may readily isomerize to give the lowerenergy structure seen in Figure 5.4b (21) and thereby reduce the reaction enthalpy, but the process still remains unfavorable due to the significant activation energy required to reach the transition state (26). Similar energetics with slightly higher barriers for all processes were found for a model serine compound. Thus, the calculations outlined support both the data and proposed mechanism in concluding that hydrogen bond constraints favor  $N-C_{\alpha}$ bond cleavage at serine and threonine residues.

#### 5.5.3 Nitrogen-Centered Radical Chemistry via Loss of H<sub>2</sub>O

The processes examined above fail to explain the presence of  $x_m$ -H<sub>2</sub>O and  $a_n$ -H<sub>2</sub>O ions. Therefore, calculations were also utilized to investigate the energetics of the mechanism shown in Scheme 5.3c, in which dissociation occurs through  $\beta$ -cleavage from a nitrogencentered radical initially formed by water loss. The low-energy structures of the C<sub> $\alpha$ </sub>centered radical for the threonine and *O*-methylthreonine model compounds are shown in Figure 5.3c,e, respectively. For both model compounds, the low-energy structures (**4** and **7**) do not possess interactions between the side-chain hydroxyl and the N-terminal amide nitrogen that would give a clear pathway to the neutral loss of water, though such structures can be found for threonine (**5**) and *O*-methylthreonine (**8**) at relative enthalpies of 2.9 and 3.6 kcal/mol, respectively. In these conformations, the side-chain oxygen acts as hydrogen bond acceptor, with the backbone N–H group functioning as hydrogen bond donor. Water loss is found to occur from these structures via concerted cleavage of the side-chain  $C_{\beta}$ –OH bond and abstraction of hydrogen from the amide nitrogen, as shown in Figure 5.5a,b. This process occurs via a tight transition state of 32-36 kcal/mol for both molecules, although the overall enthalpy of reaction in each case was favorable (10-20 kcal/mol).

Beginning from either threonine or *O*-methylthreonine, loss of water or methanol yields a nitrogen-centered radical and double bond between  $C_{\alpha}$  and  $C_{\beta}$  (structures **30** and **32**, respectively). The structure shown in Figure 5.3g seeks to replicate these features in a model system to study the energetics of  $\beta$ -cleavage of the  $C_{\alpha}$ -C bond. As illustrated in Figure 5.5c, cleavage of the  $C_{\alpha}$ -C bond from the low-energy structure (**10**) was found to occur with a tight transition state but favorable reaction enthalpy as the N-C-O bond angle changes from 123° to 174° during formation of the double bond.



**Figure 5.5.** Energetics of alternate pathways to cleavage of the N-terminal  $C_{\alpha}$ -C bond via formation of a nitrogen-centered radical: (a) water loss from threonine model compound; (b) methanol loss from *O*-methylthreonine model compound; (c) cleavage of  $C_{\alpha}$ -C bond from nitrogen-centered radical. These processes have low reaction enthalpies but possess high activation energies.

This pathway to dissociation of the  $C_{\alpha}$ –C bond is essential to explain the experimental results for the *O*-methylthreonine model peptide (Figure 5.1c). Methylation of the hydroxyl group prevents any conformation in which the side-chain acts as hydrogen bond donor, thereby suggesting that the two mechanisms presented in Scheme 5.3a,b are inoperable for this molecule. In agreement with this assertion, it was found that cleavage of the N–C<sub> $\alpha$ </sub> bond via  $\beta$ -cleavage from the low-energy C<sub> $\beta$ </sub>-centered radical structure (**6**, Figure 5.3d) is not energetically favorable. However,  $\beta$ -cleavage from a nitrogen-centered radical in *O*-methylthreonine residues gives a viable route to the observed [ $a_{n-1}$ +H]<sup>•</sup> ion even in the absence of a side-chain hydrogen bond donor. In addition, the energetics for this process are only slightly higher than those calculated for C<sub> $\alpha$ </sub>–C bond cleavage, further supporting the feasibility of this pathway.

In the case of the threonine model compound possessing a free hydroxyl group, there exists a significant disparity between the experimental observation of the water loss pathway outlined in Scheme 5.3c and the theoretical activation energies for this reaction. The theoretical barrier to water loss of 32-36 kcal/mol is 7-14 kcal/mol higher than those calculated for the processes outlined in Scheme 5.3a,b and therefore would not be expected to compete. Since the experimental data provide compelling evidence for the proposed reaction pathway leading to water loss, especially the MS<sup>4</sup> spectra shown in Figure 5.2, it appears that these calculations overestimate the activation energy for this pathway. A conformation with a lower barrier to water loss may exist, but extensive conformational searching failed to yield a transition state lower in energy than the one shown in Figure 5.5a. Alternately, the activation energy for this process may be lowered by cleavage of the backbone  $C_{\alpha}$ -C bond in concert with hydrogen atom transfer from the amide to the

hydroxyl group, a process somewhat analogous to that proposed in Scheme 5.3b. A concerted reaction would also explain the low abundance of water loss in the FRIPS spectra, but such processes were not examined using DFT calculations due to the difficulty in searching the expanded conformational space to identify the minimum energy pathway.

## 5.5.4 Why Hydrogen Bonding Favors N-C<sub>a</sub> Bond Cleavage

Although calculations can quantify the energetics of the proposed processes leading to cleavage of the stronger N–C<sub> $\alpha$ </sub> bond in the mechanisms shown in Scheme 5.3a,b, it is useful to provide a simpler but more transposable rationalization for why these reactions occur by examining the molecular orbitals involved. Strong orbital overlap connecting reactants to products is generally associated with favorable chemical reactions, whereas little overlap leads to energetically unfavorable processes.<sup>333</sup> Parts a and b of Figure 5.6 show schematically the interaction between the singly occupied  $\pi$  orbital of the C<sub> $\beta$ </sub>-radical site and the N–C<sub> $\alpha$ </sub>  $\sigma$  bond for the N-terminal hydrogen bonding and C-terminal hydrogen bonding complexes, respectively. In either structure, the hydrogen bonding interactions strongly align the  $\pi$  orbital on C<sub> $\beta$ </sub> with the N–C<sub> $\alpha$ </sub>  $\sigma$  bond, resulting in facile cleavage of the N–C<sub> $\alpha$ </sub> bond. In contrast, the C<sub> $\alpha$ </sub>–C  $\sigma$  bond lies nearly orthogonal to the  $\pi$  orbital on C<sub> $\beta$ </sub> in both conformations, and the cleavage of this bond is therefore unfavorable without the destruction of the stabilizing hydrogen bonding interactions.



**Figure 5.6**. Alignment of the singly occupied  $\pi$  orbital on  $C_{\beta}$  with the N– $C_{\alpha}$   $\sigma$  bond via hydrogen bonding interactions of threonine with either the N-terminal (a) or C-terminal (b) carbonyl oxygen. Left: schematic representation of computational structures, with orbital alignment denoted by the dashed line, Center: computational structures with schematic representation of the singly occupied  $\pi$  orbital on  $C_{\beta}$ , Right: Newman projection along the axis of the  $C_{\beta}$ – $C_{\alpha}$  bond.

In addition to orbital overlap, the transition state structures for both conformations are stabilized by enhanced hydrogen bonding. This observation is illustrated in Scheme 5.4, which gives the bond lengths in Angstroms for the N-terminal (a) and C-terminal (b) hydrogen bonding conformations in both the ground state (GS) and transition state (TS). In the C-terminal hydrogen bonding structure, the distance between the carbonyl oxygen and the hydroxyl group hydrogen decreases from 1.78 Å in the ground state to 1.63 Å in the transition state, imparting a stabilizing partial double bond character to the hydroxyl group. In the case of the N-terminal hydrogen bonding conformer, the hydrogen atom approaches to a distance of only 1.4 Å from the carbonyl at the transition state for N–C<sub>a</sub> bond cleavage, forming a stabilizing intermediate with increased bond order on the side-

chain  $C_{\beta}$ –O bond. These hydrogen bonding interactions therefore play a crucial role in

dictating the favored pathway of the reaction.

**Scheme 5.4.** Calculated Bond Lengths (Å) during N–C<sub> $\alpha$ </sub> Bond Cleavage for the Threonine Model Compound Possessing an (a) N-Terminal or (b) C-Terminal Hydrogen Bond



### 5.5.5 Comparison of FRIPS to ECD/ETD

The process of hydrogen atom transfer along with N–C<sub>a</sub> bond cleavage (Scheme 5.3b) examined in this work gives product ions with remarkable resemblance to those observed in ECD and ETD. In the UW mechanism, which is supported by numerous experimental and theoretical reports, an aminoketyl intermediate may be formed either through direct electron capture to the amide  $\pi^*$  orbital or through intramolecular electron transfer from electron capture in a Rydberg orbital at a positively charged site, followed by proton abstraction by the strongly basic amide anion radical.<sup>255,265-267,275</sup> The mechanism proposed here is more comparable to the Cornell mechanism, in which hydrogen atom transfer is suggested to form the aminoketyl intermediate, with backbone cleavage occurring

separately.<sup>255,268,269,273</sup> However, the proposed mechanism for radical-initiated cleavage at serine and threonine is absent of any aminoketyl intermediate, as the transfer of the hydrogen atom occurs in concert with N–C $_{\alpha}$  bond cleavage. Further examination of the potential energy surface failed to find a pathway in which aminoketyl intermediate formation was more energetically favorable than concerted hydrogen atom transfer and N–C $_{\alpha}$  bond cleavage, making this dissociation mechanism unlikely for this system. In addition, the Cornell mechanism is unlikely to be prevalent in ECD and ETD, as the energetic barriers to hydrogen atom transfer are greater than those to intramolecular electron transfer followed by proton abstraction.<sup>255,260</sup> Although not directly applicable to the ECD or ETD case, the hydrogen atom transfer observed here is interesting in that it illuminates yet another unique mechanism for the cleavage of backbone N–C $_{\alpha}$  bonds via a decrease in the energetic requirements for this process.

## 5.6 Conclusions

The calculations and experiments detailed in this work provide a comprehensive analysis of the factors responsible for the unique free radical chemistry of serine and threonine. Cleavage of the N–C<sub> $\alpha$ </sub> bond may occur directly via  $\beta$ -cleavage from a C<sub> $\beta$ </sub>-centered radical, whereas dissociation of the N-terminal C<sub> $\alpha$ </sub>–C bond may occur either from a nitrogencentered radical or by loss of isocyanic acid following N–C<sub> $\alpha$ </sub> bond cleavage. As a result of the lower energetic barriers associated with odd-electron versus even-electron cleavage processes, these reaction pathways are more susceptible to influence by weak interactions such as hydrogen bonding that constrain molecular geometry in reactants, intermediates, and products. This principle has significant implications for gas-phase free radical sequencing applications, in which many peptides and proteins may possess stable gasphase noncovalent interactions that may direct radical-induced dissociation, an attribute that Julian and co-workers have leveraged to elucidate the gas phase structure of proteins.<sup>294,296,313</sup> In other situations, such effects may be masked by rapid migration of the radical site following formation.<sup>315</sup> Nevertheless, it is abundantly clear from this work that free radical chemistry can be highly sensitive to local conformation at the radical site, as no fewer than five unique dissociation pathways can occur at serine and threonine residues via dissociation from different conformations.